

REV. 3-90-B

SPECIMEN LABEL



ACCEPTED  
U.S. Environmental Protection Agency  
Reregistration Act, as amended for the  
pesticide registered under  
EPA Reg. No. 56077-37

# PROPANIL WDG

## Herbicide

**For Postemergence Control of Broadleaf and Grass Weeds in Rice Fields.**

**ACTIVE INGREDIENTS:**

3', 4' Dichloropropionanilide	50.00
Inert Ingredients:	50.00
TOTAL	100.00

This product contains 0.5 lb. of 3', 4' Dichloropropionanilide (Propanil) per pound of formulated product.

EPA Reg. No. 56077-37

EPA Est. No.

PRECAUTION AL USUARIO: SI USTED NO LEE INGLES. NO USE ESTE PRODUCTO HASTA QUE LA ETIQUETA LE HAYA SIDO EXPLICADA AMPLIAMENTE.

**KEEP OUT OF REACH OF CHILDREN**

### CAUTION-CAUCION

#### STATEMENT OF PRACTICAL TREATMENT

**If Swallowed:** Induce vomiting immediately by drinking 1 or 2 glasses of water and sticking finger down throat. Get medical attention.

**If on Skin:** Wash contaminated areas thoroughly with soap and water. Do not reuse contaminated clothing until washed. Get medical attention if irritation persists.

**If yes:** Flush with copious amounts of clean water for 10-15 minutes. Get medical attention.

### PRECAUTIONARY STATEMENTS CAUTION

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed, absorbed through the skin or inhaled. Causes eye irritation. Avoid breathing spray mist or dust. Avoid contact with skin, eyes, and clothing. Wash thoroughly with soap and water after handling. In case of contact, immediately flush eyes or skin with plenty of water. Remove contaminated clothing and wash before reuse. Get medical attention if irritation persists. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

**ENVIRONMENTAL HAZARDS:** Do not apply directly to water except as indicated on this label. Do not contaminate water when disposing of equipment washwaters. This product is toxic to fish. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water intended for irrigation or domestic purposes. Do not apply when weather conditions favor drift from target areas.

**FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE CALL TOLL FREE  
1-800-424-9300**

**AGRICULTURAL CHEMICAL**  
DO NOT SHIP OR STORE WITH FOODS, DRUGS, OR CLOTHING

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with this labeling.

**REENTRY STATEMENT:** Do not enter treated areas without protective clothing until sprays have dried. Protective clothing should include: hat or other suitable head covering, long sleeved shirt and long legged trousers, or a coverall type garment, shoes and socks.

Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. The front panel PRECAUTIONARY STATEMENTS should be read to workers as well as the instruction not to enter until sprays have dried. When oral warnings are given, warnings shall be given in language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information:

**CAUTION:** Area treated with PROPANIL on (date of application). Do not enter without appropriate protective clothing until sprays have dried. (Insert here Statements of Practical Treatment as on front panel.)

## STORAGE AND DISPOSAL

**STORAGE:** Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not store this product near fertilizers, seeds, insecticides, or fungicides. Containers should not be stacked more than 3 containers high. Reclose all partially used containers by thoroughly tightening screw cap. Damaged or leaking containers which contain product that cannot be used immediately should be transferred to suitable sound containers and properly marked. Any spilled material should be thoroughly swept up and transferred to the new container or disposed of as indicated under Pesticide Disposal. For safety and prevention of unauthorized use all pesticides should be stored in locked facilities. To prevent accidental misuse, different pesticides should be stored in separate areas with enough distance between to provide clear identification. Opened, partially used pesticides should be stored in original containers when possible. Keep containers closed when not in use.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent), adding rinsate to spray tank. Offer rinsed containers for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, if allowed by state and local authorities or by burning. If burned, stay out of smoke.

## WHERE TO USE

CEDAR CHEMICAL'S PROPANIL WDG is used for postemergence control of broadleaf and grass weeds in RICE fields.

### WEEDS CONTROLLED

Barnyardgrass (watergrass), brachians, coffeeweed, crabgrass, croton, curly indigo, foxtail, goosegrass, gulf cockspur, mexicanweed, millet (Texas), paragrass, pigweed, sourdock, spearhead, wiregrass.

(PROPANIL WDG will not control arrowhead, bermudagrass, cattail, ducksalad, johnsongrass, nutgrass, red rice and sprangletop.)

## GENERAL INFORMATION

Several important factors should be taken into account to achieve a high efficiency of selective weed control with PROPANIL WDG. These include uniform application, growth stage and weather conditions. To assure uniform application mix the prescribed amount of PROPANIL WDG with a sufficient volume of water to provide thorough coverage of target area. For aerial application use approximately 10 gallons of water, or for surface (ground) applications 20-30 gallons of water per acre at sufficient spray pressure. Agitate tank mixes thoroughly and continuously. Avoid over and under application.

Growth stage of weeds is very important. Best results for selective weed control are obtained when most grasses have reached the 1 to 3 leaf stage. Proper field preparation is essential to ascertain a relatively clod free and level

212  
surface and to obtain uniform d levels and growth. Fields may be flushed prior to treatment to produce uniform and vigorous grass germination and growth. Drain water from fields prior to applying PROPANIL WDG. Higher rates are recommended to control larger grasses or exposed weeds when rice fields are not completely drained. Inspect rice fields regularly to select the correct application time.

DO NOT apply this product through any type of irrigation system.

Do not apply this product (directly or indirectly) to any crop except rice.

Do not apply when wind conditions will allow drift to adjacent, susceptible crops such as beans, soybeans, cotton, safflower, cucurbits, vegetables, orchards (such as almonds, prunes and grapes) and other sensitive crops.

Weather conditions must be observed closely. Under cool weather conditions higher rates are required to achieve satisfactory control. Avoid application if rain threatens with 6 to 8 hours, or if wind velocities are high enough to cause drift and irregular spray patterns.

When Propanil WDG is used alone (not in combination with any other postemergent rice herbicide), a suitable non-ionic surfactant or crop oil may be used to improve wetting of foliage and weed control. Refer to the surfactant or crop oil label for proper use rates.

Consult Extension Service for detailed application advice.

## RECOMMENDED BROADCAST RATE

Apply 6 pounds of PROPANIL WDG per acre when most grasses have reached 1 to 3 leaf stage. Use 8 to 12 pounds of PROPANIL WDG per acre when the grasses are large (4 to 6 leaf stage) or when unseasonably cool weather conditions prevail or in cases where rice fields have not been drained completely and where weeds are large.

Barnyardgrass may be controlled up to 30 to 45 days after planting, before rice plants have reached the fully tillered growth stage.

**NOTE:** PROPANIL WDG applied to rice after the 4 leaf stage may cause visible injury under some climatic conditions. Rice plants usually outgrow such injury.

**NOTE:** Water drained from treated rice fields must not be used to irrigate other crops or be released within 1/2 mile upstream of a potable water intake in flowing water (i.e. river, stream, etc) or within 1/2 mile of a potable water intake in a standing body of water such as a lake, pond or reservoir.

**IN CALIFORNIA—Use PROPANIL WDG only where rice fields are not completely drained. If higher water level is desired, relood field after 12 hours and before 7 days after treatment. This will discourage new weed infestations. Do not apply PROPANIL WDG within 14 days before or after insecticide applications. Serious injury to rice may occur.**

## CONDITIONS OF SALE AND WARRANTY

CEDAR AND SELLER OFFER THIS PRODUCT AND THE BUYER AND USER ACCEPTS THIS PRODUCT UNDER THE FOLLOWING AGREED CONDITIONS OF SALE AND WARRANTY.

The directions for use of this product are believed to be reliable and should be followed carefully. However, it is impossible to take into account all variables and to eliminate all risks associated with its use. Injury or damage may result because of conditions which are beyond the control of Cedar or the Seller. Cedar warrants only that this product conforms to the chemical description on the label and is believed to be reasonably fit for the purposes referred to in the Directions for Use when used as directed under normal conditions. CEDAR MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. In no case shall Cedar or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. Any variation or exception from this warranty must be in writing and signed by an authorized Cedar representative.

**CEDAR CHEMICAL CORPORATION**  
5100 POPLAR, 24th FLOOR  
MEMPHIS, TENNESSEE 38137, USA