

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS & DOMESTIC ANIMALS

CAUTION

May cause irritation of eyes, nose, throat and skin. Avoid breathing dust. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling.

ENVIRONMENTAL HAZARD

Do not apply directly to any body of water. Do not contaminate water by cleaning of equipment or disposal of waste.

DIRECTIONS FOR USE

It is a violation of federal laws to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE

To maximize the shelf life of this product, store in dry area where the container is not likely to come in contact with moisture.

DISPOSAL

Do not contaminate water by cleaning of equipment or disposal.

Wastes resulting from this product may be disposed of on site or at an approved waste disposal facility.

(For Plastic Buckets) Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(For Bags) Completely empty bag into application equipment. Then dispose of bag in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

SPRAKIL SK-13 GRANULAR WEED KILLER

ACTIVE INGREDIENTS

Tenuthiaron 1 (5-tert-butyl-1,3,4-trimethyl-2,4,6-triazine-5-thione) 100.00% 100.00%

Diuron 3 (3,4-dichlorophenyl 1,1-dimethylurea) 5.00% 5.00%

INERT INGREDIENTS

Total 100.00% 100.00%

KEEP OFF AND AWAY FROM CHILDREN

CAUTION!

STATEMENT OF PRACTICAL TREATMENT

If in eyes: Flush with plenty of water. Get medical attention if irritation persists.

If on skin: Wash with plenty of water. Get medical attention if irritation persists.

See side panel for additional precautionary statements.

MANUFACTURED FOR

SSI INDUSTRIES, INC.
A SUBSIDIARY OF MOBLEY INDUSTRIES, INC.

1000 E. FREDERICK ROAD

HUNTINGTON, W. VA. 25701

EPA Registration No. 34473-1

EPA Establishment No. 68064-1

NET CONTENT: 50 POUNDS

NON-CROP WEED CONTROL INDUSTRIAL USE ONLY

Sprakil SK-13 Granular is a fast selective herbicide for controlling a wide range of annual and perennial weeds and grasses. It is recommended only for non-cropland areas such as railroad right-of-way and industrial areas. 150 to 400 pound applications of Sprakil SK-13 Granular per acre usually result in a nonproductive condition of the soil for a period of a year or more. The duration of non-productivity is dependent upon rainfall, soil type and other conditions.

Sprakil SK-13 Granular should be applied as furnished with a seed spreader, a fertilizer spreader, a shaker type applicator or with any equipment which will distribute the chemical uniformly over the area to be treated. Application equipment is available from SSI Industries, Inc. For the control of annual and most perennial weeds and grasses, apply Sprakil SK-13 Granular at the rate of 150 to 400 pounds per acre, on smaller plots, 0.35 to 1.0 pounds per 100 square feet. Repeat spot treatment may be required when deep rooted perennial weeds are present. To obtain the best results, Sprakil SK-13 Granular should be applied to the ground where it will be absorbed by the roots. Applications made early in the season have been found to give results superior to applications at a later seasonal date. Sprakil SK-13 Granular may, however, be applied at any time of the year. Maximum effectiveness in any region is obtained when applications are made just prior to the rainy season. Users should consult state agricultural experimental stations or extension service weed specialists for recommendations as to use in their particular area. Write to SSI Industries, Inc.

IMPORTANT: Do not apply on or near valuable woody or herbaceous plants in areas where their roots may extend because of possible injury to such plants. Thoroughly clean spreading equipment with a suitable chemical cleaner before using for other purposes (do not use same spreading equipment for other purposes). Do not contaminate irrigation ditches or water used for domestic purposes. Do not use on croplands or any land to be used for subsequent cropping, trees, animals, or treated areas. Do not re-use this container (even when empty). Do not store in areas where it might be used as fungicides.

ACCEPTED

JUL 25 1986

U.S. DEPARTMENT OF AGRICULTURE
EPA REG. NO. 34473-1

for Postemergence Control of Broadleaf and Grass Weeds in Rice Fields.

Active Ingredient:

3',4'Dichloropropionanilide	42.12
Inert Ingredients:	57.88
TOTAL	100.00

This product contains 4 lbs. of 3',4' Dichloropropionanilide (Propanil) per pound of formulated product

EPA Reg. No. 56077-Y

EPA Est. No. 42761-MS-1

PRECAUTION AL USUARIO: SI USTED NO LEE INGLES. NO USE ESTE PRODUCTO HASTA QUE LA ETIQUETA LE HAYA SIDO EXPLICADA AMPLIAMENTE

KEEP OUT OF REACH OF CHILDREN



CAUTION-CAUTION

STATEMENT OF PRACTICAL TREATMENT:

If Swallowed: Induce vomiting immediately by drinking 1 or 2 glasses of water and sticking finger down throat. Get medical attention.

If on Skin: Wash contaminated areas thoroughly with soap and water. Do not reuse contaminated clothing until washed. Get medical attention if irritation persists.

If in Eyes: Flush with copious amounts of clean water for 10-15 minutes;...Get medical attention.



2

ENVIRONMENTAL HAZARDS: Do not apply directly to water except as indicated on this label. Do not contaminate water when disposing of equipment washwaters. This product is toxic to fish. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water intended for irrigation or domestic purposes. Do not apply when weather conditions favor drift from target area.

DO NOT SHIP OR STORE WITH FOOD, FEEDS,
DRUGS, OR CLOTHING.

FOR CHEMICAL SPILL, LEAK, FIRE OR
EXPOSURE, CALL TOLL FREE 1-800-424-9300

3

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with this labeling.

REENTRY STATEMENT: Do not enter treated areas without protective clothing until sprays have dried. Protective clothing should include: hat or other suitable head covering, long sleeved shirt and long legged trousers, or a coverall type garment, shoes and socks.

Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. The front panel PRECAUTIONARY STATEMENTS should be read to workers as well as the instruction not to enter until sprays have dried. When oral warnings are given, warnings shall be given in language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information:

CAUTION: Area treated with PROPANIL on (date of application). Do not enter without appropriate protective clothing until sprays have dried. (Insert here Statements of Practical Treatment as on front panel.)

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not store this product near fertilizers, seeds, insecticides, or fungicides. Containers should not be stacked more than 3 containers high. Reclose all partially used containers by thoroughly tightening screw cap. Damaged or leaking containers which contain product that cannot be used immediately should be transferred to suitable sound containers and properly marked. Any spilled material should be thoroughly absorbed with a suitable absorbant, swept up and transferred to a new or waste container for disposal as indicated under Pesticide Disposal.

For safety and prevention of unauthorized use, all pesticides should be stored in locked facilities.

To prevent accidental misuse, different pesticides should be stored in separate areas with enough distance between to provide clear identification. Opened, partially used pesticides should be stored in original containers when possible. When transfer to another container is necessary because of leakage or damage, carefully mark and identify contents of the new container.

1,6

4

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of federal law. Wastes resulting from the use of this product may be disposed of at an approved waste disposal facility. If these wastes cannot be disposed of by use according to label instructions, contact your state pesticide or environmental control agency or the hazardous waste representative at the nearest EPA regional office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent), adding rinsate to spray tank. Offer rinsed containers for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

WHERE TO USE

CEDAR CHEMICAL'S PROPANIL FLOWABLE is used for postemergence control of broadleaf and grass weeds in RICE fields.

WEEDS CONTROLLED

Barnyardgrass (watergrass), brachiaria, coffeeweed, crabgrass, croton, curly indigo, foxtail, goosegrass, gulf cockspur, mexicanweed, millet (Texas), paragrass, pigweed, sourdock, spearhead, wiregrass.

(PROPANIL FLOWABLE will not control arrowhead, bermudagrass, cattail, ducksalad, johnsongrass, nutgrass, red rice and sprangletop.)

GENERAL INFORMATION

Several important factors should be taken into account to achieve a high efficiency of selective weed control with PROPANIL FLOWABLE. These include uniform application, growth stage and weather conditions. To assure uniform application mix the prescribed amount of PROPANIL FLOWABLE with a sufficient volume of water to provide thorough coverage of target area. For aerial applications use approximately 10 gallons of water, or for surface (ground) applications 20-30 gallons of water per acre at sufficient spray pressure. Agitate tank mixes thoroughly and continuously. Avoid over and under application.

Growth stage of weeds is very important. Best results for selective weed control are obtained when most grasses have reached the 1 to 3 leaf stage.

Proper field preparation is essential to ascertain a relatively clod free and level surface and to obtain uniform flood levels and growth. Fields may be flushed prior to treatment to produce uniform and vigorous grass germination and growth. Drain water from fields prior to applying PROPANIL FLOWABLE. Higher

15/6

rates are recommended to control larger grasses or exposed weeds when rice fields are not completely drained. Inspect rice fields regularly to select the correct application time.

DO NOT apply this product through any type of irrigation system.

Weather conditions must be observed closely. Under cool weather conditions higher rates are required to achieve satisfactory control. Avoid application if rain threatens within 6 to 8 hours, or if wind velocities are high enough to cause drift and irregular spray patterns.

When Propanil Flowable is used alone (not in combination with any other postemergent rice herbicide), a suitable non-ionic surfactant or crop oil may be used to improve wetting of foliage and weed control. Refer to the surfactant or crop oil label for proper use rates.

Consult Extension Service for detailed application advice.

RECOMMENDED BROADCAST RATE

Apply 3 quarts of PROPANIL FLOWABLE per acre when most grasses have reached 1 to 3 leaf stage. Use 4 to 6 quarts of PROPANIL FLOWABLE per acre when the grasses are large (4 to 6 leaf stage) or when unseasonably cool weather conditions prevail or in cases where rice fields have not been drained completely and where weeds are large.

Barnyardgrass may be controlled up to 30 to 45 days after planting, before rice plants have reached the fully tillered growth stage.

NOTE: PROPANIL FLOWABLE applied to rice after the 4 leaf stage may cause visible injury under some climatic conditions. Rice plants usually outgrow such injury.

NOTE: Water drained from treated rice fields must not be used to irrigate other crops or be released within 1/2 mile upstream of a potable water intake in flowing water (i.e. river, stream, etc.) or within 1/2 mile of a potable water intake in a standing body of water such as a lake, pond or reservoir

IN CALIFORNIA--Use PROPANIL FLOWABLE only where rice fields are not completely drained. If higher water level is desired, reflood field after 12 hours and before 7 days after treatment. This will discourage new weed infestations. Do not apply PROPANIL FLOWABLE within 14 days before or after insecticide applications. Serious injury to rice may occur.

CAUTION

Do not apply this product (directly or indirectly) to any crop except rice.

Do not apply when wind conditions will allow drift to adjacent, susceptible crops such as beans, soybeans, cotton, safflower, cucurbits, vegetables, orchards (such as almonds, prunes and grapes) and other sensitive crops.