

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0080. Approval expires 05-31-98



United States
Environmental Protection Agency
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number
253598

Application for Pesticide - Section I

1. Company/Product Number 56077-52	2. EPA Product Manager Joanne Miller	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Butoxone 7500	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) Cedar Chemical Corp. 5100 Poplar Ave., Suite 2414 Memphis, TN 38137 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.) To add the general notice that INJURY TO CROPS FROM 2,4-DB PRODUCTS MAY OCCUR. IF YOU ARE NOT PREPARED TO ACCEPT SOME DEGREE OF CROP INJURY, DO NOT USE THIS PRODUCT, and to add the same notice in the soybeans section stating that INJURY TO SOYBEANS FROM 2,4-DB PRODUCTS...

NOTIFICATION
JAN 2 1997

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	If "Yes" Package wgt. No. per container		<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container			
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		<input type="checkbox"/> Other _____			

Section - IV


1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Jeanine H. Betsher	Title Mgr., Regulatory Affairs	Telephone No. (Include Area Code) 901-685-5348
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment both under applicable law.		6. Date Application Received (Stamped)
2. Signature <i>Jeanine H. Betsher</i>	3. Title Mgr., Regulatory Affairs	
4. Typed Name Jeanine H. Betsher	5. Date 12/10/96	

We have had 5 soybean crop injury complaints in the last 6 - 7 years. When local universities have been consulted regarding the incidents, they have offered no physiological/environmental explanation, but insisted that the injury, though rare, is consistent with 2,4-DB use on soybeans.

Cedar's Butoxone labels have use warnings of injury due to 2,4-DB use under adverse growing conditions; however, when used according to label directions, these warnings offer Cedar no defense to the crop injury complaints mentioned above, particularly when university extension personnel consider this type of injury "classic 2,4-DB injury".

Therefore, we propose the addition of a statement in 2 locations on the label: first, in the general information section with other Do Not statements, and second, in the Soybeans section.

BEST AVAILABLE COPY



375
After



Butoxone® 7500

Herbicide

For the control of certain seedling broadleaf weeds in peanuts, soybeans, and seedling stands of alfalfa, and for use on established stands of alfalfa.

ACTIVE INGREDIENT:	
4-(2,4-dichlorophenoxy) butyric acid	75.00%
INERT INGREDIENTS: 25.00%	
TOTAL	100.00%

*This product contains 0.75 lb. of 4-(2,4-dichlorophenoxy) butyric acid per pound of formulated product.

NOTIFICATION

EPA Registration No. 56077-52
EPA Establishment No. 62171-MS-01

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching the back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting.

If On Skin: Wash with plenty of soap and water. Get medical attention.

If In Eyes: Flush with plenty of water. Call a physician if irritation persists.

FOR CHEMICAL SPILL, LEAK, FIRE OR EXPOSURE CALL TOLL FREE
1-800-424-9300

AGRICULTURAL CHEMICAL
DO NOT SHIP OR STORE WITH FOODS, FEEDS, DRUGS, OR CLOTHING

PRECAUTIONARY STATEMENTS

CAUTION

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:
• long-sleeved shirt and long pants
• waterproof gloves
• shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:
• Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
• Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not contaminate water intended for irrigation or domestic purposes. Do not apply when weather conditions favor drift from target area.

BEST AVAILABLE COPY

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- coveralls
- waterproof gloves
- shoes plus socks

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not store this product near fertilizers, seeds, insecticides, or fungicides. Containers should not be stacked more than six (6) high. Reclose all partially used containers by tying bag top shut. Damaged or leaking containers which cannot be used immediately should be transferred to suitable sound containers and properly marked. Any spilled material should be thoroughly swept up and transferred to the new container or disposed of as indicated under "Pesticide Disposal".

For safety and prevention of unauthorized use, all pesticides should be stored in locked facilities.

To prevent accidental misuse, different pesticides should be stored in separate areas with enough distance between to provide clear identification.

Opened, partially used pesticides should be stored in original labeled containers when possible. When transfer to another container is necessary because of leakage or damage, carefully mark and identify contents of the new container.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your state pesticide or environmental control agency or the hazardous waste representative at the nearest EPA regional office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Apply this product only as specified on this label.

NOTE: Avoid spray drift to susceptible plants and crops: cotton, tobacco, tomatoes, ornamentals, etc. Coarse sprays are less likely to drift.

Spray equipment previously used to apply another phenoxy must be thoroughly cleaned with alkali and water prior to use with Butoxone 7500. Similarly, after using Butoxone 7500 wash sprayer thoroughly before using again to spray susceptible crops. Do not use near greenhouses. Observe all caution and limitations on labeling of all products used in mixtures.

Use agitation to keep solution well mixed, especially if it has been

allowed to stand unagitated.

Calibrate sprayer accurately.

INJURY TO CROPS FROM 2,4-DB PRODUCTS MAY OCCUR. IF YOU ARE NOT PREPARED TO ACCEPT SOME DEGREE OF CROP INJURY, DO NOT USE THIS PRODUCT.

DO NOT APPLY WHEN CROP IS STRESSED FROM LACK OF MOISTURE.

DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM.

GROUNDWATER CONTAMINATION: Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-DB have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-DB pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will prevent groundwater contamination.

ALFALFA

APPLICATION:

For use in seedling alfalfa, spray when the crop has reached the 1 to 2 trifoliate leaf stage and growing conditions are good. In established alfalfa certain weeds will emerge in the fall and over winter in a rosette stage. Best control of these weeds will result from application in late fall or early winter rather than in the spring. Do not apply after flowering.

WHEN TO APPLY:

Weeds must be in the young seedling stage and actively growing to achieve satisfactory results. For best results spray weeds in the 2 to 5 leaf stage of growth.

AMOUNT TO USE:

Apply Butoxone 7500 as an overall spray by ground sprayer or airplane. Apply at rates listed below according to weed problems. Use the higher rate if weeds are past the seedling stage. Aerial applications should be made in at least 5 gallons of water per acre, while ground sprayer applications should be in a minimum of 10 gallons of water per acre. Higher spray gallonage per acre of 10 gallons by aerial application and 20 to 30 gallons for ground sprayer applications will give better coverage and weed control. Spray gallonage should give adequate coverage of the weeds without run-off. The use of a non-ionic or non-ionic/anionic surfactant or crop oil concentrate approved for agricultural uses at their label rates will usually result in better weed control, especially if weeds are beyond the seedling stage.

BROADLEAF WEEDS	Rate of Butoxone 7500 per Acre
Annual Morningglory (Ipomoea spp.)	1.33 pounds
Cocklebur (Xanthium spp.)	
Common Lambsquarters (Chenopodium album)	
Jimsonweed (Datura stramonium)	
Kochia or Mexican Fireweed (Kochia scoparia)	
Pigweed (Amaranthus spp.)	
Velvetleaf (Abutilon theophrasti)	
Wild Turnip (Brassica campestris)	
Black Mustard (Brassica nigra)	2.0 pounds
Buckhorn Plantain (Plantago lanceolata)	
Common Ragweed (Ambrosia artemisiifolia)	
Curly Dock (Rumex crispus)	
Field Pennycress (Fanweed or Stinkweed) (Thlaspi arvense)	
Hedge Smartweed (Polygonum scandens)	
Ladysthumb (Polygonum persicaria)	
Prickly Lettuce (Lactuca serriola)	
Shepherdspurse (Capsella bursa-pastoris)	
Sweetclover (Volunteer plants) (Melilotus spp.)	
Wild Beet (Beta maritima)	
Wild Mustard (Brassica kaber)	

established alfalfa to livestock within 30 days after application. Do not graze or feed SEEDLING alfalfa within 60 days after application. Irrigation, in particular overhead sprinkler irrigation, should be delayed as long as possible (10 days or more) following application of Butoxone 7500 to avoid washing the chemical into the root zone. Do not apply when crop is stressed from lack of moisture. DO NOT use on peas.

Apply as a postemergent spray. When properly timed there is little or no effect on the crop. In established alfalfa there may be some twisting of stems and malformation of leaves. This condition is usually outgrown.

Do not spray when daytime temperatures are expected to exceed 90°F within the next 2 or 3 days. Do not apply if temperatures are likely to fall below 40°F during or shortly after treatment.

Use of a surfactant or crop oil concentrate in the desert areas of California and Arizona may cause some crop injury under certain climatic and crop stress conditions. Do not use a surfactant unless possible crop injury is acceptable.

PEANUTS

APPLICATION:

Apply Butoxone 7500 postemergent as an overall spray by ground sprayer or airplane. With ground applications use a boom sprayer with flat fan-type nozzle. Adjust the height of the boom above the crop so the growing terminals of weeds are sprayed. Use low pressure (about 20 to 40 lbs. per square inch). Do not use boom jet nozzles. With airplane applications Butoxone 7500 has been successfully applied using 5 gallons of total solution per acre. The height of the application should insure THOROUGH COVERAGE of the tops of the weeds. Best control has resulted from airplane application with coarse spray when cocklebur is at least crop high and actively growing. Other weed species should be in the seedling stage and actively growing. Airplane application is not recommended in the vicinity of susceptible crops.

Do not apply if peanuts are suffering from lack of moisture.

Amount to Use: Apply the recommended rates as indicated in the table below.

BROADLEAF WEEDS	Rate of Butoxone 7500 per Acre
Cocklebur (Xanthium spp.)	0.3 pound
Annual Morningglory (Ipomoea spp.)	0.5 pound
Common Lambsquarters (Chenopodium album)	
Common Ragweed (Ambrosia artemisiifolia)	
Jimsonweed (Datura stramonium)	
Sicklepod (Coffeeweed) (Cassia obtusifolia)	
Velvetleaf (Abutilon theophrasti)	

Do not apply more than two (2) applications per season; second application should be made no later than the late bloom stage of peanuts (about 90 to 100 days after planting). Do not feed treated peanut vines or peanut hay to livestock. Do not apply to peanuts within 30 days of harvest.

SOYBEANS

APPLICATION: Apply Butoxone 7500 postemergent as an overall spray by ground sprayer or airplane. With ground applications use a boom sprayer with flat fan-type nozzles. Adjust the height of the boom above the crop so the growing terminals of all weeds are sprayed. Use low pressure (about 20 to 40 lbs. per square inch). Do not use boom jet nozzles. With airplane applications Butoxone 7500 has been successfully applied using 5 gallons of total spray solution per acre. The height of the application should insure thorough coverage of the tops of the weeds. Best control has resulted when cocklebur is at least crop high and actively growing. Airplane application is not recommended in the vicinity of susceptible crops. Do not apply if soybeans are suffering from lack of moisture.

TOPICAL (OVERHEAD) APPLICATIONS

Apply from 7 to 10 days before soybeans bloom through mid-bloom. A good indication that the beans are about to bloom is

Do not graze established alfalfa, nor feed straw or hay from

595

when the soybean plants have turned a dark green color. If application is made 7 to 10 days before bloom, apply 2.33 pounds of Butoxone 7500 per 10 acres; if application is made from early-bloom through mid-bloom, apply 2.33 pounds of Butoxone 7500 per 8 acres. Use sufficient water to obtain adequate coverage. The rates recommended are for the control of cocklebur. These rates will also stunt or partially control certain other weeds including annual morningglory, velvetleaf, and jimsonweed. In the Midwest use only a directed application since topical application may reduce soybean yields.

NOTICE: INJURY TO SOYBEANS FROM 2,4-DB PRODUCTS MAY OCCUR. IF YOU ARE NOT PREPARED TO ACCEPT SOME DEGREE OF CROP INJURY, DO NOT USE THIS PRODUCT. Application should not be made to sparsely foliated stands stressed because of disease or lack of moisture. It is essential that the canopy above the soybeans be complete. To apply the product otherwise could result in crop damage such as stalk and stem splitting, and reduction in yield.

DIRECTED APPLICATIONS

For best results apply when weeds do not exceed 3 inches in height. Good coverage of the growing terminals of weeds is essential for effective control. Do not allow spray pattern to contact more than the lower one-third of the soybean plants.

Do not apply more than two (2) applications per season. Precise application is essential to prevent damage to the crop. In order to maintain the correct spraying height, nozzles must be mounted on oiling shoes, skid shoes, or on cultivators with gauge wheels. Do not use booms with drop nozzles.

First Application: Apply as a directed spray when soybeans are 8 to 12 inches tall and weeds have emerged.

Second Application: Apply as a directed spray no later than mid-bloom stage.

Amount to Use: Apply the recommended rates as indicated in the table below.

BROADLEAF WEEDS	Rate of Butoxone 7500 per Acre
Cocklebur (<i>Xanthium</i> spp.)	0.3 pound
Annual Morningglory (<i>Ipomoea</i> spp.)	0.5 pound
Common Lambsquarters (<i>Chenopodium album</i>)	
Common Ragweed (<i>Ambrosia artemisiifolia</i>)	
Jimsonweed (<i>Datura stramonium</i>)	
Sicklepod (Coffeeweed) (<i>Cassia obtusifolia</i>)	
Velvetleaf (<i>Abitilon theophrasti</i>)	

While some temporary twisting of soybean plants may follow an application, this will have little or no permanent effect on the crop.

Applications made under adverse growing conditions, such as periods of drought, will result in injury to the beans; under such conditions the crop should not be sprayed.

Do not spray Butoxone 7500 on soybeans showing a general infestation of *Phytophthora*.

Do not apply more than the recommended amount.

Do not harvest soybeans within 60 days after application.

Do not use on soybeans grown West of the Rocky Mountains.

DIRECTED BAND APPLICATIONS OF MIXTURES OF BUTOXONE 7500 AND LOROX* 50 W

Application: Apply directed spray when beans are at least 8 inches tall and when weeds do not exceed two inches in height. Direct spray to cover weed foliage with minimum contact of the soybean plant. Do not spray higher than 3 inches on the soybean stem or crop injury may result. Use an 8002T-Jet (or equivalent) or larger with minimum of 25 gallons per broadcast acre and spray pressure of 20 to 25 psi, to avoid spray drift. For each 25 gallons of spray mixture a pint of surfactant such as Surfactant WK may be added. Continuous agitation in the spray tank is required to keep the material in suspension. Agitate by mechanical or hydraulic means; if by-pass or return line is used, it should terminate at the bottom of the tank to minimize foaming. Openings in nozzle screen should be equal to or larger than 50 mesh. Apply with sprayer nozzles mounted on skid shoes, oiling shoes, or on cultivators with gauge wheels. Do not mount on booms with drop nozzles or on cultivators without gauge wheels.

Amount to Use: Apply the recommended rates as indicated in the table below.

BROADLEAF WEEDS	Rate of Butoxone 7500 and Lorox per broadcast acre
Annual Morningglory (<i>Ipomoea</i> spp.)	0.3 pound
Cocklebur (<i>Xanthium</i> spp.)	1 pound
Common Lambsquarters (<i>Chenopodium album</i>)	
Common Ragweed (<i>Ambrosia artemisiifolia</i>)	
Crabgrass (<i>Digitaria</i> spp.) (2 leaf)	
Jimsonweed (<i>Datura stramonium</i>)	
Prickly Sida (Teaweed) (<i>Sida spinosa</i>)	
Sesbania (<i>Sesbania</i> spp.)	
Velvetleaf (<i>Abitilon theophrasti</i>)	

Use a proportional amount of the two chemicals on the band (for example, on a 12 inch band in 36 inch rows use 0.1 pound of Butoxone 7500 and ¼ lb. of Lorox per acre). If a new flush of weeds occurs after initial treatment, make a second application but do not apply within 60 days of harvest or make more than 2 applications per growing season. Do not allow spray or spray drift to contact growing terminals of beans, as excessive crop injury will result. Do not feed soybean forage or hay to livestock from fields treated with this mixture. Harvested soybeans may be used for feed or oil purposes.

NOTE: Do not use on sands, loamy sands, gravelly soils, or on exposed sub-soils. Do not use on soils containing less than ½ % organic matter as crop injury may result. Observe all cautions and limitations of labeling of all products used in mixture.

NOTE: The Lorox 50W, Butoxone 7500 tank mix is not approved for use in California.

* Lorox 50W is a registered trademark of E.I. duPont de Nemours & Co. Inc.

CONDITIONS OF SALE AND WARRANTY

CEDAR AND SELLER OFFER THIS PRODUCT AND THE BUYER AND USER ACCEPTS THIS PRODUCT UNDER THE FOLLOWING AGREED CONDITIONS OF SALE AND WARRANTY.

The directions for use of this product are believed to be reliable and should be followed carefully. However, it is impossible to take into account all variables and to eliminate all risks associated with its use. Injury or damage may result because of conditions which are beyond the control of Cedar or the Seller. Cedar warrants only that this product conforms to the chemical description on the label and is believed to be reasonably fit for the purposes referred to in the Directions for Use when used as directed under normal conditions. CEDAR MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. In no case shall Cedar or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. Any variation or exception from this warranty must be in writing and signed by an authorized Cedar representative.

**CEDAR CHEMICAL CORPORATION
5100 POPLAR AVENUE
24TH FLOOR
MEMPHIS, TENNESSEE 38137, USA**

BEST AVAILABLE COPY

