



U.S. ENVIRONMENTAL PROTECTION AGENCY  
Office of Pesticide Programs  
Registration Division (HT305C)  
421 "M" St., S.W.  
Washington, D.C. 20460

EPA Reg.  
Number:  
56077-68

Date of Issuance:  
MAR 21 2002

NOTICE OF PESTICIDE:

Registration  
Reregistration

(under FIFRA, as amended)

Term of Issuance:

Conditional, with  
Expiration Date of  
March 21, 2007

Name of Pesticide Product:

INTER Plus II  
Herbicide

Name and Address of Registrant (include ZIP Code):

Cedar Chemical Corporation  
5100 Poplar Ave., Suite 2414  
Memphis, TN 38137

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Add the phrase "EPA Registration No. 56077-68 to the label before you release the product for shipment.

COMMENTS CONTINUED ON PAGE 2 OF THIS NOTICE OF REGISTRATION

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product under the enclosed stamped copy of the label constitutes acceptance of these conditions.

Enclosure

Joanne I. Miller  
Product Manager (23)  
Herbicide Branch  
Registration Division (7505C)

Signature of Agent or Official:

*Joanne I. Miller*

Date:

MAR 21 2002

Comments Continued:

2. Submit the following data required for the reregistration of this pesticide product within 36 months of the date of this Notice of Registration:

<u>EPA Guideline Data Number</u>	<u>Guideline Descriptor</u>
71-4(a)	Avian Reproduction in Bobwhite Quail
72-4(b)	Aquatic Invertebrate Life-Cycle
164-1	Turf Field Dissipation
132-1(a)	Foliar Residue Dissipation
133-3	Dermal Passive Dosimetry Exposure

3. Submit the following data required for the reregistration of this pesticide product within 48 months of the date of this Notice of Registration:

<u>EPA Guideline Data Number</u>	<u>Guideline Descriptor</u>
166-1	Small Scale Prospective Ground Water Monitoring Study

4. The statement in the "Rotational Crops; "Inter Plus II Alone" section that reads: "All other rotational crops may be planted 12 months after application." must be revised to read: "All other rotational crops may be planted 12 months after a layby application."
5. Submit and/or cite all data required for the registration of this product when the Agency requires all registrants of similar products to submit data; and submit acceptable responses required for reregistration of this product under FIFRA, section 4.
6. Submit one (1) copy of the final printed labeling before you release the product for shipment
-

Rec'd 03-30-02  
via. E-mail  
Jmw.

# INTER Plus II

## HERBICIDE

For weed control in corn, cotton, peanuts, pod crops, potatoes, safflower, grain or forage sorghum, and soybeans

**Active Ingredient:**

Metolachlor: 2-chloro- N-(2-ethyl-6-methylphenyl)-N-(2-methoxy-1-methylethyl) acetamide

84.4%

**Inert Ingredients:**

15.6%

Total:

100.0%

INTER Plus II contains 7.8 lbs. of active ingredient per gallon.  
See directions for use in attached booklet.

EPA Reg. No. 56077-AI  
EPA Est. No. 56077-AR-001

### KEEP OUT OF REACH OF CHILDREN

### CAUTION

#### FIRST AID

<b>IF INHALED</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration preferably by mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>IF ON SKIN</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF IN EYES</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 15 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF SWALLOWED</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
FOR CHEMICAL SPILL, LEAK, FIRE OR EXPOSURE CALL TOLL FREE 1-800-424-9300	

Cedar Chemical Corp.  
5100 Poplar Avenue, Suite 2414  
Memphis, Tennessee 38137

#### NET CONTENTS:

**ACCEPTED  
with COMMENTS  
In EPA Letter Dated**

MAR 21 2002

**Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.  
56077-68**

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION**

Harmful if swallowed, inhaled, or absorbed through skin. Causes eye irritation. Avoid contact with skin, eyes, or clothing. Avoid breathing vapor or spray mist.

**PERSONAL PROTECTIVE EQUIPMENT**

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category H on an EPA chemical resistance category selection chart.

**Applicators and other handlers must wear:**

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves, such as barrier laminate or viton
- Chemical-resistant footwear plus socks
- Chemical-resistant headgear for overhead exposure
- Chemical-resistant apron when cleaning equipment

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**Engineering Control Statements**

Mixers and loaders supporting aerial applications are required to use closed systems. The closed system must be used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4)). When using the closed system, the mixers and loaders PPE requirements may be reduced or modified as specified in the WPS.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**User Safety Recommendations**

**Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters or rinsate.

**Ground Water Advisory:**

This chemical is known to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

**Surface Water Advisory:**

Metolachlor can contaminate surface water through ground spray drift. Under some conditions, metolachlor may also have a high potential for runoff into surface water (primarily via dissolution in runoff water), for several months post-application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas over-laying extremely shallow ground water, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-laying tile drainage systems that drain to surface water.

**Mixing/Loading Instructions:**

Care must be taken when using this product to prevent back-siphoning into wells, spills, or improper disposal of excess pesticide, spray mixtures, or rinsates.

Check-valves or antisiphoning devices must be used on all mixing and/or irrigation equipment.

This product may not be mixed or loaded within 50 ft. of perennial or intermittent streams and rivers, natural or impounded lakes and reservoirs. This product may not be mixed/loaded or used within 50 ft. of all wells, including abandoned wells, drainage wells, and sink holes. Operations that involve mixing, loading, rinsing, or washing of this product into or from pesticide handling or application equipment or containers within 50 ft. of any well are prohibited unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or wash-water, and rain water that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self-contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above-specified minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading site.

**STORAGE AND DISPOSAL****Pesticide Disposal**

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Wastes resulting from the use of this product are toxic. Improper disposal of unused pesticide, spray mixture, or rinsate is a violation of Federal law. Pesticide, spray mixture, or rinsate that cannot be used according to label instructions must be disposed of according to Federal, state, or local procedures. For guidance in proper disposal methods, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office.

**Container Disposal**

Do not reuse empty container. Triple rinse (or equivalent), puncture and dispose of in a sanitary landfill, or by incineration, or by open burning, if allowed by state and local authorities. If burned, keep out of smoke. This product may be stored at temperatures down to 30 degrees below 0° F.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

**DIRECTIONS FOR USE**

This product is intended for use in weed control in corn, peanuts, pod crops, potatoes, safflower, grain or forage sorghum, and soybeans.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.** Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter

the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or viton
- Shoes plus socks

**FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.**

To avoid spray drift, do not apply under windy conditions. Avoid spray overlap, as crop injury may result.

**A. General Information**

Observe all precautions and limitations on the labels of each product used in tank mixtures. Tank mixtures are permitted only in those states where the tank mix partner is registered.

INTER Plus II is a selective herbicide recommended as a preplant surface-applied, preplant incorporated, or preemergence treatment in water or fluid fertilizer for control of most annual grasses and certain broadleaf weeds in corn (all types), cotton, peanuts, pod crops, potatoes, safflower, grain or forage sorghum, and soybeans.

**Note:** Do not use in nurseries, turf, or landscape plantings.

Do not apply under conditions which favor runoff or wind erosion of soil containing this product to non-target areas.

To prevent off-site movement due to runoff or erosion,

1. Avoid treating powdery dry or light sand soils when conditions are favorable for wind erosion. Under these conditions, the soil surface should first be settled by rainfall or irrigation.
2. Do not apply to impervious substrates such as paved or highly compacted surfaces.
3. Do not use tailwater from the first flood or furrow irrigation of treated fields to treat non-target crops unless at least 1/2 inch of rainfall has occurred between application and the first irrigation.

Where directions specify an INTER Plus II tank mixture with AAtrex formulations, other brands of atrazine may be used. Follow the rates, recommendations, and limitations on the AAtrex or respective atrazine product label, if other brands of atrazine are used.

**Note:** Certain states may have established rate limitations for atrazine within specific geographical areas. Consult your state lead pesticide control agency for additional information. It is a violation of this label to deviate from state use regulations.

If INTER Plus II is incorporated, any supplemental tillage before planting must not exceed the depth of incorporation.

Dry weather following preemergence application of INTER Plus II or a tank mixture may reduce effectiveness. Cultivate if weeds develop.

Where reference is made to weeds partially controlled, partial control can either mean erratic control from good to poor, or consistent control at a level below that generally considered acceptable for commercial weed control.

*Precaution: Injury may occur to crops other than corn following the use of INTER Plus II under abnormally high soil moisture conditions during early development of the crop.*

**1. Soil Textures and Herbicide Rates**

Where rates are based on coarse-, medium-, or fine-textured soils, it is understood that soil textural classes are generally categorized as follows:

**Coarse**  
Sand  
Loamy sand  
Sandy loam

**Medium**  
Loam  
Silt loam  
Silt

**Fine**  
Sandy clay loam  
Silty clay loam  
Clay loam

**Fine (Cont'd)**  
Sandy clay  
Silty clay  
Clay

Within rate ranges in the rate tables and elsewhere on this label, use the lower rate on soils relatively coarse-textured or low in organic matter; use the higher rate on soils relatively fine-textured or high in organic matter.

**Note:** INTER Plus II may be applied preemergence alone, or in combination with tank mix partners specified on this label, following preplant incorporated herbicides when used according to their label recommendations, provided that such use is not prohibited on the respective labels.

Thoroughly clean sprayer or other application device before using. Dispose of cleaning solution in a responsible manner. Do not use a sprayer or applicator contaminated with any other materials, or crop damage or clogging of the application device may result.

## 2. Mixing Instructions

**INTER Plus II Alone:** Mix INTER Plus II with water or fluid fertilizer and apply as a spray. Fill the spray tank 1/2 - 3/4 full with water or fluid fertilizer, add the proper amount of INTER Plus II, then add the rest of the water or fluid fertilizer. Provide sufficient agitation during mixing and application to maintain a uniform emulsion.

**Tank Mixtures:** Fill the spray tank 1/4 full with water, and start agitation; add 2,4-D, AAtrex, Balan, Banvel, Canopy, Caparol 4L, Command, Cotoran, Eptam, Gemini, Lexone, Lorox, Lorox Plus, Marksman, Preview, Princep, Prowl, Pursuit, AAtrex + Princep, Scepter, Sencor, Sonalan, or Treflan, and allow it to become dispersed; then add INTER Plus II; then add Gramoxone Extra, Landmaster BW, or Roundup if these products are being used; and finally the rest of the water. For tank mixtures with AAtrex, Banvel, Canopy, Caparol 4L, Command, Cotoran\*, Eptam, Gemini, Lexone, Lorox, Lorox Plus, Marksman, Preview, Princep, Prowl \*, Pursuit, AAtrex + Princep, Scepter, Sencor, Sonalan, or Treflan, fluid fertilizers may replace all or part of the water as carrier, except in the AAtrex postemergence and the Banvel postemergence tank mixes. For tank mixtures with AAtrex, see additional mixing instructions on the AAtrex label. For each mixture, check compatibility with fluid fertilizer, as described below, before mixing in spray tank. For all tank mixtures, agitate during mixing and application to maintain a uniform suspension.

\*See **Special Mixing Instructions** for tank mixtures with Cotoran, and with AAtrex or Princep + Prowl under the appropriate tank mixture section.

For directions on how to conduct a compatibility test, see **Appendix A**.

## 3. Application Procedures

### Application Timing

INTER Plus II alone or in some tank mixtures with other labeled herbicides may be applied for weed control in certain crops at various times. Refer to the given crop section of the label to determine if application timings listed below are recommended.

**a. Preplant Surface-Applied:** For minimum-tillage or no-tillage systems only, INTER Plus II alone and some INTER Plus II tank mixtures may be applied up to 45 days before planting certain crops. Use only split applications for treatments made 30-45 days before planting, with 2/3 the recommended broadcast rate for the crop and soil texture applied initially and the remaining 1/3 at planting. Treatments less than 30 days before planting may be made either as a split or a single application. Refer to individual crop to determine if early preplant surface application is recommended. If weeds are present at the time of treatment, apply in a tank mixture combination with a contact herbicide (for example, Gramoxone Extra or Roundup). Observe directions for use, precautions, and restrictions on the label of the contact herbicide.

To the extent possible, do not move treated soil out of the row or move untreated soil to the surface during planting, or weed control will be diminished.

**b. Preplant Incorporated:** Apply INTER Plus II to the soil and incorporate into the top 2 inches of soil within 14 days before planting, using a finishing disk, harrow, rolling cultivator, or similar implement capable of providing uniform 2-inch incorporation. Use a preplant incorporated application if furrow irrigation is used or when a period of dry weather after application is expected. If crop will be planted on beds, apply and incorporate INTER Plus II after bed formation, unless specified otherwise.

**c. Preemergence:** Apply INTER Plus II during planting (behind the planter) or after planting, but before weeds or crops emerge.

#### 4. Special Application Procedures

**Preplant Incorporated—CA Only (Corn, Safflower, Pod Crops):** Broadcast INTER Plus II alone or with approved tank mix partners to the soil and thoroughly incorporate with a disk or similar implement set to till 4 - 6 inches deep. For more thorough incorporation, till the soil in 2 different directions (cross-till). Crops may be planted on flat surface or on beds. Caution should be used when forming the beds that only soil from the INTER Plus II treated zone is used (i.e., untreated soil should not be brought to soil surface). If the application is made to preformed beds, incorporate INTER Plus II with a tillage implement set to till 2 - 4 inches deep. Care should be taken during tilling to keep the tilled (INTER Plus II treated) soil on the beds.

**Preemergence:** Apply INTER Plus II after planting. Water with sprinkler or flood irrigation within 7-10 days.

**Fall Application (Only in IA, MN, ND, SD, WI, and portions of NE and IL - see specific instructions in the Corn, Soybeans, and Pod Crops sections of this label for timing of application and other information):** Do not apply to frozen ground. Use on medium and fine soils with greater than 2.5% organic matter that will be planted to corn or soybeans the next spring. Ground may be tilled before or after application. Do not exceed a 2 to 3-inch incorporation depth if tilled after treatment. **Note:** If a spring application is made, the total rate of the fall plus spring applications must not exceed the maximum total rate for the specific crop, or illegal residues may result.

**Ground Application:** Apply INTER Plus II alone or in tank mixtures by ground equipment in a minimum of 10 gals. of spray mixture per acre, unless otherwise specified.

Use sprayers that provide accurate and uniform application. For INTER Plus II tank mixtures with wettable powder or dry flowable formulations, screens and strainers should be no finer than 50-mesh. Rinse sprayer thoroughly with clean water immediately after use.

Calculate the amount of herbicide needed for band treatment by the formula:

$$\frac{\text{band width in inches}}{\text{row width in inches}} \times \text{broadcast rate per acre} = \text{amount needed per acre of field}$$

**NOTE:** For information on applying in lower volumes of carrier, see **Low Carrier Application in Appendix B.**

For application by air or through center pivot systems, see **Appendices C and D.** **Appendix C** includes **Aerial Drift Management and Aerial Drift Reduction Advisory** sections.

For information on impregnating dry fertilizer, see **Appendix E.**

#### B. INTER Plus II Applied Alone

##### 1. Weeds Controlled:

barnyardgrass  
(watergrass)  
bristly foxtail  
crabgrass  
crowfootgrass  
fall panicum



foxtail millet  
 giant foxtail  
 goosegrass  
 green foxtail  
 prairie cupgrass  
 red rice  
 robust foxtails  
 (purple, white)  
 signalgrass  
 (*Brachiaria*)  
 southwestern cupgrass  
 wild proso millet\*  
 witchgrass  
 woolly cupgrass\*  
 yellow foxtail  
 yellow nutsedge  
 carpetweed  
 common waterhemp  
 Eastern black  
 nightshade  
 Florida pusley  
 galinsoga  
 pigweed  
 tall waterhemp

\*For control of these weeds in corn only, refer to the **Corn - Woolly Cupgrass and Wild Proso Millet Control Program** section of this label.

**2. Weeds Partially Controlled\*:**

common purslane	hairy nightshade	Texas panicum***
eclipta	sandbur	volunteer sorghum
Florida beggarweed **	seedling johnsongrass	wild proso millet
	shattercane	wolly cupgrass

\*See **General Information** section. Control of these weeds can be erratic, due partially to variable weather conditions. Control may be improved by following these suggested procedures:

1. In corn, use 2 - 2.67 pts /A or the preplant surface-applied rates for INTER Plus II alone or in tank mixture, if allowed, when making preplant incorporated or preemergence applications.
2. **Thoroughly till moist soil** to destroy germinating and emerged weeds. If INTER Plus II is to be applied preplant incorporated, this tillage may be used to incorporate INTER Plus II if uniform 2-inch incorporation is achieved as recommended under **Application Procedures**.
3. Plant crop into moist soil **immediately after tillage**. If INTER Plus II is to be used preemergence, apply at planting or immediately after planting.
4. If available, **sprinkler irrigate** within 2 days after application. Apply 1/2 - 1 inch of water. Use lower water volume (1/2 inch) on *coarse-textured soils* and higher volume (1 inch) on *fine-textured soils*. Also, refer to the section on **Center Pivot Irrigation Application** for this method of applying INTER Plus II.
5. If irrigation is not possible and rain does not occur within 2 days after planting and application, weed control may be decreased. Under these conditions, a uniform, shallow cultivation is recommended as soon as weeds emerge.

\*\*For partial control of this weed, use a minimum of 2 pts /A and apply preemergence.

\*\*\*For partial control of this weed, use a minimum of 2 pts /A and apply through a center pivot irrigation system.

**2. Rotational Crops:**

**INTER Plus II Alone:** (1) If crop treated with INTER Plus II alone is lost, any crop on this label may be replanted immediately. Do not make a second broadcast application of INTER Plus II. If the original application was banded and the second crop is planted in the untreated row middles, a second banded treatment may be applied. (2) Barley, oats, rye, or wheat may be planted 4 1/2 months following treatment; alfalfa may be planted 4 months following application. Tomatoes may be planted 6 months following application. (3) Any crop on this label, in addition to root crops, tobacco, barley, buckwheat, milo, oats, rice, rye, wheat, cabbage, peppers, stone fruits, or tree nuts, may be planted in the spring following treatment. Clover may be seeded 9 months following application. Do not graze or feed forage

or fodder from cotton to livestock.(4) Following a lay-by treatment or multiple treatments applied the previous season, any crop on this label, in addition to tobacco, cabbage, peppers, stone fruits, or tree nuts may be planted in the spring. All other rotational crops may be planted 12 months after application.

**INTER Plus II Tank Mixtures:** For **Rotational Crops** restrictions for INTER Plus II used in tank mixtures, refer to the statements/restrictions above for INTER Plus II and to the respective product labels of any mixing partner(s) for additional statements/restrictions.

**Important Notes:** To avoid injury to rotational alfalfa or clover: (1) Do not apply more than 2 lbs. a.i. of metolachlor per acre (2 pts. of INTER Plus II) preemergence (including preplant surface, preplant incorporated, postplant incorporated, etc.), and (2) Do not make lay-by or other postemergent applications of INTER Plus II.

---

## CROPS:

---

### A. Corn (All Types)

#### INTER Plus II Alone

Apply INTER Plus II either preplant surface, preplant incorporated, preemergence, or lay-by, using the appropriate rate specified below.

##### 1. Preplant Surface-Applied:

Refer to instructions for use of INTER Plus II alone under **Application Procedures**.

**a. Fall Application (Apply after September 30 in MN, ND, SD, WI and north of Route 30 in IA; Apply after October 15 north of Route 91 in NE and south of Route 30 in IA; Apply after October 31 north of Route 136 in IL):** In all locations, apply to crop stubble after harvest when the sustained soil temperature at a 4-inch depth is less than 55 ° F and falling. In minimum-till or no-tillage systems on soils having greater than 2.5% organic matter, use 1.67 - 2.0 pts./A on *medium-textured* and 2.0 pts./A on *fine-textured soils*. Do not apply to frozen ground. A tillage operation may precede the application. A fall and/or a spring tillage may follow application, but do not exceed an incorporation depth greater than 2 - 3 inches. Minimize furrow and ridge formation in the tillage operations. **Note:** If a spring application is made, the total rate of the fall plus spring applications must not exceed the maximum total rate for corn, or illegal residues may result.

**b. Use on medium- and fine-textured soils with minimum-tillage or no-tillage systems in CO, IA, IL, IN, KS, KY, MN, MO, MT, ND, NE, SD, TN, WI, and WY.** Apply 2/3 the recommended rate of INTER Plus II(1.67 pts./A on *medium soils* and 2.0 pts./A on *fine soils*) as a split treatment 30 - 45 days before planting and the remainder at planting. Applications made less than 30 days prior to planting may be as either a split or single treatment. Apply 1.33 pts./A on *coarse soils* not more than 2 weeks prior to planting.

**c. On medium- and fine-textured soils with minimum- or no-tillage systems in CT, DE, MA, MD, ME, MI, NH, NY, OH, PA, RI, VA, VT, and WV,** preplant surface applications may be applied following the directions for use above. If the amount of rainfall results in unsatisfactory length of weed control following the earlier treatment, a postemergence application of an appropriately labeled broadleaf and/or grass weed herbicide may be used, i.e., AAtrex, Beacon<sup>®</sup>, Bicep<sup>®</sup>, Exceed<sup>®</sup>, Accent<sup>®</sup>, Banvel, Basagran<sup>®</sup>, bromoxynil (Brominal<sup>®</sup> or Buctril<sup>®</sup>), or 2,4-D. If the postemergence treatment includes the herbicide used preplant surface-applied, do not exceed the total labeled rate for corn on a given soil texture. Observe all directions for use, precautions, and limitations on the label of the postemergent herbicide.

##### 2. Preplant Incorporated or Preemergence:

Follow instructions for use of INTER Plus II alone under **Application Procedures**. On *coarse soils*, apply 1.0 - 1.33 pts./A of INTER Plus II if organic matter content is less than 3%, or 1.33 pts./A if organic matter content is 3% or greater. On *medium soils*, apply 1.33 - 1.67 pts./A of INTER Plus II. On *fine soils*, apply 1.33 - 1.67 pts./A of INTER Plus II if organic matter content is less than 3%, or 1.67 - 2.0 pts./A if organic matter content is 3% or greater.

**a. Lay-by:** To extend the duration of weed control in corn, a maximum rate of 2.0 pts./A of INTER Plus II

may be applied after corn emergence until the corn plants reach 40 inches in height, following any preplant surface-applied, preplant incorporated, or preemergence herbicide application, including INTER Plus II. For best results, applications should be made to soil free of emerged weeds and directed towards the base of corn plants in excess of 5 inches tall. The total INTER Plus II rate applied on corn during any one crop year should not exceed 4 pts./A, depending on soil texture.

**Note for all applications to corn:** To avoid possible illegal residues, do not graze or feed forage from treated areas for 30 days following application.

**b. Shattercane, Wild Proso Millet, Woolly Cupgrass, and Eclipta – Partial Control:** For more consistent partial control of shattercane, wild proso millet, woolly cupgrass, or eclipta, apply 2.0 - 2.55 pts./A as a single application; or apply 1.0 - 1.33 pts./A of INTER Plus II preplant incorporated followed by 1.0 - 1.33 pts./A of INTER Plus II preemergence; however, do not apply more than a total of 2.55 pts./A. Make the preemergence application during or after planting, but before weeds and corn emerge. Apply the 1.33 pts./A rate of INTER Plus II when a heavy infestation of shattercane, wild proso millet, woolly cupgrass, or eclipta is expected. A shallow cultivation may be needed to control any late emerging weeds.

**c. Woolly Cupgrass and Wild Proso Millet Control Program:** For control of these species, use the following 3-step program: (1) Apply INTER Plus II early preplant, preplant incorporated, or preemergence at 1.67 pts./A on *medium soils* and 2.0 pts./A on *fine-textured soils*, up to the maximum label rate. Lightly incorporate with a rotary hoe if rainfall does not occur within 5 - 7 days; (2) Apply a postemergence tank mix of Beacon at 0.38 oz./A or Exceed at 1 packet per 4 acres plus Accent SP at 0.33 oz./A plus 1 qt. of crop oil concentrate plus 1 gal./A of 28% nitrogen, or the equivalent amount of ammonium sulfate, when grasses are 2-3 inches tall and the corn is at least 4 inches tall; and (3) Cultivate 14-21 days after the postemergence application.

**Notes:** (1) Do not apply more than the labeled application rate for a given soil texture per year, either as a single or split treatment, or illegal residues may result. (2) In corn, INTER Plus II may be used up to 2.75 pts./A as either a preplant surface, preplant incorporated, or preemergence treatment on soils having an organic matter content between 6% and 20% or up to 2.0 pts./A on any soil for extended residual control and where severe stands of problem weeds are expected. (3) In the event of escape of annual weeds following a preplant surface, preplant incorporated, or preemergence treatment of INTER Plus II, follow with a postemergence application of an appropriately labeled broadleaf and/or grass weed herbicide, i.e., AAtrex, Beacon, Bicep, Exceed®, Accent, Banvel, Basagran, Brominal, Buctril, or 2,4-D. If the postemergence treatment includes the herbicide used in the earlier treatment i.e., AAtrex, do not exceed the total labeled rate for corn on a given soil texture. (4) Brominal or Buctril may be applied postemergence alone or in tank-mix combination with AAtrex. Do not exceed 1.2 lbs. a.i./A of AAtrex in tank-mix combination with Brominal or Buctril postemergence. Refer to the AAtrex, Brominal, and Buctril labels for specific rates and precautions. (5) Do not use INTER Plus II on peat or muck soils.

## A. Corn

### INTER Plus II Combinations

INTER Plus II in any tank mixture for corn (except INTER Plus II + AAtrex postemergence and INTER Plus II+ Banvel postemergence) may be applied in water or fluid fertilizer. Use only water in the INTER Plus II+ AAtrex or the INTER Plus II+ Banvel postemergence tank mixes.

**Note:** For all applications to corn, do not graze or feed forage from treated areas for 30 days following application, or possible illegal residues may result.

**IMPORTANT: FOR TANK MIXTURES WITH AATREX (OR OTHER BRANDS OF ATRAZINE) –** If applying INTER Plus II in tank mixture with AAtrex, all the restrictions and rate limitations on the AAtrex label must be followed if more restrictive/protective than those on this label. In addition, if AAtrex is/must be applied at rates lower than those recommended on this label, broadleaf weed control may be affected. Refer to the AAtrex label for weeds controlled at the reduced rates.

### Chart 1: INTER Plus II Tank Mixtures for Corn – Additional Weeds Controlled and Special Instructions

	<b>INTER Plus II + Aatrex and/or Princep (Preplant Surface, PPI, PRE)</b>	<b>INTER Plus II + Aatrex (Post)</b>
<b>Section</b>	D1	D2
<b>Special Mixing Instructions</b>		
<b>Comments</b>	3,4,5, 6,9,10	3,4,5,6
Browntop panicum	•	
Cocklebur	•	o
Common purslane	•	
Hairy nightshade	•	
Jimsonweed		o
Kochia		•
Lambsquarters	•	•
Morningglory	•	o
Mustard		•
Pigweed		
Prickly sida		•
Ragweed	•	•
Smartweed	•	•
Velvetleaf	•	•

**Chart 1 Continued: INTER Plus II Tank Mixtures for Corn - Additional Weeds Controlled and Special Instructions**

	INTER Plus II + Banvel (Field Corn)	INTER Plus II + AAtrex + Lorox	INTER Plus II + AAtrex or Princep + Prowl	INTER Plus II + Marksman	INTER Plus II + Broadstrike + Inter 8E II
Section	D5	D6	D7	D11	D12
Special Mixing Instructions			2		
Comments		3,4,5 6,8	3,4,5,6	9	9
Browntop panicum		•	•		
Cocklebur	o	•	•	•	o•
Common purslane		•	•	•	•
Hairy nightshade		•	•	•	•
Jimsonweed	o			•	o•
Kochia				•	•
Lambsquarters	•	•	•	•	•
Morningglory	o	•	•	•	o•
Mustard				•	•
Pigweed		•	•	•	•
Prickly Sida				•	•
Ragweed	•	•	•	•	o•
Smartweed	•	•	•	•	•
Velvetleaf	o	•	•	•	•

• = control; o = partial control; o• = partial to full control depending on ratio of products used or on weed population

**Comments**

**1. Special Mixing Instructions for INTER Plus II + AAtrex or Princep and Prowl**

(1) Fill the spray tank 1/4 full with water or fluid fertilizer and start agitation. (2) To aid compatibility, add a compatibility agent, such as Unite or X-77 • , at 4 pts./100 gals. of spray mixture. (3) Then add the AAtrex or Princep and allow it to become dispersed. (4) Then add INTER Plus II and Prowl 4E. (5) Add the rest of the water.

**2. Although a single formulation for AAtrex or Princep is listed in the rate tables, other formulations may be substituted, using the following formula:**

- 1 lb. of AAtrex Nine-O® or Princep Caliber 90® = 1.8 pts. of AAtrex 4L or Princep 4L.

**3.** Although directions specify AAtrex formulations in tank mixture with INTER Plus II, other brands of atrazine may be used. Follow the rates, recommendations, and limitations on the atrazine label.

**4.** See additional mixing instructions on the AAtrex label.

**5. Precaution:** Do not exceed a total of 2.5 lbs. a.i. of atrazine per acre per year. However, certain states may have established rate limitations for atrazine within specific geographical areas. Consult your state lead pesticide control agency for additional information. It is a violation of this label to deviate from state use regulations.

**6.** Other formulations of Lorox can be used: 1 lb. of Lorox DF = 1 pt. of Lorox L.

**7.** In Minimum-Tillage and No-Tillage systems, mix with Gramoxone Extra for control of most emerged annual weeds and suppression of perennial weeds; or with Landmaster BW for suppression of emerged field bindweed and control or suppression of annual weeds; or with Roundup for control of most emerged annual and perennial weeds.

**8.** Refer to label section A7 for specific directions for 2,4-D or Banvel burndown combinations in Minimum-Tillage and No-Tillage systems.

INTER Plus II in any tank mixture for corn may be applied in water or fluid fertilizer, except as noted.

**Notes:** (1) For all applications to corn, do not graze or feed forage from treated areas for 30 days following application, or possible illegal residues may result. (2) When applying INTER Plus II in tank mixture with AAtrex, do not exceed a total of 2.5 lbs. a.i. of atrazine per acre per year. (3) Refer to Section A, **Corn (All Types) - INTER Plus II Alone**, Note 3 for recommended sequential postemergence treatments if escape weeds develop. (4) In corn, INTER Plus II may be used up to 2.0 pts. / A in combinations on any soil for extended residual control and where severe stands of problem weeds are expected.

**1. Tank Mixture with AAtrex or Princep, or AAtrex + Princep - Preplant Surface, Preplant Incorporated, or Preemergence**

In addition to the weeds controlled by INTER Plus II alone, INTER Plus II+ AAtrex or Princep, or INTER Plus II+ AAtrex + Princep, applied preplant surface, preplant incorporated, or preemergence, also controls the following weeds: browntop panicum, cocklebur, common purslane, hairy nightshade, lambsquarters, morningglory, ragweed, smartweed, and velvetleaf.

Apply INTER Plus II+ AAtrex or Princep, or INTER Plus II+ AAtrex + Princep either preplant surface, preplant incorporated, or

preemergence.

**Preplant Surface-Applied:** Follow instructions for use of INTER Plus II alone under **Application Procedures** and under application instructions for INTER Plus II alone on corn. Apply INTER Plus II+ AAtrex or Princep, or INTER Plus II+ AAtrex + Princep on *medium soils* (1.67 pts./A of INTER Plus II+ 3.2 - 4 pts./A of AAtrex 4L or Princep 4L, or AAtrex 4L + Princep 4L combined) and on *fine soils* (1.67 - 2.0 pts./A of INTER Plus II+ 4 - 5 pts./A of AAtrex 4L or Princep 4L, or AAtrex 4L + Princep 4L combined) in minimum-tillage and no-tillage systems in CO, IA, IL, IN, KS, KY, MN, MO, MT, ND, NE, SD, TN, WI, and WY. Apply the tank mixtures as a split or single treatment in those states and as indicated in the **INTER Plus II Alone - Preplant Surface-Applied** section of the label for corn. On *coarse soils*, apply 1.33 pts./A of INTER Plus II and 3.2 pts./A of AAtrex 4L or Princep 4L, or AAtrex 4L + Princep 4L combined.

**Preplant Incorporated or Preemergence:** Follow instructions for use of INTER Plus II alone under **Application Procedures**. Apply INTER Plus II+ AAtrex or Princep, or INTER Plus II+ AAtrex + Princep, using the appropriate rates from **Table 1**.

**Note:** Do not apply more than the labeled rate for a given soil texture per year, either as a split or single treatment, or illegal residues may result.

**Shattercane, Wild Proso Millet, Woolly Cupgrass, and Eclipta - Partial Control**

For more consistent partial control of shattercane, wild proso millet, woolly cupgrass, or eclipta, where INTER Plus II is applied in tank mixture or sequentially with other registered corn herbicides apply 2.0 - 2.33 pts. as a single application, or the following applications may be made:

1. Apply 1.0 - 1.33 pts./A of INTER Plus II+ 2 lbs. a.i./A of AAtrex or Princep preplant incorporated, followed by 1.0 - 1.33 pts./A of INTER Plus II preemergence. Make the preemergence application during or after planting, but before weeds and corn emerge.
2. Apply INTER Plus II at 1.33 pts./A alone or in tank mix combination with up to 2 lbs. a.i./A of AAtrex or Princep, preplant incorporated. Do not exceed the total rate of triazine herbicide recommended for corn grown on a given soil texture. Follow with a post-directed application of Evik® 80W at 2.5 lbs./A. Refer to the Evik 80W label for specific directions for the post-directed application.
3. Apply Eradicane® or Sutan® (or equivalent EPTC or butylate formulations) at labeled rates preplant incorporated, followed by a preemergence application of INTER Plus II at 1.0 - 1.33 pts./A. Do not use Eradicane or Sutan on soils where rapid degradation has been shown to occur. Make the preemergence application during or after planting, but before weeds and corn emerge.

*Precaution: When following the application regimes in numbers 1-4 above, a shallow cultivation may be needed after the preemergence or postemergence application to help control any late emerging shattercane or wild proso millet plants.*

**Table 1: INTER Plus II+ AAtrex or Princep, or INTER Plus II+ AAtrex + Princep, Preplant Incorporated or Preemergence - Corn (All Types)**

Soil Texture	Broadcast Rates Per Acre			
	Less than 3% Organic matter		3% Organic Matter or Greater	
	INTER Plus II + AAtrex Nine-O* OR Princep Caliber 90*	INTER Plus II + AAtrex Nine-O** + Princep Caliber 90**	INTER Plus II + AAtrex Nine-O* OR Princep Caliber 90*	INTER Plus II + AAtrex Nine-O** + Princep Caliber 90**
COARSE	0.85 - 1.0 pts. + 1.1 - 2.2 lbs.	0.85 - 1.0 pts. + 0.6 - 1.1 lbs. + 0.6 - 1.1 lbs.	1.0 pts. + 1.3 - 2.2 lbs.	1.0 pts. + 0.7 - 1.1 lbs. + 0.7 - 1.1 lbs.
MEDIUM	1.0 - 1.33 pts. + 1.3 - 2.2 lbs.	1.0 - 1.33 pts. + 0.7 - 1.1 lbs. + 0.7 - 1.1 lbs.	1.33 pts. + 1.8 - 2.2 lbs.	1.33 pts. + 0.9 - 1.1 lbs. + 0.9 - 1.1 lbs.
FINE	1.33 pts. + 1.8 - 2.2 lbs.	1.33 pts. + 0.9 - 1.1 lbs. + 0.9 - 1.1 lbs.	1.33 - 1.67 pts. + 1.8 - 2.2 lbs.***	1.33 - 1.67 pts. + 0.9 - 1.1 lbs.*** + 0.9 - 1.1 lbs.
Muck or Peat (soils with more than 20% organic matter)	DO NOT USE			

\*Use Princep in preference to AAtrex when heavy infestations of crabgrass or fall panicum are expected. On soils having between 6% and 20% organic matter, INTER Plus II may be used up to 2.33 pts./A in tank mix combination with 2.2 lbs./A of AAtrex Nine-O, or equivalent rates of AAtrex 4L. Refer to the AAtrex label for weeds controlled at this reduced rate.

\*\*When using the tank mixture of INTER Plus II+ AAtrex Nine-O + Princep Caliber 90, use equal rates of each as shown when heavy broadleaf weed infestations are expected. When heavy infestations of crabgrass or fall panicum are expected, use a 1:2 ratio of AAtrex + Princep instead of the 1:1 ratio given in Table 1. (Example: Total AAtrex Nine-O + Princep Caliber 90 = 1.2 lbs./A, use 0.4 lb. of AAtrex + 0.8 lb. of Princep, respectively.) Refer to Comment No. 3 following Chart 1 for AAtrex 4L and Princep 4L conversions.

\*\*\*For cocklebur, yellow nutsedge, and velvetleaf control on *fine-textured soils* above 3% organic matter, apply 2.25 lbs./A of AAtrex Nine-O, or equivalent rates of AAtrex 4L, or the same total amount of AAtrex + Princep with 1.33 - 1.67 pts./A of INTER Plus II.

**2. Tank Mixture with AAtrex - Postemergence**

**a. Weeds Controlled**

barnyardgrass (watergrass)		yellow foxtail	prickly sida
crabgrass	kochia	Jimsonweed	purslane
crowfootgrass		ragweed	smartweed
fall panicum		lambsquarters	velvetleaf
giant foxtail		mustard	
green foxtail		pigweed	

**b. Weeds Partially Controlled:** cocklebur, morningglory, yellow nutsedge

Apply 1.0 pts./A of INTER Plus II+ 1.3 lbs./A of AAtrex Nine-O\* on *coarse soils*, 1.33 pts./A of INTER Plus II+ 1.8 lbs./A of AAtrex Nine-O on *medium soils*, or 1.33 - 1.67 pts./A of INTER Plus II + 1.8 - 2.2 lbs./A\*\* of AAtrex Nine-O on *fine soils*. Apply this tank mixture before grass and broadleaf weeds pass the 2-leaf stage and before corn exceeds 5 inches in height. Application to weeds larger than the 2-leaf stage will generally result in unsatisfactory control.

**Lay-by:** Apply to corn plants not more than 12 inches tall. Applications to corn in excess of 5 inches should be directed to the base of the corn plants, whereas, applications to corn plants less than 5 inches tall may be made over the top. Occasionally, some corn leaf burn may result, but this should not affect later growth or yield. Do not apply this postemergence tank mixture in fluid fertilizer, or severe crop injury may occur.

\*When using AAtrex 4L, use equivalent rates. One lb. of AAtrex Nine-O equals 1.8 pts. of AAtrex 4L.

\*\*For better control of cocklebur, morningglory, velvetleaf, and yellow nutsedge on *fine-textured soils* above 3% organic matter, apply 2.2 lbs./A of AAtrex Nine-O, or equivalent rate of AAtrex 4L, with 1.33 - 1.67 pts./A of INTER Plus II.

Tank mixtures of INTER Plus II+ AAtrex may be applied following use of any registered preplant surface-applied, preplant incorporated, or preemergence corn herbicide, including INTER Plus II+ AAtrex.

**Note:** The total INTER Plus II rate should not exceed 4 pts., nor the AAtrex rate more than 2.5 lbs. a.i./A during any one crop year, or illegal residues may result. Refer to the AAtrex label for geographic, soil-texture, and rotational restrictions.

**3. Tank Mixture with Banvel**

**Preemergence:** Use this tank mixture only on field corn which is flat-planted (no furrows) in CO, IA, IL, IN, KS, MN, NE, OH, SD, and WI.

In addition to the weeds controlled by INTER Plus II alone, INTER Plus II+ Banvel, applied preemergence, also controls lambsquarters, ragweed, smartweed, cocklebur\*, jimsonweed\*, morningglory\*, and velvetleaf\*.

\*Partially controlled.

Apply INTER Plus II+ Banvel preemergence. Broadcast 1 pt./A of Banvel with 1.33 pts./A of INTER Plus II on *medium soils*, or with 1.33 - 1.67 pts./A of INTER Plus II on *fine soils*. Do not apply on *coarse soils* or on soils with less than 2.5% organic matter. Apply this tank mixture to the soil surface at planting or after planting, but before corn emerges. Plant corn at least 1.5 inches deep and apply behind planting equipment, avoiding incorporation by the planter wheel or other seed covering device. Do not incorporate before corn emergence. If it is necessary to rotary hoe to break the soil crust, do not disturb the soil more than 1/2 inch deep.

*Precautions: (1) Avoid drift to sensitive non-target plants, such as soybeans, during application, or injury may occur. (2) Do not apply with aircraft.*

**Postemergence for Control of Pigweed (Mid-Atlantic states, including DE, MD, PA, VA, and WV):** Apply 1.0 - 1.50 pts. of INTER Plus II+ 0.5 - 1 pt./A of Banvel by ground equipment when pigweed plants are less than 3 inches tall and before corn exceeds 5 inches in height in a minimum of 20 gals. of spray per acre. Use the lower rate on coarse-textured and low organic matter soils. Use the higher rate on fine-textured and high organic matter soils.

Banvel SGF® and Clarity® may be used at equivalent lbs. of active ingredient per acre.

*Precautions: (1) Avoid drift to sensitive non-target plants, such as soybeans, during application, or injury may occur. (2) Do not apply with aircraft.*

**4. Tank Mixture with AAtrex and Lorox for Control of Lambsquarters and Pigweed**

For prolonged control of lambsquarters and pigweed in DE, MD, NJ, NY, PA, VA, and WV, INTER Plus II may be applied preemergence in tank mix combination with AAtrex + Lorox. Apply INTER Plus II and AAtrex according to the rates in Table 1 and Lorox according to the following rates.

Soil Texture	Broadcast Rate Per Acre
Sandy loam (1 - 3% organic matter)	0.87 lb. Lorox
Sandy loam (3 - 6% organic matter)	1.0 lb. Lorox
Medium- and fine-textured soils (1-6% organic matter)	1.0 lb. Lorox

Observe all directions for use, precautions, and limitations on the INTER Plus II, AAtrex, and Lorox labels when applying these products in tank mix combinations.

**5. Tank Mixture with AAtrex or Princep + Prowl for Prolonged Control of Lambsquarters and Pigweed in Field Corn Only (Northeast U.S., including MI, IN, KY, and States East of These)**

For prolonged control of lambsquarters and pigweed, in addition to a broad spectrum of annual broadleaf and grass weeds, INTER Plus II in tank mix combination with AAtrex\* or Princep + Prowl 4E may be applied after planting, but before corn or weeds emerge. Apply by ground equipment in a minimum of 10 gals. of water or 20 gals. of liquid fertilizer. Apply by air in a minimum of 5 gals. of water. Refer to Table 1 of this label for rates of INTER Plus II, AAtrex, or Princep to be applied. Apply Prowl 4E according to the following rates in Table 2.

\* Do not apply INTER Plus II in tank mix combination with AAtrex 80W + Prowl, as this combination is not compatible. Other AAtrex formulations may be used.

Mixing Instructions: See Comment No. 1 following Chart 1.

Table 2: Prowl 4E – Broadcast Rates Per Acre

Soil Texture	Percent Organic Matter in Soil		
	Less Than 1.5%	1.5 - 3%	Over 3%
COARSE	1.5 - 2.0 pts.	2.0 pts.	3.0 pts.
MEDIUM	2.0 pts.	3.0 pts.	3.0 pts.
FINE	2.0 pts.	3.0 pts.	3.0 pts.

Observe all directions for use, precautions, and limitations on the respective product labels when applying these products in tank mix combination. Refer to the Prowl 4E label for replanting instructions in the event of crop loss.

**6. Tank Mixture with AAtrex, or Princep, AAtrex + Princep, with Gramoxone Extra, Landmaster BW, or Roundup for**



## Minimum-Tillage or No-Tillage Systems

\*See Comment No. 1 following Chart 1 for special mixing instructions.

In minimum-tillage or no-tillage systems where corn is planted directly into a cover crop, stale seedbed, established sod, or previous crop residues, the contact herbicides Gramoxone Extra, Landmaster BW, or Roundup may be added to a tank mix of INTER Plus II+ AAtrex or Princep, or INTER Plus II+ AAtrex + Princep. See Comment No. 7 following Chart 1. The INTER Plus II+ AAtrex or Princep, or INTER Plus II+ AAtrex + Princep portion of the tank mixture provides preemergence control of the weeds listed on this label in the tank mixture section for INTER Plus II+ AAtrex or Princep, or INTER Plus II+ AAtrex + Princep – Preplant Surface, Preplant Incorporated, or Preemergence.

**Application:** Apply before, during, or after planting, but before the corn emerges, at the rates specified below. Add Gramoxone Extra, Landmaster BW, or Roundup at the following broadcast rates:

**Gramoxone Extra:** 1.5 - 2.0, 2.0 - 2.5, or 2.5 - 3.0 pts./A to 1 - 3 inches, 3 - 6 inches, or 6-inch tall weeds, respectively. Apply surfactant at 1 or 2 pts./100 gals. of spray mixture with 75% or greater or 50 - 74% nonionic active ingredient, respectively. This treatment will not control weeds taller than 6 inches.

**Note:** Do not apply combinations containing Gramoxone Extra in suspension-type liquid fertilizers, because the activity of paraquat will be reduced.

**Landmaster BW:** 27 - 54 oz./A depending on weed species and size. See the Landmaster BW label for weeds controlled, recommended rates for specific weeds, and other information concerning use.

**Roundup:** See the Roundup or Roundup RT label for weeds controlled, recommended rates, and other use directions.

Apply in 20 - 60 gals. of water or fluid fertilizer per acre with ground equipment.

On *coarse soils*, apply 1.0 pts./A of INTER Plus II with 1.3 lbs. of AAtrex Nine-O\* or Princep Caliber 90\*, or with 0.7 lb. of AAtrex Nine-O\*\* + 0.7 lb. of Princep Caliber 90\*\*. On *medium soils*, apply 1.33 pts./A of INTER Plus II with 1.8 lbs. of AAtrex Nine-O or Princep Caliber 90, or with 0.9 lb. of AAtrex Nine-O + 0.9 lb. of Princep Caliber 90. On *fine soils*\*\*\*, apply 1.33 - 1.67 pts./A of INTER Plus II with 1.8-2.2 lbs. of AAtrex Nine-O or Princep Caliber 90, or with 0.9 - 1.1 lbs. of AAtrex Nine-O + 0.9 - 1.1 lbs. of Princep Caliber 90.

\*Use Princep in preference to AAtrex when heavy infestations of crabgrass or fall panicum are expected.

\*\*When using the tank mixture of INTER Plus II + AAtrex Nine-O + Princep Caliber 90, use equal rates of AAtrex and Princep as shown when heavy broadleaf weed infestations are expected. When heavy infestations of crabgrass or fall panicum are expected, use a 1:2 ratio of AAtrex + Princep instead of the 1:1 ratio given. ( *Example:* Total AAtrex Nine-O + Princep Caliber 90 = 1.8 lbs./A, use 0.6 lb. of AAtrex + 1.2 lbs. of Princep, respectively.) Refer to Comment No. 2 following Chart 1 for AAtrex 4L and Princep 4L conversions.

\*\*\*For cocklebur, yellow nutsedge, and velvetleaf control on *fine-textured soils* above 3% organic matter, apply 2.25 lbs./A of AAtrex Nine-O, or equivalent rate of AAtrex 4L, or the same total amount of AAtrex + Princep, with 1.33 - 1.67 pts./A of INTER Plus II.

### 7. Tank Mixture with AAtrex; or AAtrex + 2,4-D; or AAtrex + 2,4-D + Banvel for Minimum-Tillage or No-Tillage Systems

In minimum-tillage or no-tillage systems where corn is planted directly into a cover crop, stale seedbed, established sod, or previous crop residues, INTER Plus II applied in combination with AAtrex will kill most emerged small annual weeds. Apply INTER Plus II+ AAtrex before, during, or after planting, but before corn emerges, according to the rates in Table 1.

Where heavy crop residues exist, add 0.8 - 1.6 pts./A of an appropriately labeled 3.8 lbs. a.i./gal. 2,4-D amine (such as Weedar 64, Weedar 64A, DMA-4 Herbicide, or Formula 40) to the spray tank last and apply in a minimum of 25 gals. of carrier per acre.

As carriers, nitrogen solutions and complete liquid fertilizers, applied before corn emergence, enhance burndown of existing weeds, and therefore, are recommended instead of water. Add X-77 surfactant at 1.0 - 2.0 qts./100 gals. of diluted spray, or another appropriate surfactant at its recommended rate, or add crop oil concentrate plus 28% liquid nitrogen (or equivalent). Apply before weeds exceed 3 inches in height. If alfalfa is present, add Banvel to the spray mixture at 0.33 - 0.5 pt./A and apply before alfalfa exceeds 6 inches in height.

For fields with existing sod grasses (e.g., bromegrass, orchardgrass, rye, or timothy), when existing weeds exceed 3 inches in height or when very dry conditions exist, add Gramoxone Extra at the rate of 2.5 pts./A in place of or in addition to 2,4-D, as indicated above. Do not apply Gramoxone Extra in suspension-type liquid fertilizer. Observe all directions for use, precautions, and limitations on the respective product labels when applying these products in tank mix combination.

### 8. Tank Mixture with Marksman in Conservation Tillage - Field and Silage Corn

In conservation tillage systems where corn is planted directly into a cover crop or previous crop residue, INTER Plus II+ Marksman will kill most emerged small annual weeds. Apply INTER Plus II+ Marksman before, during, or after planting, but before corn emergence on *medium* and *fine soils* with greater than 2.5% organic matter. For fields with existing vegetation exceeding 3 inches in height or when very dry conditions exist, add Gramoxone Extra at its standard rate. INTER Plus II+ Marksman may be applied postemergence to corn less than 3 inches tall and before weedy grasses exceed the 2-leaf stage. As carriers, nitrogen solutions and complete liquid fertilizers, applied before corn emergence, enhance burndown of existing weeds. Do not apply Gramoxone Extra in suspension-type liquid fertilizer or use on emerged corn.

Refer to the Marksman label and follow all directions, limitations, precautions, and information regarding application and use in corn.

### 9. Tank Mixture with Broadstrike + INTER 8E II

For preplant surface, preplant incorporated, or preemergence application where severe grass populations are expected on medium- or fine-textured soils with relatively high organic matter content, Broadstrike + INTER 8E II may be spiked with INTER Plus II for optimum performance. Refer to the Broadstrike + INTER 8E II label for its use rate and the amount of metolachlor active ingredient it contains. INTER Plus II may be added up to, but not to exceed, the maximum alone INTER Plus II label rate for the soil classification. Refer to the respective labels and follow all directions, limitations, and restrictions for each product.

## B. Cotton

### INTER Plus II Alone

**1. Application:** Apply INTER Plus II preemergence only in Area 1\* at the rate of 0.50 - 1.0 pts./A on sandy loams, .67- 1.33 pts./A on *medium soils*, or 1.0 - 1.33 pts./A on *fine soils*. Apply INTER Plus II preplant incorporated or preemergence in Area 2\*\* at 1.0 pts./A on sandy loams, 1.0 - 1.33 pts./A on medium soils, or 1.33 pts./A on fine soils. **Do not use on sands and loamy sand.**

\*Area 1 = AR, LA, MS, TN, and Bootheel of MO

\*\*Area 2 = NM, OK, and TX

**2. Preplant Incorporated (NM, OK, and TX Only):** Apply to the soil and incorporate into the top inch of soil immediately before planting, at planting, or after planting, but before crop or weeds emerge. Use a rolling cultivator or similar implement to uniformly incorporate not more than 1 inch deep. Use a preplant incorporated application if furrow irrigation is used or when a period of dry weather after application is expected. Where furrow irrigation is used, wet the top of the bed for best results. If the crop is to be planted on beds, apply and incorporate after bed formation. Cotton should be planted below the zone of incorporation; i.e., at least 1 inch on *fine soils* and 1.5 inches on *coarse* and *medium soils*. If incorporated prior to planting, use a planter that will result in a minimum of soil disturbance.

**Note:** For best control of yellow nutsedge and suppression of seedling johnsongrass, apply INTER Plus II preplant incorporated at the maximum rate for the soil texture, whether applied alone or mixed with Caparol 4L.

**3. Preemergence:** Apply to the soil surface at planting or after planting, but before weeds or crop emerge.

*Precautions: To avoid crop injury, (1) Do not apply Inter Plus II on sand or loamy sand soils, or in areas where water is likely to "pond" over the bed; (2) To avoid concentration in the seed furrow, do not make broadcast applications of Inter Plus II to cotton planted in furrows more than 2 inches deep. Band applications may be made to cotton planted in furrows deeper than 2 inches, but band width should not exceed the width of the bottom of the furrow, and (3) Do not apply on Taloka silt loam. (4) Do not use in Gaines County, TX.*

**Note:** To avoid possible illegal residues, do not graze or feed forage or fodder from cotton to livestock.

## B. Cotton

### INTER Plus II Combinations

#### 1. Tank Mixture with Caparol 4L

INTER Plus II tank mixtures with Caparol 4L may be applied in water or fluid fertilizer. When fluid fertilizer is used as a carrier for INTER Plus II, either alone or in combination with Caparol 4L, mix only the amount that will be sprayed in one operation. These mixtures should not be allowed to stand without agitation.

In addition to those weeds controlled by INTER Plus II alone, INTER Plus II+ Caparol 4L, applied preplant incorporated or preemergence, also controls the following weeds: junglerice, wild oats, annual morningglory, groundcherry, hairy nightshade, lambsquarters, malva, mustard, prickly sida (teaweed), purslane, ragweed, and shallow-germinating seedlings of cocklebur and coffeeweed.

**Preplant Incorporated or Preemergence:** Apply INTER Plus II+ Caparol 4L, either preplant incorporated or preemergence, using the appropriate rate from Table 3. Cotton should be planted below the zone of incorporation; i.e., at least 1.0 inch on *fine soils* and 1.5 inches on *coarse* and *medium soils*. If incorporated before planting, use a planter that will result in a minimum of soil disturbance.

**Table 3: INTER Plus II+ Caparol 4L - Cotton (NM, OK, TX)**

Use Areas	Soil Texture	Broadcast Rates Per Acre	
		INTER Plus II	Caparol 4L
ALL	Sand, loamy sand	DO NOT USE	
OK, and Blacklands and Gulf Coast of TX	Loams	0.85 - 1.33 pts.	2.4 pts.
	Clays	1.33 pts.	4.8 pts.
Rio Grande Valley of TX	Loams	0.85 - 1.33 pts.	3.2 pts.
	Clays	1.33 pts.	4.8 pts.

NM; High Plains, Rolling Plains, Edwards Plateau of TX, and Southwest TX	Sandy loam	0.85 - 1.0 pts.	1.6 pts.
	Loams	0.85 - 1.33 pts.	2.4 pts.
	Sandy clay loams	1.33 pts.	2.4 pts.
	Other clay soils	1.33 pts.	3.2 pts.

**Precautions:** (1) To avoid concentration in the seed furrow, do not make broadcast applications of INTER Plus II + Caparol 4L to cotton planted in furrows more than 2 inches deep. Band applications may be made to cotton planted in furrows deeper than 2 inches, but band width should not exceed the width of the bottom of the furrow. To avoid crop injury, (2) Do not apply on sand or loamy sand soils, or in areas where water is likely to "pond" over the bed; (3) Do not apply in cut areas of newly leveled fields, or in areas of excess salt. (4) Do not apply to glandless cotton varieties; and (5) Do not apply on Taloka silt loam. (6) Do not use in Gaines County, TX.

**Note:** To avoid possible illegal residues, do not graze or feed forage or fodder from cotton to livestock.

Refer to the Caparol 4L label for further instructions and limitations.

## 2. Tank Mixture with Cotoran DF

INTER Plus II may be applied in tank mixture with Cotoran DF preemergence for control of those weeds controlled by INTER Plus II alone and those as listed on the Cotoran DF label. This combination will also control spotted spurge, hyssop spurge, nodding spurge, and prostrate spurge. Apply to the soil surface at planting or after planting, but before weeds or crop emerge, using the appropriate rates from Table 4.

**Mixing Instructions:** Incompatibility may occur when tank mixing INTER Plus II and Cotoran DF. To help overcome this condition, fill the spray tank 1/4 full with water or fluid fertilizer and start agitation, add the Cotoran DF and allow it to become dispersed. Add X-77 at 0.5% volume/volume final spray (4 pts./100 gals.), then add the INTER Plus II and finally the rest of the water or fluid fertilizer. Agitate during mixing and application to maintain a uniform suspension.

**Table 4: INTER Plus II + Cotoran DF – Cotton**

Soil Texture	Broadcast Rates Per Acre		Cotoran DF***
	INTER Plus II (pts.)		
	Area 1	Area 2	lbs.
Sand, loamy sand		DO NOT USE	
Sandy loam	0.50 - 1.0	0.85 - 1.0	1.2
Loam, silt, silt loam	0.67 - 1.33	1.0 - 1.33	1.2 - 1.9
Fine soil	1.0 - 1.33	1.33	1.9 - 2.4

\*Area 1 = AR, LA, MS, Bootheel of MO, and TN

\*\*Area 2 = Eastern OK, Gulf Coast, Rio Grande Valley, and Eastern TX

\*\*\*When using Cotoran 4L, use equivalent rates. Multiply lbs. of Cotoran DF by 1.7 to get pts. of Cotoran 4L.

**Precautions:** (1) Do not apply INTER Plus II + Cotoran on sand or loamy sand soils, or in areas where water is likely to "pond" over the bed, or crop injury may occur. (2) To avoid concentration in the seed furrow, do not make broadcast applications of INTER Plus II + Cotoran to cotton planted in furrows more than 2 inches deep. Band applications may be made to cotton planted in furrows deeper than 2 inches, but band width should not exceed the width of the bottom of the furrow. (3) The use of Cotoran following the use of a systemic insecticide at planting may result in crop injury. (4) Do not use on Taloka silt loam, or crop injury may occur. (5) Do not use in Gaines County, TX.

Refer to the Cotoran labels for further instructions, precautions, and limitations.

**Note:** To avoid possible illegal residues, do not feed treated forage or gin trash to livestock, or graze treated areas.

## 3. Tank Mixture of INTER Plus II or INTER Plus II + Cotoran with Gramoxone Extra or Roundup for Minimum-Tillage or No-Tillage Systems

In minimum-tillage or no-tillage systems where cotton is planted directly into a cover crop, stale seedbed, or previous crop residues, the contact herbicides Gramoxone Extra or Roundup may be added to a tank mix of either INTER Plus II or INTER Plus II + Cotoran. When used as directed, the Gramoxone Extra portion of the tank mixture controls most emerged weeds and suppresses many perennial weeds. Roundup combinations will control emerged annual and perennial weeds when applied as directed on the Roundup label. The INTER Plus II and INTER Plus II + Cotoran portion of the tank mixture provides preemergence control of the weeds listed on this label in the INTER Plus II and INTER Plus II + Cotoran sections, respectively.

Refer to the label of each product used in combination and observe the planting details, information regarding application, geographical restrictions, and all other precautions and limitations. Refer to **Mixing Instructions** under the **Tank Mixture with Cotoran DF** section.

**Application:** Apply before, during, or after planting, but before the cotton emerges, at the rates specified below. Apply INTER Plus II at 0.85 - 1.0 pts./A on sandy loams, medium-, and fine-textured soils. Refer to Table 4 for the Cotoran DF rates.

Add Gramoxone Extra or Roundup at the following broadcast rates:

**Gramoxone Extra:** 1.5 - 2.0, 2.0 - 2.5, or 2.5 - 3.0 pts./A to 1 - 3 inches, 3 - 6 inches, or 6-inch tall weeds, respectively. Apply

surfactant at 1 or 2 pts./100 gals. of spray mixture with 75% or greater or 50 - 74% nonionic active ingredient, respectively. This treatment will not control weeds taller than 6 inches.

**Note:** Do not apply combinations containing Gramoxone Extra in suspension-type liquid fertilizers, as the activity of paraquat will be reduced.

**Roundup:** See the Roundup label for weeds controlled, recommended rates, and other use directions.

Apply in 20-60 gals. of water or fluid fertilizer per acre with ground equipment.

**Precautions:** (1) If heavy rain occurs soon after application, crop injury may result, especially in poorly drained areas where water stands for several days, or where the seeding slit has not been properly closed. (2) Refer to the Cotoran labels and the Tank Mixture with Cotoran DF section of this label for further instructions, precautions, and limitations. (3) Do not use in Gaines County, TX.

### C. Peanuts

#### INTER Plus II Alone

Apply INTER Plus II, either preplant incorporated, postplant incorporated or preemergence using the appropriate rate specified below. **Preplant Incorporated or Preemergence:** Follow instructions for use of INTER Plus II alone under **Application Procedures**. **Postplant Incorporated:** Apply and shallowly incorporate INTER Plus II into the soil after planting, but before peanut germination. Incorporation depth and incorporating implements must be kept above the seed, or seed will be damaged.

Apply INTER Plus II alone, preplant incorporated, postplant incorporated or preemergence at a broadcast rate of 1.0 - 1.33 pts./A in the Southeast\* and 0.85 - 1.33 pts./A in NM, OK, and TX.

\*In the Southeast, use 1.33 - 2.0 pts./A and apply preemergence for partial control of Florida beggarweed.

**Notes:** (1) INTER Plus II alone may be applied as directed after any of the following preplant incorporated herbicides when used according to their label recommendations: Balan at 3 - 4 qts./A; Treflan E.C. at 1 pt./A; Vernam <sup>®</sup> at 2.33 - 3 pts./A; Sonalan at 1.25 - 3 pts./A; Pursuit at 0.25 pt./A; or Prowl at 1 - 2 pts./A. (2) Do not graze or feed peanut forage or fodder to livestock for 30 days following application, and (3) Do not apply within 90 days of harvest, or illegal residues may result.

### C. Peanuts

#### INTER Plus II Combinations

##### 1. Tank Mixture with Balan L.C.

INTER Plus II+ Balan tank mixture applied preplant incorporated controls those weeds listed under INTER Plus II Applied Alone and those weeds as listed on the Balan label.

Apply 1.0 - 1.33 pts./A of INTER Plus II+ 3 - 4 qts./A of Balan in a minimum of 10 gals. of spray volume per acre for ground application or in a minimum of 5.0 gals. of spray volume per acre for aerial application. Follow the recommended procedures for Balan on the Balan label for soil preparation and incorporation of this tank mix. Apply and incorporate INTER Plus II+ Balan up to 14 days prior to planting.

**Note:** Follow all restrictions and precautions on the Balan label.

##### 2. Tank Mixture or Sequentially with Pursuit

The tank mixture or sequential treatment of INTER Plus II and Pursuit controls all weeds controlled by INTER Plus II alone and by Pursuit alone. Refer to the INTER Plus II Applied Alone section for weeds controlled by INTER Plus II and to the Pursuit label for weeds controlled by Pursuit.

Refer to the respective labels for application methods, timing, rates, restrictions, and precautions; and use in accordance with the more restrictive label.

##### 3. Tank Mixture with Sonalan

The tank mixture controls all weeds controlled by INTER Plus II alone and by Sonalan alone. Refer to the INTER Plus II Applied Alone section for weeds controlled by INTER Plus II and to the Sonalan label for weeds controlled by Sonalan.

Apply INTER Plus II+ Sonalan preplant incorporated, using the appropriate rate from Table 5. Follow recommended soil preparation procedures for Sonalan. Refer to the Peanut Sonalan/INTER 8E II Tank Mixture label for incorporation specifications.

**Table 5: INTER Plus II+ Sonalan - Peanuts**

	Broadcast Rates Per Acre			
	Southeast		NM, OK, TX	
Soil Texture	INTER Plus II	Sonalan	INTER Plus II	Sonalan
COARSE	1.0 - 1.33 pts.	1.25 - 2 pts.	0.85 - 1.33 pts.	1.25 - 2 pts.
MEDIUM	1.0 - 1.33 pts.	1.75 - 2.5 pts.	0.85 - 1.33 pts.	1.75 - 2.5 pts.

FINE	1.0 - 1.33 pts.	2.25 - 3 pts.	0.85 - 1.33 pts.	2.25 - 3 pts.
------	-----------------	---------------	------------------	---------------

**Note:** Follow all use directions, limitations, precautions, and information regarding application to peanuts on the INTER Plus II and Sonalan labels.

#### 4. Tank Mixture with Prowl

INTER Plus II+ Prowl applied preplant incorporated controls all weeds controlled by INTER Plus II alone plus Texas panicum, field sandbur, johnsongrass from seed, lambsquarters, kochia, annual spurge, and other species on the Prowl label. Apply INTER Plus II+ Prowl by ground or by aerial equipment within 14 days before planting. Incorporate into the top 1 - 2 inches of soil before planting and within 7 days of application, using a finishing disk or similar implement capable of providing uniform incorporation. If peanuts will be planted on beds, apply and incorporate after bed formation. Refer to the Incorporation instructions of the respective labels for additional directions.

Apply INTER Plus II+ Prowl preplant incorporated, using the appropriate rates from Table 6.

**Table 6: INTER Plus II + Prowl - Peanuts**

Soil Texture	Broadcast Rates Per Acre	
	NM, OK, TX	Other Peanut Growing States
	INTER Plus II + Prowl	INTER Plus II + Prowl
Sand, loamy sand	0.85 + 1 - 1.5 pts	1.0 - 1.33 + 1.5 - 2 pts.
Sandy loam	0.85 - 1.0 + 1 - 1.5 pts.	1.0 - 1.33 + 1.5 - 2 pts.
Fine soil	1.33 + 1 - 1.5 pts.	1.33 + 1.5 - 2 pts.

**Note:** Follow all use directions, limitations, precautions, and information regarding application to peanuts on the INTER Plus II and Prowl labels.

#### 5. Sequentially with Storm

Apply INTER Plus II according to the directions for INTER Plus II Alone and follow with a postemergence treatment of Storm as specified on its label for the control of weeds listed on the INTER Plus II label and on the Storm label. Refer to the Peanuts - INTER Plus II Alone section and to the Storm label and follow all directions, limitations, and restrictions for each product.

#### 6. Multiple Applications

Where weed pressure is heavy or where species difficult to control are expected, INTER Plus II is most effective when used as follows:

##### A. Southeast Only (AL, FL, GA, NC, SC, VA)

**1st Application:** Apply INTER Plus II preplant incorporated as directed under Peanuts - INTER Plus II Alone or apply INTER Plus II+ Balan preplant incorporated as directed previously in this section. Refer to the respective section for weeds controlled.

**2nd Application:** Apply INTER Plus II any time from preemergence to before "ground cracking" at 1.0 - 2.0 pts./A for extended control of weeds not yet emerged. Do not use INTER Plus II after peanut emergence. If peanuts have emerged, use INTER 8E II according to its label: Peanuts - INTER 8E II Combinations - Multiple Applications.

**3rd Application:** Do not use INTER Plus II. Apply INTER 8E II at lay-by as directed under Peanuts - INTER 8E II Alone section of the INTER 8E II label.

**Notes:** (1) Do not apply more than the equivalent of 4 pts. of metolachlor active ingredient per acre during any one year, or illegal residues may result. This includes INTER 8E II and INTER Plus II, but do not use INTER Plus II after peanuts have emerged. (2) Do not graze or feed peanut forage or fodder to livestock for 30 days following application, and (3) Do not apply within 90 days of harvest, or illegal residues may result.

##### B. Southwest Only (NM, OK, TX)

**1st Application:** Apply INTER Plus II preplant incorporated or preemergence to before "ground cracking" as directed under Peanuts - INTER Plus II Alone or apply INTER Plus II + Balan preplant incorporated as directed previously in this section. Do not use INTER Plus II after peanut emergence. If peanuts have emerged, use INTER 8E II according to its label.

**2nd Application:** Do not use INTER Plus II. Apply INTER 8E II at lay-by as directed under Peanuts - INTER 8E II Alone on that label. Use only when late germinating weeds are expected to be a problem. Refer to the INTER 8E II Applied Alone section for a list of weeds controlled.

**Notes:** (1) Do not apply more than the equivalent of 2.6 pts. of metolochlor active ingredient per acre during any one year, or illegal residues may result. This includes INTER 8E II and INTER Plus II, but do not use INTER Plus II after peanuts have emerged. (2) Do not graze or feed peanut forage or fodder to livestock for 30 days following application, and (3) Do not apply within 90 days of harvest, or illegal residues may result.

#### D. Pod Crops

## INTER Plus II Alone

Pod crops, including garbanzo, great northern beans, kidney beans, lima beans, mung beans, navy beans, peas (English\*; southern peas, such as blackeye, pinkeye, crowder, etc.), pinto beans, snap beans (green, wax, string), and lupines (sweet, white, white sweet, and grain).

### Fall Application

Apply after September 30 in ND, SD, MN, WI, and north of Route 30 in IA.

Apply after October 15 north of Route 91 in NE and south of Route 30 in IA.

Apply after October 31 north of Route 136 in IL.

In all locations, apply to crop stubble after harvest when the sustained soil temperature at a 4-inch depth is less than 55°F and falling. In minimum-till or no-tillage systems on soils having greater than 2.5% organic matter, use 1.65 - 2.0 pts./A on *medium-textured* and 2.0 pts./A on *fine-textured* soils. Do not apply to frozen ground. A tillage operation may precede the application. A fall and/or a spring tillage may follow application, but do not exceed an incorporation depth greater than 2-3 inches. Note: If a spring application is made, the total rate of the fall plus spring applications must not exceed the maximum total rate for pod crops, or illegal residues may result.

### Spring Application

Apply INTER Plus II, either preplant incorporated or preemergence, using the appropriate rate specified below. **Preplant Incorporated or Preemergence:** Follow instructions for use of INTER Plus II alone under **Application Procedures**. On *coarse soils* with less than 3% organic matter, apply 1.0 - 1.33 pts./A of INTER Plus II or 1.33 pts./A if organic matter is 3% or greater. On *medium soils*, apply 1.33 - 1.67 pts./A of INTER Plus II. On *fine soils*, apply 1.33 - 1.67 pts./A of INTER Plus II if organic matter content is less than 3%, or 1.67 - 2 pts./A if organic matter content is 3% or greater.

\*On English peas, use only preemergence applications. Do not use on English peas in northeastern U.S., or injury may occur.

**Note:** To avoid possible illegal residues, (1) Do not cut for hay within 120 days following an INTER Plus II application, and (2) Do not apply more than 2.0 pts./A of INTER Plus II during any one crop year, depending on soil texture.

## D. Pod Crops

### INTER Plus II Combinations

**Note:** When applying INTER Plus II in combination on pod crops, do not cut for hay within 120 days following application, or illegal residues may result.

#### 1. Tank Mixture and Sequential Applications with Eptam - Beans (Green or Dry)

This mixture controls all weeds controlled by INTER Plus II alone and by Eptam alone. Refer to the **INTER Plus II Applied Alone** section of this label for weeds controlled by INTER Plus II alone and to the Eptam label for weeds controlled by Eptam.

**Preplant Incorporated:** Follow instructions for use of INTER Plus II alone under **Application Procedures**. **Sequential:** Apply Eptam alone preplant incorporated, as specified on that label. Follow with a preemergence application of INTER Plus II, at rates specified for INTER Plus II alone, during planting (behind the planter) or after planting, but before the weeds or crop emerge.

Refer to the **General Information** section of this label and to the Eptam label for weather, cultural practices, and all other precautions and limitations that affect performance of these products.

Apply 2.5 - 4.5 pts./A of Eptam 7E\* with INTER Plus II as specified. On *coarse soils*, apply 0.85 pts./A of INTER Plus II if organic matter content is less than 3%, or 1.0 pts./A if organic matter content is 3% or greater. On *medium soils*, apply 1.0 pts./A of INTER Plus II if organic matter content is less than 3%, or 1.33 pts./A if organic matter content is 3% or greater. On *fine soils*, apply 1.33 pts./A of INTER Plus II if organic matter is less than 3%, or 1.33 - 1.67 pts./A if organic matter is 3% or greater.

\*Refer to the Eptam label for rate limitations depending on geographical area, and for species and varietal restrictions.

*Precaution: Do not exceed 3.5 pts./A of Eptam 7E on small white beans or green beans grown on coarse-textured soils.*

#### 2. Tank Mixture with Treflan - Beans (Dry - Kidney, Navy, Pinto, etc.; Lima; and Snap)

INTER Plus II+ Treflan tank mix applied preplant incorporated controls those weeds listed under **INTER Plus II Applied Alone** and those weeds listed for Treflan alone on the Treflan label. INTER Plus II+ Treflan may be applied by ground or by aerial equipment and incorporated up to 14 days prior to planting. Follow the recommended procedures on this label and on the respective Treflan label using equipment that provides uniform 2-inch incorporation.

Apply INTER Plus II+ Treflan tank mix using the appropriate INTER Plus II rate specified for INTER Plus II alone, and the Treflan rate from the Dry Beans, and the Lima and Snap Beans sections of the respective Treflan label. Choose the product rate for the specific soil texture, organic matter classification and weed species expected.

**Note:** Follow all restrictions and precautions on the respective Treflan label and in the **Pod Crops - INTER Plus II Alone** section of this label.

## E. Potatoes

### INTER Plus II Alone

Apply INTER Plus II, either incorporated or preemergence, according to directions specified below for control of weeds listed under the **General Information** section. Within a rate range, use the lower rate on soils relatively coarse-textured or low in organic matter;

use the higher rate on soils relatively fine-textured or high in organic matter. Effectiveness will be reduced if later cultural practices expose untreated soil.

**Incorporated:** Apply INTER Plus II at 1.0 - 2.0 pts./A to the soil and incorporate into the top 3 inches before planting, using a finishing disk, harrow, rolling cultivator, or similar implement. Planting and later cultural practices should not bring untreated soil to the surface. Postplant incorporated application may be made any time after planting to drag-off, but before potato emergence. Use an implement that evenly distributes INTER Plus II in the top 2 inches of soil. Do not damage potato seed pieces or sprouts with incorporation equipment.

**Preemergence:** Apply INTER Plus II at 1.0 - 2.0 pts./A, either after planting as a preemergence, delayed preemergence, after drag-off or hilling treatment, but before weeds emerge. Up to 2.75 pts./A of INTER Plus II alone may be used where soil organic matter is between 6% and 20%.

*Precautions: (1) Do not use on muck or peat soils. If cool, wet soil conditions occur after application, INTER Plus II may delay maturity and/or reduce yield of Superior and other early maturing potato varieties. To avoid crop injury, (2) Do not use on sweet potatoes or yams; (3) Do not apply both as a preemergence and an incorporated treatment; and (4) Do not use in Kern County, CA.*

**Note:** Potatoes treated with INTER Plus II should not be harvested within 60 days after the at-planting to drag-off application, or illegal residues may result.

## E. Potatoes

### INTER Plus II Combinations

#### 1. Tank Mixture with Sencor or Lexone

In addition to those weeds controlled by INTER Plus II alone, INTER Plus II applied in tank mix combination with, or sequentially with, any of the registered Sencor or Lexone formulations, also controls the following broadleaf weeds: cocklebur\*, hairy nightshade\*, hemp sesbania, jimsonweed\*, lambsquarters, prickly sida, ragweed, smartweed, velvetleaf, Venice mallow, and wild mustard.

\*Partially controlled.

INTER Plus II at 1.0 - 2.0 pts./A plus the labeled Sencor/Lexone use rate may be used preemergence. Apply 1.0 - 1.33 pts./A of INTER Plus II on coarse soils and 1.33 - 2.0 pts./A on other soil textures. Within this rate range, use the lower rate on soils relatively coarse-textured or low in organic matter; use the higher rate on soils relatively fine-textured or high in organic matter. Effectiveness will be reduced if later cultural practices expose untreated soil. INTER Plus II will not control emerged weeds.

Refer to the Sencor or Lexone labels for precautionary statements, restrictions, application information, and weeds controlled.

*Precautions: (1) Do not use INTER Plus II + Sencor or Lexone on potatoes in Kern County, CA. (2) Do not apply to sweet potatoes or yams. (3) Do not use this tank mixture on muck or peat soils.*

**Notes:** Potatoes treated with INTER Plus II in tank mixture with Sencor or Lexone cannot be harvested within 60 days after application, or illegal residues may result.

#### 2. INTER Plus II+ Lorox Tank Mixture (East of Rocky Mountains)

INTER Plus II may be applied in a tank-mix combination with any of the registered Lorox formulations as a preemergence broadcast application to potatoes. Apply to the soil surface after planting and before emergence of the crop or after final drag-off according to the rates specified in Table 7.

Table 7: INTER Plus II+ Lorox - Potatoes (East of Rocky Mountains)

Soil Texture	Broadcast Rates Per Acre			
	1% to Less Than 3% Organic Matter		3 - 5 % Organic Matter	
	INTER Plus II	Lorox*	INTER Plus II	Lorox*
<b>COARSE</b> Sandy loam	1.0 pts	1 - 1.5 lbs	1.33 pts	1.5 - 2 lbs
<b>MEDIUM</b> Loam, silt loam silt	1.33 pts	1.5 - 2 lbs	1.67 - 2.0 pts	2 - 2.5 lbs

\*When using Lorox L or Lorox DF use equivalent rates. One pt. of Lorox L equals 1 lb. of Lorox DF

*Precautions: To avoid crop injury, (1) Do not use on sands or loamy sands, and (2) Do not incorporate or spray over the top of emerged potatoes.*

Refer to the **General Information** section of this label and to the Lorox label for precautionary statements, restrictions, application information, and weeds controlled.

### 3. Tank Mixture with Prowl 4E

In addition to the weeds controlled by INTER Plus II alone, this tank mixture with Prowl 4E controls such problem species as kochia, lambsquarters, purslane, annual spurge, stinging nettle, and others specified on the Prowl 4E Alone label. Apply INTER Plus II+ Prowl 4E preemergence or preemergence incorporated, according to the specific directions on the Prowl 4E label, using the rates in Table 8.

**Table 8: INTER Plus II+ Prowl 4E - Potatoes**

	Broadcast Rates Per Acre	
	Less Than 3% Organic Matter	More Than 3% Organic Matter
<b>Soil Texture</b>	<b>INTER Plus II+ Prowl 4E*</b>	<b>INTER Plus II+ Prowl 4E*</b>
<b>COARSE</b>	1.0 - 1.33 pts. + 1 - 1.5 pts.	1.0 - 1.33 pts. + 1 - 1.5 pts.
<b>MEDIUM</b>	1.33 pts + 1.5 - 2 pts.	1.33 - 1.67 pts. + 2 - 3 pts.
<b>FINE</b>	1.33 - 1.67 pts. + 2 - 3 pts.	1.67 - 2.0 pts. + 3 pts.

\*When using other formulations of Prowl, use equivalent rates of active ingredient.

Refer to the INTER Plus II and Prowl 4E labels and observe all directions, timings, limitations, precautions, and restrictions concerning the use of these products on potatoes and follow the most restrictive.

### 4. Tank Mixture with Prowl 4E + Eptam

In addition to the weeds controlled by INTER Plus II alone, this tank mixture will control those species on the Prowl 4E and Eptam labels. Refer to the INTER Plus II+ Prowl 4E labels for rates of those products and add Eptam 7E at 3.5-7.0 pts./A, depending on geographical area. Refer to the respective INTER Plus II, Prowl 4E, and Eptam labels and observe all directions, limitations, precautions, and restrictions concerning the use of these products on potatoes and follow the most restrictive.

### F. Safflower

#### INTER Plus II Alone

**Preplant Incorporated or Preemergence:** Follow instructions for use of INTER Plus II alone under **Application Procedures**. **On coarse soils**, apply 1.0 - 1.33 pts./A of INTER Plus II if organic matter content is less than 3%, or 1.33 pts./A if organic matter is 3% or greater. **On medium soils**, apply 1.33 - 1.67 pts./A of INTER Plus II. **On fine soils**, apply 1.33 - 1.67 pts./A of INTER Plus II if organic matter content is less than 3%, or 1.67 - 2.0 pts./A if organic matter content is 3% or greater.

### G. Grain or Forage Sorghum (Seed Treated with Concep or Screen)

#### INTER Plus II Alone

Apply INTER Plus II, either preplant surface, preplant incorporated, or preemergence, using the appropriate rate specified below. Apply INTER Plus II alone only when the sorghum seed has been properly treated by the seed company with Concep or Screen. **Preplant Surface-Applied:** Refer to instructions for use of INTER Plus II under **Application Procedures**. For minimum-tillage or no-tillage systems only, INTER Plus II may be applied up to 45 days before planting in CO, IA, IL, KS, MO, NE, and SD. Use only split applications for treatments made 30 - 45 days prior to planting, with 2/3 of the broadcast rate applied initially and the remaining 1/3 at planting. Apply 1.50 pts./A of INTER Plus II on *medium soils* or 1.67 pts./A on *fine soils*. Treatments less than 30 days prior to planting may be made either as a split or single application. Apply 1.33 pts./A of INTER Plus II on *coarse soils* not more than 2 weeks prior to planting. Under dry conditions, irrigation after application is recommended to move INTER Plus II into the soil.

**Preplant Incorporated or Preemergence:** Refer to instructions for use of INTER Plus II under **Application Procedures**. Broadcast 1.0 - 1.33 pts./A of INTER Plus II on *coarse soils*, 1.33 - 1.50 pts./A on *medium soils*, or 1.33 - 1.67 pts./A on *fine soils*.

*Precautions (1) If sorghum seed is not properly treated with Concep or Screen, INTER Plus II will severely injure the crop. (2) Under high soil moisture conditions prior to sorghum emergence, injury may occur following the use of INTER Plus II. The crop will normally outgrow this effect. (3) Do not use INTER Plus II on sorghum grown under dry mulch tillage or injury may occur. (4) Except for the split preplant surface treatment, do not make more than one application per year, or illegal residues may result.*

### G. Grain or Forage Sorghum (Seed Treated with Concep or Screen)

#### INTER Plus II Combinations

INTER Plus II tank mixtures with AAtrex may be applied in water or fluid fertilizer. Apply INTER Plus II in tank mixtures only when



the sorghum seed has been properly treated by the seed company with Concep or Screen.

**IMPORTANT: FOR TANK MIXTURES WITH AATREX (OR OTHER BRANDS OF ATRAZINE) –** If applying INTER Plus II in tank mixture with AAtrex, all the restrictions and rate limitations on the AAtrex label **must** be followed if more restrictive/protective than those on this label. In addition, if AAtrex is/must be applied at rates lower than those recommended on this label, broadleaf weed control may be affected. Refer to the AAtrex label for weeds controlled at the reduced rates.

*Precautions: (1) Applications of INTER Plus II + AAtrex on highly alkaline soils or on eroded areas where calcareous subsoils are exposed may cause sorghum injury. (2) If sorghum seed is not properly treated with Concep or Screen, INTER Plus II + AAtrex may severely injure the crop. (3) Under high soil moisture conditions prior to sorghum emergence, injury may occur following the use of INTER Plus II + AAtrex. The crop will normally outgrow this effect. (4) Do not use INTER Plus II + AAtrex on sorghum grown under dry mulch tillage, or injury may occur. (5) Except for the split preplant surface treatment, do not make more than one application per year, or illegal residues may result.*

#### **1. Tank Mixture with AAtrex**

In addition to the weeds controlled by INTER Plus II alone, INTER Plus II+ AAtrex also controls the following broadleaf weeds when applied either preplant surface, preplant incorporated, or preemergence: cocklebur, common purslane, hairy nightshade, lambsquarters, morningglory, ragweed, smartweed, and velvetleaf.

**Preplant Surface-Applied:** Refer to instructions for use of INTER Plus II under **Application Procedures**. For minimum-tillage or no-tillage systems only, INTER Plus II+ AAtrex may be applied up to 45 days prior to planting in IA, IL, eastern KS, MO, NE, and SD. Use only split applications for treatments made 30 - 45 days prior to planting, with 2/3 of the broadcast rate applied initially and the remaining 1/3 at planting. Apply 1.50 pts./A of INTER Plus II+ 1.7 - 2 lbs./A of AAtrex Nine-O\* on *medium soils* with 1.5% organic matter or greater. Apply 1.50 pts./A of INTER Plus II+ 1.7 - 2 lbs./A of AAtrex Nine-O on *fine soils* with less than 1.5% organic matter, or apply 1.67 pts./A of INTER Plus II+ 2 - 2.2 lbs./A of AAtrex Nine-O on *fine soils* with 1.5% organic matter or greater. Treatments less than 30 days prior to planting may be made either as a split or single application. Under dry conditions, irrigation after application is recommended to move INTER Plus II + AAtrex into the soil.

*Precautions: To avoid crop injury, (1) Do not use on coarse soils, and (2) Do not use on medium soils with less than 1.5% organic matter.*

**Preplant Incorporated or Preemergence:** Refer to instructions for use of INTER Plus II under **Application Procedures**. On *medium soils* with 1.5% organic matter or greater, apply 1.0 pts./A of INTER Plus II + 1.3 lbs./A of AAtrex Nine-O\*. On *fine soils* with less than 1.5% organic matter, apply 1.0 pts./A of INTER Plus II + 1.3 lbs./A of AAtrex Nine-O, on *fine soils* with 1.5% organic matter or greater, apply 1.2 - 1.33 pts./A of INTER Plus II + 1.6 - 1.8 lbs./A of AAtrex Nine-O.

\*When using AAtrex 4L, use equivalent rates. One lb. of AAtrex Nine-O equals 1.8 pts. of AAtrex 4L.

*Precautions: To avoid crop injury, (1) Do not use on coarse soils; (2) Do not use on medium soils with less than 1.5% organic matter; (3) Do not use in NM, OK, or TX, except in northeast OK and the TX Gulf Coast and Blacklands areas; and (4) Do not apply preplant incorporated in AZ or the Imperial Valley of CA.*

#### **2. Tank Mixture of INTER Plus II or INTER Plus II+ AAtrex, with Gramoxone Extra, Landmaster BW, or Roundup for Minimum-Tillage or No-Tillage Systems**

In minimum-tillage or no-tillage systems where sorghum (seed treated with Concep or Screen) is planted directly into a cover crop, stale seedbed, established sod, or previous crop residues, the contact herbicides Gramoxone Extra, Landmaster BW, or Roundup may be tank mixed with INTER Plus II or INTER Plus II + AAtrex. See Comment No. 7 following Chart 1. The INTER Plus II or INTER Plus II+ AAtrex portion of the tank mixture provides preemergence control of the weeds listed on this label under the respective sections.

Refer to the label of each product used in combination and observe the planting details, restrictions, and all other precautions and limitations.

**Application:** Apply before, during, or after planting, but before sorghum emerges, at the appropriate rates listed under Grain or Forage Sorghum - INTER Plus II Alone or - INTER Plus II Combinations - INTER Plus II + AAtrex, respectively. Add Gramoxone Extra, Landmaster BW, or Roundup at the following broadcast rates:

Gramoxone Extra: 1.5 - 2, 2 - 2.5, or 2.5 - 3 pts. / A to 1 - 3 inches, 3 - 6 inches, or 6-inch tall weeds, respectively. Apply surfactant at 1 or 2 pts. / 100 gals. of spray mixture with 75% or greater or 50 - 74% nonionic active ingredient, respectively. This treatment will not control weeds taller than 6 inches.

Landmaster BW: 27 - 54 oz. / A depending on weed species and size. See the Landmaster BW label for weeds controlled, recommended rates for specific weeds, and other information concerning use.

Roundup: See the Roundup or Roundup RT label for weeds controlled, recommended rates, and other use directions. Apply in a minimum of 20 gals. of water per acre with conventional spray equipment.

#### **H. Soybeans**

## INTER Plus II Alone

Apply INTER Plus II, either preplant surface-applied, preplant incorporated, or preemergence, using the appropriate rate specified below. **Preplant Surface-Applied, Preplant Incorporated, or Preemergence:** Follow instructions for use of INTER Plus II alone under **Application Procedures**.

### Preplant Surface-Applied

**A. Fall Application (Apply after September 30 in MN, ND, SD, WI and north of Route 30 in IA; Apply after October 15 north of Route 91 in NE and south of Route 30 in IA; Apply after October 31 north of Route 136 in IL):**

In all locations, apply to crop stubble after harvest when the sustained soil temperature at a 4-inch depth is less than 55 ° F and falling. In minimum-till or no-tillage systems on soils having greater than 2.5% organic matter, use 1.67 - 2.0 pts./A on *medium-textured* and 2.0 pts./A on *fine-textured* soils. Do not apply to frozen ground. A tillage operation may precede the application. A fall and/or a spring tillage may follow application, but do not exceed an incorporation depth greater than 2 - 3 inches. Minimize furrow and ridge formation in the tillage operations. **Note:** If a spring application is made, the total rate of the fall plus spring applications must not exceed the maximum total rate for soybeans, or illegal residues may result.

**B. Use on *medium* and *fine* soils** with minimum-tillage or no-tillage systems in CO, CT, DE, IA, IL, IN, KS, KY, MA, MD, ME, MI, MN, MO, MT, ND, NE, NH, NY, OH, PA, RI, SD, TN, VA, VT, WI, WV, and WY. Apply 2/3 the recommended rate of INTER Plus II (1.67 pts./A on *medium* soils and 2.0 pts./A on *fine* soils) as a split treatment 30 - 45 days prior to planting and the remainder at planting. Applications made less than 30 days before planting may be as either a split or single treatment. Apply 1.33 pts./A on *coarse* soils not more than 2 weeks prior to planting.

**Preplant Incorporated or Preemergence:** On *coarse* soils, apply 1.0 - 1.33 pts./A of INTER Plus II if organic matter content is less than 3%, or 1.33 pts./A if organic matter content is 3% or greater. On *medium* soils, apply 1.33 - 1.67 pts./A of INTER Plus II. On *fine* soils, apply 1.33 - 1.67 pts./A of INTER Plus II if organic matter content is less than 3%, or 1.67 - 2.0 pts./A if organic matter content is 3% or greater.

**Note:** On soybeans, INTER Plus II may be used up to 2.55 pts./A as a preplant surface-applied, preplant incorporated, or preemergence treatment on soils having an organic matter content between 6% and 20%. The total INTER Plus II rate applied to soybeans during any one crop should not exceed 2.55 pts./A.

## H. Soybeans

### INTER Plus II Combinations

Water or fluid fertilizer may be used as carrier for INTER Plus II in combination with Sencor, Lexone, Lorox, Lorox Plus, Gemini, Canopy, Preview, Pursuit, Scepter, Sonalan, or Command.

**Note:** For all of the following combinations, INTER Plus II may be used up to 2.3 pts./A on soils having an organic matter content between 6% and 20%. The total INTER Plus II rate applied to soybeans during any one crop year should not exceed 2.55 pts./A.

#### 1. Tank Mixture with Sencor or Lexone

In addition to those weeds controlled by INTER Plus II alone, INTER Plus II+ Sencor or Lexone, when applied as directed, also controls the following broadleaf weeds: cocklebur\*, hairy nightshade, hemp sesbania, jimsonweed\*, lambsquarters, prickly sida, ragweed, smartweed, velvetleaf, Venice mallow, and wild mustard.

\*Partially controlled.

Apply INTER Plus II and Sencor or Lexone preplant incorporated or preemergence, using the appropriate rates from Table 9. **Preplant Incorporated or Preemergence:** Follow instructions for use of INTER Plus II alone under **Application Procedures**.

**Sequential:** Apply INTER Plus II alone **Preplant Incorporated**, as specified in Table 9 for this tank mixture. Follow with a preemergence application of Sencor or Lexone during planting (behind the planter) or after planting, but before weeds or soybeans emerge.

Refer to the Sencor or Lexone labels for planting details and soybean variety restrictions.

Table 9: INTER Plus II+ Sencor or Lexone - Soybeans

Soil Texture**	Broadcast Rates Per Acre					
	0.5% to Less Than 3% Organic Matter			3% Organic Matter or Greater		
	INTER Plus II	+	Sencor / Lexone DF*	INTER Plus II	+	Sencor / Lexone DF*
<b>COARSE</b> Loamy sand (over 2% organic matter), sandy loam	0.85 - 1.0 pts	+	0.33 lb.	1.0 pts	+	0.5 lb.
<b>MEDIUM</b>	1.0 - 1.33 pts	+	0.5 lb.	1.33 pts	+	0.67 lb.***
<b>FINE</b>	1.33 pts.	+	0.67 lb.	1.33 - 1.67 pts.	+	0.67 lb.
<b>Mississippi Delta only</b> Silty clay, clay	1.33 pts.	+	1.0 lb.	1.33 - 1.67 pts.	+	1.0 lb.

Muck or Peat (soils with more than 20% organic matter)	DO NOT USE
--	------------

\*When using Sencor 4 or Lexone 4L, multiply lbs. of DF by 1.5 to get pts./A.

\*\*On all sand and on loamy sand with less than 2% organic matter, do not use this tank mixture preemergence or the sequential treatment. Do not use the tank mixture preplant incorporated on any sand, loamy sand, or sandy loam, or crop injury may occur.

\*\*\*Use 0.5 lb./A if applied preplant incorporated.

*Precautions: (1) Do not use the tank mix or sequential application on soil with less than 0.5% organic matter or on alkaline soil with a pH over 7.4, or crop injury may occur. (2) If heavy rain occurs soon after application, crop injury may result, especially in poorly drained areas where water stands for several days.*

## 2. Tank Mixture with Lorox

In addition to those weeds controlled by INTER Plus II alone, INTER Plus II+ Lorox, applied preemergence, also controls the following broadleaf weeds: cocklebur\*, jimsonweed\*, lambsquarters, morningglory\*, prickly sida, ragweed, smartweed, velvetleaf\*, Venice mallow, and wild mustard.

\*Partially controlled.

**Preemergence:** Apply during planting (behind planter) or after planting, but before weeds or soybeans emerge. Refer to the Lorox label for planting details. Apply the appropriate rates from Table 10.

*Precaution: Do not use on soil with less than 0.5% organic matter, or crop injury may occur.*

**Table 10: INTER Plus II+ Lorox - Soybeans**

Soil Texture*	Broadcast Rates Per Acre					
	0.5% to Less Than 3% Organic Matter			3% Organic Matter or Greater		
	INTER Plus II	+	Lorox DF***	INTER Plus II	+	Lorox DF***
COARSE	0.85 pts.	+	1 lb.	1.0 pts.	+	1 - 1.5 lbs.
MEDIUM	1.0 pts.	+	1 - 1.5 lbs.	1.33 pts.	+	1.5 - 2 lbs.
FINE	1.33 pts.	+	2 lbs.	1.33 - 1.67 pts.	+	2.5 - 3 lbs.
Muck or Peat (soils with more than 20% organic matter)	DO NOT USE					

\*Do not use on sand, gravelly soils, or exposed subsoils.

\*\*Do not use on loamy sand, except in the northeastern U.S. on loamy sand with over 1% organic matter.

\*\*\*When using Lorox L or Lorox DF, use equivalent rates. One pt. of Lorox L equals 1 lb. of Lorox DF.

## 3. Tank Mixture with Treflan

INTER Plus II+ Treflan tank mix applied preplant incorporated controls those weeds listed under the INTER Plus II Applied Alone section and those weeds listed for Treflan Alone on the Treflan label. INTER Plus II+ Treflan may be applied by ground or by aerial equipment and incorporated up to 14 days before planting. Follow the recommended procedures on the Treflan and INTER Plus II labels, using equipment that provides uniform 2-inch incorporation.

Apply INTER Plus II+ Treflan tank mix, using the appropriate rate from the Soybeans - INTER Plus II Alone section of this label and the Treflan Alone section of the Treflan label for the specific soil texture/organic matter classification and weed species expected.

To control DNA-resistant goosegrass\* and other species on the respective labels where the soil organic matter is 3% or less, apply the rate listed in Table 11.

**Table 11: INTER Plus II+ Treflan - Organic Matter Content Less Than 3%**

Soil Texture	Broadcast Rates Per Acre		
	INTER Plus II Organic Matter Less Than 3%	Treflan E.C.** Organic Matter	
		Less Than 2%	2 - 3 %
COARSE*	0.85 - 1.0 pts.	1 pt.	1.5 pts.
MEDIUM	1.0 pts.	1.5 pts.	1.5 pts.
FINE	1.33 pts.	2 pts.	2 pts.

\*Where a range of rates is given for INTER Plus II, use the minimum rate where DNA-resistant goosegrass is the predominant species.

\*\*When Treflan MTF or Treflan 5 is used, use comparable rates. Multiply pts. of Treflan E.C. by 1 for Treflan MTF and by 0.8 for Treflan 5.

**Note:** Follow all restrictions and precautions on the respective Treflan label and in the Soybeans – INTER Plus II Alone section of this label.

**4. Tank Mixture with Scepter**

This tank mixture controls all weeds controlled by INTER Plus II alone and by Scepter alone. Refer to the INTER Plus II Applied Alone section for weeds controlled by INTER Plus II and to the Scepter label for weeds controlled by Scepter. Refer to the Scepter label for geographical locations where this tank mixture may be applied.

Apply INTER Plus II+ Scepter preplant incorporated or preemergence, using rates in Table 12. Follow use directions under Application Instructions on the Scepter label. For preplant incorporated applications, apply and incorporate within 30 days before planting. Observe all other precautions and limitations on the Scepter labels.

**Table 12: INTER Plus II+ Scepter – Soybeans**

Soil Texture	Broadcast Rates Per Acre			
	Less Than 3% Organic Matter		3% or More Organic Matter	
	INTER Plus II	Scepter	INTER Plus II	Scepter
COARSE	0.85 pts.	0.67 pt.	1.0 pts.	0.67 pt.
MEDIUM	1.0 pts.	0.67 pt.	1.33 pts.	0.67 pt.
FINE	1.33 pts.	0.67 pt.	1.33 - 1.67* pts.	0.67 pt.
Muck or Peat (soils with more than 20% organic matter)	DO NOT USE			

\*Use the higher rate of INTER Plus II if heavy weed infestations are expected.

**Notes:** (1) Do not apply within 90 days of harvest, and (2) Do not graze or feed treated soybean forage, hay, or straw to livestock, or illegal residues may result.

**5. Tank Mixture with Lorox Plus**

This tank mixture controls all weeds controlled by INTER Plus II alone and by Lorox Plus alone. Refer to the INTER Plus II Applied Alone section for weeds controlled by INTER Plus II and to the Lorox Plus label for weeds controlled by Lorox Plus.

Apply INTER Plus II+ Lorox Plus preemergence after planting, but before soybeans emerge, using rates in Table 13.

**Note:** Follow all use directions, limitations, precautions, information regarding application to soybeans, and rotational restrictions on the INTER Plus II and Lorox Plus labels.

**Table 13: INTER Plus II+ Lorox Plus – Soybeans**

Soil Texture	Broadcast Rates Per Acre	
	0.5 - 3% Organic Matter	
	INTER Plus II	Lorox Plus (60DF)
COARSE	0.85 pts.	12 - 14 oz.
MEDIUM	1.0 pts.	14 - 16 oz.
FINE	1.33 pts.	16 - 18 oz.

*Precaution: Do not apply to sand, or to any soil with less than 0.5% organic matter, or to any soil with pH greater than 6.8*

**6. Tank Mixture with Gemini**

This tank mixture controls all weeds controlled by INTER Plus II alone and by Gemini alone. Refer to the INTER Plus II Applied Alone section for weeds controlled by INTER Plus II and to the Gemini label for weeds controlled by Gemini.

Apply INTER Plus II + Gemini preemergence after planting but before soybeans emerge, using rates in Table 14

**Note:** Follow all use directions, limitations, precautions, information regarding application to soybeans, and rotational restrictions on the INTER Plus II and Gemini labels.

**Table 14: INTER Plus II+ Gemini – Soybeans**

Soil Texture	Broadcast Rates Per Acre	
	0.5 - 3% Organic Matter	
	INTER Plus II	Gemini (60DF)
COARSE (Sandy loam only)	0.85 pts.	12 - 16 oz.
MEDIUM	1.0 pts.	16 - 20 oz.
FINE	1.33 pts.	20 - 24 oz.

*Precaution: Do not apply to sand or loamy sand, or to any soil with less than 0.5% organic matter, or to any soil with pH greater than 7.0, except as noted on the Gemini label.*

#### 7. Tank Mixture with Canopy

This tank mixture controls all weeds controlled by INTER Plus II alone and by Canopy alone. Refer to the INTER Plus II Applied Alone section for weeds controlled by INTER Plus II and to the Canopy label for weeds controlled by Canopy.

Apply preplant incorporated or preemergence, using the appropriate rates from Table 15. **Preplant Incorporated:** Apply within 2 weeks of planting. Uniformly incorporate into the top 1-2 inches of soil before planting soybeans. **Preemergence:** Apply after planting, but before soybeans emerge.

**Note:** Follow all use directions, varietal restrictions, limitations, precautions, information regarding application to soybeans, and rotational restrictions on the INTER Plus II and Canopy labels.

**Table 15: INTER Plus II+ Canopy - Soybeans**

Soil Texture	Broadcast Rates Per Acre			
	0.5 to Less Than 3% Organic Matter		3-5% Organic Matter	
	INTER Plus II	Canopy	INTER Plus II	Canopy
COARSE	0.85 pts.	6.0 - 10 oz.	1.0 pts.	6.0 - 10 oz.
MEDIUM	1.0 pts.	8.0 - 12 oz.	1.33 pts.	8.0 - 12 oz.
FINE	1.33 pts.	10 - 14 oz.	1.33 - 1.67 pts.	10 - 14 oz.

*Precaution: Do not apply to sand, or to any soil with less than 0.5% organic matter, or to any soil with pH greater than 7.0, except as noted on the Canopy label.*

#### 8. Tank Mixture with Preview

This tank mixture controls all weeds controlled by INTER Plus II alone and by Preview alone. Refer to the INTER Plus II Applied Alone section for weeds controlled by INTER Plus II and to the Preview label for weeds controlled by Preview.

Apply preplant incorporated or preemergence, using the appropriate rates from Table 16. **Preplant Incorporated:** Apply within 2 weeks of planting. Uniformly incorporate into the top 1 - 2 inches of soil before planting soybeans. **Preemergence:** Apply after planting, but before soybeans emerge.

**Note:** Follow all use directions, varietal restrictions, limitations, precautions, information regarding application to soybeans, and rotational restrictions on the INTER Plus II and Preview labels.

**Table 16: INTER Plus II+ Preview - Soybeans**

Soil Texture	Broadcast Rates Per Acre			
	0.5% to Less Than 3% Organic Matter		3 - 5 % Organic Matter	
	INTER Plus II	Preview 75DF	INTER Plus II	Preview 75DF
COARSE	0.85 pts.	6 oz.	1.0 pts.	7 oz.
MEDIUM	1.0 pts.	7 oz.	1.33 pts.	8 oz.
FINE	1.33 pts.	8 oz.	1.33 - 1.67 pts.	9 - 10 oz.

*Precaution: Do not apply to sand, or to any soil with less than 0.5% organic matter, or to any soil with pH greater than 6.8.*

#### 9. Tank Mixture with Command\*

This tank mixture controls all weeds controlled by INTER Plus II alone and by Command alone. Refer to the INTER Plus II Applied Alone section for weeds controlled by INTER Plus II and to the Command label for weeds controlled by Command.

Apply INTER Plus II+ Command preplant incorporated, using rates in Table 17. Follow all Command application instructions as to incorporation interval, geographical location, equipment operation, soil moisture conditions, etc.

**\*Note:** Before making applications, read and strictly follow all use directions, limitations, precautions, information regarding application to soybeans, and rotational restrictions on the INTER Plus II and Command labels.

**Table 17: INTER Plus II+ Command – Soybeans**

Soil Texture	Broadcast Rates Per Acre			
	INTER Plus II		Command 4E	
	0.5 - 3% Organic Matter	Greater Than 3% Organic Matter	Northern Area	Southern Area
<b>COARSE</b>	0.85 pts.	1.0 pts.	1.5 - 2 pts.	2 - 2.5 pts.
<b>MEDIUM</b>	1.0 pts.	1.33 pts.	1.5 - 2 pts.	2 - 2.5 pts.
<b>FINE</b>	1.33 pts.	1.33 - 1.67 pts.	1.5 - 2 pts.	2 - 2.5 pts.

**10. Tank Mixture with Sonalan**

This tank mixture controls all weeds controlled by INTER Plus II alone and by Sonalan alone. Refer to the **INTER Plus II Applied Alone** section for weeds controlled by INTER Plus II and to the Sonalan label for weeds controlled by Sonalan.

Apply INTER Plus II and Sonalan preplant incorporated, using the appropriate rates from Table 18.

**Preplant Incorporated:** Follow recommended soil preparation procedures for Sonalan. Refer to the Sonalan/ INTER 8E II Tank Mixture label for incorporation specifications.

**Sequential:** Apply Sonalan alone preplant incorporated as specified on the Sonalan label. Follow with a preemergence application of INTER Plus II during planting (behind the planter) or after planting, but before weeds or soybeans emerge.

**Table 18: INTER Plus II+ Sonalan – Soybeans**

Soil Texture	Broadcast Rates Per Acre			
	Less Than 3% Organic Matter		3% or More Organic Matter	
	INTER Plus II	Sonalan	INTER Plus II	Sonalan
<b>COARSE</b>	1.0 - 1.33 pts.	1.25 - 2 pts.	1.33 pts.	1.25 - 2 pts.
<b>MEDIUM*</b>	1.33 - 1.67 pts.	1.75 - 2.5 pts.	1.33 - 1.67 pts.	1.75 - 2.5 pts.
<b>FINE*</b>	1.33 - 1.67 pts.	2.25 - 3 pts.	1.67 - 2 pts.	2.25 - 3 pts.
Muck or Peat (soils with more than 20% organic matter)	DO NOT USE			

\*For eastern black nightshade on these soils, apply Sonalan at 3 pts./A on *medium* and 3.5 pts./A on *fine-textured* soils, and follow with 2 incorporation passes.

**Note:** Follow all use directions, limitations, precautions, and information regarding application to soybeans on the INTER Plus II and Sonalan labels.

**11. Tank Mixture with Pursuit**

This tank mixture controls all weeds controlled by INTER Plus II alone and by Pursuit alone. Refer to the **INTER Plus II Applied Alone** section for weeds controlled by INTER Plus II and to the Pursuit label for weeds controlled by Pursuit. Refer to the Pursuit label for geographical locations where this tank mixture may be applied.

Apply INTER Plus II+ Pursuit early preplant, preplant incorporated, or preemergence after planting, using rates in Table 19. Application can be made in water or liquid fertilizer. Follow all use directions under Soil Applications on the Pursuit label. For early preplant and preplant incorporated applications, apply within 30 days before planting.

**Note:** Follow all use directions, limitations, precautions, information regarding application to soybeans, and rotational restrictions on the INTER Plus II and Pursuit labels.

**Table 19: INTER Plus II+ Pursuit – Soybeans**

Soil Texture	Broadcast Rates Per Acre		
	Less Than 3% Organic Matter	3% or More Organic Matter	
	INTER Plus II	INTER Plus II	Pursuit
COARSE	0.85 pts.	1.0 pts.	0.25 pt.
MEDIUM	1.0 pts.	1.33 pts.	0.25 pt.
FINE	1.33 pts.	1.33 - 1.67 pts.	0.25 pt.

**Sequential:** Apply INTER Plus II early preplant, preplant incorporated, or preemergence after planting at 0.85 pts./A on coarse soils and 1.0 pts./A on medium- and fine-textured soils. Follow with a sequential postemergence application of Pursuit to control emerged weeds according to the Pursuit label. INTER Plus II will improve the consistency and level of control from Pursuit on most grass species. Refer to the Pursuit postemergence label for a listing of weeds controlled, application rate, and growth stage limitations.

#### 12. Tank Mixture with Lexone, Sencor, Scepter, Lorox, Lorox Plus, Gemini, Canopy, Preview, or Pursuit, plus Gramoxone Extra or Roundup for Minimum-Tillage or No-Tillage Systems

In minimum-tillage or no-tillage systems where soybeans are planted directly into a cover crop, stale seedbed, established sod, or previous crop residues, the contact herbicides Gramoxone Extra or Roundup may be added to a tank mix of either INTER Plus II+ Sencor or Lexone, INTER Plus II+ Scepter, INTER Plus II+ Lorox, INTER Plus II+ Lorox Plus, INTER Plus II+ Gemini, INTER Plus II+ Canopy, INTER Plus II+ Preview, or INTER Plus II+ Pursuit. When used as directed, the Gramoxone Extra portion of the tank mixture controls most emerged weeds and suppresses many perennial weeds. Roundup combinations will control emerged annual and perennial weeds when applied as directed on the Roundup label. The INTER Plus II + Sencor / Lexone, Scepter, Lorox, Lorox Plus, Gemini, Canopy, Preview, or Pursuit portion of the tank mixture provides preemergence control of the weeds listed on this label in the tank mixture section for INTER Plus II+ Sencor / Lexone, INTER Plus II+ Scepter, INTER Plus II+ Lorox, INTER Plus II+ Lorox Plus, INTER Plus II+ Gemini, INTER Plus II+ Canopy, INTER Plus II+ Preview, and INTER Plus II+ Pursuit, respectively. Refer to the label of each product used in combination and observe the planting details, soybean variety restrictions, information regarding application to soybeans, geographical restrictions, and all other precautions and limitations.

Refer below for rates of Gramoxone Extra or Roundup, INTER Plus II+ Sencor / Lexone, INTER Plus II+ Scepter, INTER Plus II+ Lorox, INTER Plus II+ Lorox Plus, INTER Plus II+ Gemini, INTER Plus II+ Canopy, INTER Plus II+ Preview, and INTER Plus II+ Pursuit, respectively.

**Application:** Apply before, during, or after planting, but before the soybeans emerge, at the rates specified below. Add Gramoxone Extra or Roundup at the following broadcast rates:

**Gramoxone Extra:** 1.5 - 2, 2 - 2.5, or 2.5 - 3 pts./A to 1 - 3 inches, 3 - 6 inches, or 6-inch tall weeds, respectively. Apply surfactant at 1.0 or 2 pts. / 100 gals. of spray mixture with 75% or greater or 50 - 74% nonionic active ingredient, respectively. This treatment will not control weeds taller than 6 inches.

**Note:** Do not apply combinations containing Gramoxone Extra in suspension type liquid fertilizers, as the activity of paraquat will be reduced.

**Roundup:** See the Roundup or Roundup RT label for weeds controlled, recommended rates, and other use directions.

Apply in 20 - 60 gals. of water or fluid fertilizer per acre with ground equipment.

#### INTER Plus II+ Lexone / Sencor + Gramoxone Extra or Roundup

On *loamy sand* with over 2% organic matter, apply 1.0 pts. / A of INTER Plus II+ 0.33 - 0.5 lb./A of Sencor or Lexone DF. On *medium soils*, apply 1.33 pts./A of INTER Plus II + 0.5 - 0.67 lb./A of Sencor or Lexone DF. On *fine soils*, apply 1.33 - 1.67 pts./A of INTER Plus II + 0.67 lb./A of Sencor or Lexone DF.

\* When using Sencor 4 or Lexone 4L, multiply lbs. of DF by 1.5 to get pts./A.

**Precautions:** To avoid crop injury, (1) Do not use this tank mixture on soil with less than 0.5% organic matter, on alkaline soil with a pH over 7.4, or on all sand and on loamy sand with less than 2% organic matter. (2) If heavy rain occurs soon after application, crop injury may result, especially in poorly drained areas where water stands for several days, or where the seeding slit has not been properly closed.

#### INTER Plus II+ Scepter + Gramoxone Extra or Roundup

On *coarse soils*, apply 1.0 pts./A of INTER Plus II + 0.67 pt./A of Scepter. On *medium soils*, apply 1.33 pts./A of INTER Plus II+ 0.67 pt./A of Scepter. On *fine soils*, apply 1.67 pts. /A of INTER Plus II+ 0.67 pt./A of Scepter.

**Notes:** (1) Do not apply within 90 days of harvest, and (2) Do not graze or feed treated soybean forage, hay, or straw to livestock, or illegal residues may result.

#### **INTER Plus II+ Lorox + Gramoxone Extra or Roundup**

On *coarse soils*\*, apply 1.0 pts./A of INTER Plus II+ 1 - 1.5 lbs./A of Lorox DF\*\*. On *medium soils*, apply 1.33 pts./A of INTER Plus II+ 1 - 2 lbs./A of Lorox DF. On *fine soils*, apply 1.33 - 1.67 pts./A of INTER Plus II+ 2 - 3 lbs./A of Lorox DF.

\* Do not use on loamy sand, except in the northeastern U.S. on loamy sand with over 1% organic matter, or injury may occur. Do not use on sand, gravelly soils, or exposed subsoils, or injury may occur.

\*\* When using Lorox L or Lorox DF, use equivalent rates. One pt. of Lorox L equals 1.0 lb. of Lorox DF.

*Precaution: Do not use on soil with less than 0.5% organic matter, or crop injury may occur.*

#### **INTER Plus II+ Lorox Plus + Gramoxone Extra or Roundup**

Use only where soils have 0.5 - 3% organic matter. On *coarse soils*, apply 1.0 pts./A of INTER Plus II+ 12 - 14 oz./A of Lorox Plus 60DF. On *medium soils*, apply 1.33 pts./A of INTER Plus II+ 14 - 16 oz./A of Lorox Plus. On *fine soils*, apply 1.33 - 1.67 pts./A of INTER Plus II+ 16 - 18 oz./A of Lorox Plus.

*Precaution: Do not apply to sand or to any soil with pH greater than 6.8.*

#### **INTER Plus II+ Gemini + Gramoxone Extra or Roundup**

Use only where soils have 0.5 - 3% organic matter. On *coarse soils* (sandy loam only), apply 1.0 pts./A of INTER Plus II+ 12 - 16 oz./A of Gemini 60DF. On *medium soils*, apply 1.33 pts./A of INTER Plus II+ 16 - 20 oz./A of Gemini. On *fine soils*, apply 1.33 - 1.67 pts./A of INTER Plus II+ 20 - 24 oz./A of Gemini.

*Precaution: Do not apply to sand or loamy sand, or to any soil with less than 0.5% organic matter, or to any soil with pH greater than 7.0, except as noted on the Gemini label.*

#### **INTER Plus II+ Canopy + Gramoxone Extra or Roundup**

Use only where soils have 0.5 - 5% organic matter. On *coarse soils* (except sand), apply 1.0 pts./A of INTER Plus II, on *medium soils*, apply 1.33 pts./A of INTER Plus II, and on *fine soils*, apply 1.33 - 1.67 pts./A of INTER Plus II. Refer to the Canopy label for appropriate rate according to geographical location, soil and organic matter classification, pH limitations, and all other use directions.

*Precaution: Do not apply to sand, or to any soil with less than 0.5% organic matter, or to any soil with pH greater than 7.0, except as noted on the Canopy label.*

#### **INTER Plus II+ Preview + Gramoxone Extra or Roundup**

Use only where soils have 0.5 - 5% organic matter. On *coarse soils* (except sand), apply 1.0 pts./A of INTER Plus II+ 6\* - 7 oz./A of Preview 75DF. On *medium soils*, apply 1.33 pts./A of INTER Plus II+ 7\* - 8 oz./A of Preview. On *fine soils*, apply 1.33 - 1.67 pts./A of INTER Plus II+ 8\* to 9 - 10 oz./A of Preview.

\*Use this rate where the soil organic matter is in the 0.5 to less than 3.0% range.

*Precaution: Do not apply to sand, or to any soil with less than 0.5% organic matter, or to any soil with pH greater than 6.8.*

#### **INTER Plus II+ Pursuit + Gramoxone Extra or Roundup**

On *coarse soils*, apply 1.0 pts./A of INTER Plus II+ 0.25 pt./A of Pursuit. On *medium soils*, apply 1.33 pts./A of INTER Plus II+ 0.25 pt./A of Pursuit. On *fine soils*, apply 1.67 pts./A of INTER Plus II+ 0.25 pt./A



of Pursuit.

### 13. Tank Mixture with Broadstrike + Inter 8E II

For preplant surface, preplant incorporated, or preemergence application where severe grass populations are expected on *medium-* or *fine-textured soils* with relatively high organic matter content, Broadstrike + INTER 8E II may be spiked with INTER Plus II for optimum performance. Refer to the Broadstrike + INTER 8E II label for its use rate and the amount of metolachlor active ingredient it contains. INTER Plus II may be added up to, but not to exceed, the maximum alone INTER Plus II label rate for the soil classification. Refer to the respective labels and follow all directions, limitations, and restrictions for each product.

## APPENDICES

### Appendix A: Compatibility Test

Since liquid fertilizers can vary, even within the same analysis, always **check compatibility with herbicide(s) each time before use**. Be especially careful when using **complete** suspension or fluid fertilizers, as serious compatibility problems are more likely to occur. Commercial application equipment may improve compatibility in some instances. The following test assumes a spray volume of 25 gals./A. For other spray volumes, make appropriate changes in the ingredients. Check compatibility using this procedure:

1. Add 1 pt. of fertilizer to each of 2 one-qt. jars with tight lids.
2. To **one** of the jars, add  $\frac{1}{4}$  tsp. or 1.2 milliliters of a compatibility agent approved for this use, such as Compex or Unite ( $\frac{1}{4}$  tsp. is equivalent to 2 pts./100 gals. spray). Shake or stir gently to mix.
3. To **both** jars, add the appropriate amount of herbicide(s). If more than one herbicide is used, add them separately with dry herbicides first, flowables next, and emulsifiable concentrates last. After each addition, shake or stir gently to thoroughly mix. The appropriate amount of herbicides for this test follows:

**Dry herbicides:** For each pound to be applied per acre, add 1.5 level teaspoons to each jar.

**Liquid herbicides:** For each pint to be applied per acre, add 0.5 teaspoon or 2.5 milliliters to each jar.

**Note:** For INTER Plus II tank mixtures with AAtrex + Princep, use  $\frac{1}{3}$  -  $\frac{1}{2}$  the amount of AAtrex specified above and the remainder as Princep, depending on whether the 1:2 or 1:1 ratio of AAtrex to Princep is to be applied.

4. After adding all ingredients, put lids on and tighten, and invert each jar 10 times to mix. Let the mixtures stand 15 minutes and then look for separation, large flakes, precipitates, gels, heavy oily film on the jar, or other signs of incompatibility. Determine if the compatibility agent is needed in the spray mixture by comparing the 2 jars. If either mixture separates, but can be remixed readily, the mixture can be sprayed as long as good agitation is used. If the mixtures are incompatible, test the following methods of improving compatibility: (A) slurry the dry herbicide(s) in water before addition, or (B) add  $\frac{1}{2}$  of the compatibility agent to the fertilizer and the other  $\frac{1}{2}$  to the emulsifiable concentrate or flowable herbicide before addition to the mixture. If incompatibility is still observed, do not use the mixture.

### Appendix B: Low Carrier Application For Broadcast Ground Application Only

Use sprayers, such as Ag-Chem RoGator<sup>®</sup>, Hagie, John Deere Hi-Cycle<sup>T</sup>, Melroe Spra-Coupe, Tyler Patriot<sup>TM</sup>, or Willmar Air Ride<sup>®</sup>, that provide accurate and uniform application. **Only water may be used as a carrier**. Screens in suction and in-line strainers should be 50-mesh. Manufacturers may require that tip screens as fine as 100-mesh be used with some nozzles. Use a pump with capacity to: (1) maintain up to 35-40 psi at the nozzles, and (2) provide sufficient agitation in tank to keep mixture in suspension. Use a minimum of 5 gals. of spray mixture per acre. Maximum recommended sprayer speed is 15 mph. Rinse sprayer thoroughly with clean water immediately after each use.

**Note:** Low pressure nozzles are recommended to reduce drift and increase application accuracy. Care should be taken when using automatic rate controlling devices to spray the material within the rated working pressure and flow ranges of the nozzles selected. Nozzle screens should be used when recommended by the manufacturer. All nozzles should be placed on 20-inch centers, except flooding

types which should be placed on 40-inch centers. When Flat Fan-type nozzles are used, angles of 80° or 110° are recommended. Always read and follow the manufacturer's directions for optimum setup and performance of their nozzles or tips.

### **Appendix C: Aerial Application**

Apply INTER Plus II in water alone or in tank mixtures with AAtrex, Lexone, Lorox, or Sencor in a minimum total volume of 2 gals./A by aircraft. INTER Plus II may also be applied by air in combination with Balan, Prowl, or Treflan. Avoid application under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur. In order to assure that spray will be controllable within the target area when used according to label directions, make applications at a maximum height of 10 ft., using low-drift nozzles at a maximum pressure of 40 psi, and restrict application to periods when wind speed does not exceed 10 mph. To assure that spray will not adversely affect adjacent sensitive nontarget plants, apply INTER Plus II alone or INTER Plus II+ AAtrex by aircraft at a minimum upwind distance of 400 ft. from sensitive plants, or apply INTER Plus II+ Lexone, Lorox, or Sencor at a minimum upwind distance of 300 ft. from sensitive plants.

Avoid application to humans or animals. Flagmen and loaders should avoid inhalation of spray mist and prolonged contact with skin.

### **Aerial Drift Management:**

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment- and weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target drift movements from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
2. Nozzles must always point backwards parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the **Aerial Drift Reduction Advisory Information** section below.

### **Aerial Drift Reduction Advisory Management:**

#### **Information on Droplet Size**

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see **Wind, Temperature and Humidity, and Temperature Inversions**).

#### **Controlling Droplet Size**

- Volume - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- Pressure - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- Number of Nozzles - Use the minimum number of nozzles that provide uniform coverage.
- Nozzle Orientation - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant

- deflection from horizontal will reduce droplet size and increase drift potential.
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

### **Boom Length**

For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

### **Application Height**

Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

### **Swath Adjustment**

When applications are made with a crosswind, the swath will be displaced downward. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.).

### **Wind**

Drift potential is lowest between speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind directions and high inversion potential.

**Note:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

### **Temperature and Humidity**

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

### **Temperature Inversions**

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

### **Sensitive Areas**

This pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive area).

### **Appendix D: Center Pivot Irrigation Application**

INTER Plus II alone or in tank mixture with other herbicides on this label, which are registered for center pivot application, may be applied in irrigation water preemergence (after planting, but before weeds or crop emerge) at rates recommended on this label. Apply this product only through a center pivot irrigation system. Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension specialists, equipment

manufacturers, or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system, unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

#### **Operating Instructions**

1. The system must contain a functional check-valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check-valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump or piston pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.
8. Prepare a mixture with a minimum of 1 part of water to 1 part herbicide(s) and inject this mixture into the center pivot system. Injecting a larger volume of a more dilute mixture per hour will usually provide more accurate calibration of metering equipment. Maintain sufficient agitation to keep the herbicide in suspension.
9. Meter into irrigation water during entire period of water application.
10. Apply in  $\frac{1}{2}$  - 1 inch of water. Use the lower water volume ( $\frac{1}{2}$  inch) on *coarse-textured soils* and the higher volume (1 inch) on *fine-textured soils*. More than 1 inch of water at application may reduce weed control by moving the herbicide below the effective zone in the soil.

*Precaution for center pivot applications: Where sprinkler distribution patterns do not overlap sufficiently, unacceptable weed control may result. Where sprinkler distribution patterns overlap excessively, crop injury may result.*

#### **Appendix E: Dry Bulk Granular Fertilizers**

Many dry bulk granular fertilizers may be impregnated or coated with INTER Plus II alone or selected INTER Plus II tank mixtures which are registered for preplant incorporated or preplant surface application which are used to control weeds in crops on the INTER Plus II label and are not prohibited from use on dry bulk granular fertilizers.

When applying INTER Plus II or INTER Plus II mixtures with dry bulk granular fertilizers, follow all directions for use and precautions on the respective product labels regarding target crops, rates per acre, soil texture, application methods (including timing of application), and rotational crops.

All individual state regulations relating to dry bulk granular fertilizer blending, registration, labeling, and application are the responsibility of the individual and/or company selling the herbicide/fertilizer mixture. Prepare the herbicide/fertilizer mixtures by using any closed drum, belt, ribbon, or other commonly used dry bulk fertilizer blender. Nozzles used to spray INTER Plus II and INTER Plus II mixtures onto the fertilizer must be placed to provide uniform spray coverage. Care should be taken to aim the spray directly onto the fertilizer only and to avoid spraying the walls of the blender.

If the herbicide/fertilizer mixture is too wet, add a highly absorptive material, such as Agsorb<sup>®</sup> or Celatom MP-79<sup>®</sup>, or similar granular clay or diatomaceous earth materials, to obtain a dry, free-flowing mixture.

Absorptive materials should be added only after the herbicide has been thoroughly blended into the fertilizer mixture. Best application results will be obtained by using a granule of 6/30 particle size or of a size similar to that of the fertilizer material being used. Generally, less than 2% by weight of absorptive material will be needed. Avoid using more than 5% absorptive material by weight.

Calculate amounts of INTER Plus II, AAtrex, AAtrex + Princep, Princep, Sencor, Lexone, or Sonalan by the following formula:

$$\frac{2,000}{\text{lbs. of fertilizer per acre}} \times \text{pts./A of liquid or flowable product} = \text{pts. of liquid or flowable product per ton of fertilizer}$$

$$\frac{2000}{\text{lbs. of fertilizer per acre}} \times \text{lbs./A of dry product} = \text{lbs. of dry product per ton of fertilizer}$$

**Pneumatic (Compressed Air) Application (INTER Plus II Alone):** High humidity, high urea concentrations, low fertilizer use rates, and dusty fertilizer may cause fertilizer mixture to build up or plug the distributor head, air tubes, or nozzle deflector plates. To minimize buildup, premix INTER Plus II with Exxon Aromatic 200 at a rate of 1 - 4 pts./gal. of INTER Plus II. Aromatic 200 is a noncombustible/nonflammable petroleum product. Aromatic 200 may be used in either a fertilizer blender or through direct injection systems. Drying agents should not be used when using Aromatic 200.

**Notes:** (1) Mixtures of INTER Plus II and Aromatic 200 must be used on dry fertilizer only. Poor results or crop injury may result if these mixtures are used in water or liquid fertilizer solutions for spraying applications. (2) When impregnating INTER Plus II in a blender before application, a drier mixture can be attained by substituting a drying agent for Aromatic 200. The use of Agsorb FG or drying agents of 6/30 particle size are recommended. (3) Drying agents are not recommended for use with On-The-Go impregnation equipment.

**Precautions:** To avoid potential for explosion, (1) Do not impregnate INTER Plus II or INTER Plus II mixtures on ammonium nitrate, potassium nitrate, or sodium nitrate, either alone or in blends with other fertilizers. (2) Do not use INTER Plus II or INTER Plus II mixtures on straight limestone, since absorption will not be achieved. Fertilizer blends containing limestone can be impregnated.

#### **Application**

Apply 200-700 lbs. of the herbicide/fertilizer mixture per acre. For best results, apply the mixture uniformly to the soil with properly calibrated equipment immediately after blending. Uniform application of the herbicide/fertilizer mixture is essential to prevent possible crop injury. Non-uniform application may also result in unsatisfactory weed control. In areas where conventional tillage is practiced, a shallow incorporation of the mixture into the soil may improve weed control. On *fine-* or *medium-textured soils* in areas where soil incorporation is not planned, i.e., reduced tillage situations or in some conventional till situations, make applications approximately 30 days before planting to allow moisture to move the herbicide/fertilizer mixture into the soil. On *coarse-textured soils*, make applications approximately 14 days prior to planting.

**Precaution:** To avoid crop injury, do not use the herbicide/fertilizer mixture on crops where bedding occurs.

AAtrex, Beacon, Bicep, Caliber 90, Caparol, Concep, Cotoran, Evik, Exceed, Nine-O, and Princep trademarks of Novartis Crop Protection, Inc.

Accent, Canopy, Gemini, Lexone, Lorox, Lorox Plus, and Preview trademarks of E. I. duPont de Nemours and Company, Inc.

Ag-Chem RoGator trademark of Ag-Chem Equipment Company

Agsorb trademark of Oil-Dri Corporation

Balan, Broadstrike\*, Sonalan, and Treflan trademarks of DowElanco

Banvel, Banvel SGF, Basagran, Clarity, Marksman, and Storm trademarks of BASF AG

Brominal, Buctril, trademarks of Rhône-Poulenc Ag Company

Sutan trademark of Cedar Chemical

Celatom MP-79 trademark of Eagle-Picher Industries, Inc.

Command trademark of FMC Corporation

Compex trademark of KALO Agricultural Chemicals, Inc.

Eptam, Eradicane, Gramoxone, and Vemam trademarks of Zeneca Ag Products

Hi-Cycle<sup>T</sup> trademark of John Deere Company

Landmaster, Roundup, and Screen trademarks of Monsanto Company

Prowl, Pursuit, and Scepter trademarks of American Cyanamid

Sencor trademark of Bayer AG

Tyler Patriot<sup>TM</sup> trademark of Tyler Ltd. Partnership

Unite trademark of HACO, Inc.

Willmar Air Ride trademark of Willmar Manufacturing

X-77 trademark of Loveland Industries, Inc.

#### **CONDITIONS OF SALE AND WARRANTY**

**Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.**

Conditions: The directions for use of this product are believed to be adequate and should be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Cedar Chemical Corporation. All such risks shall be assumed by the user or buyer.

Disclaimer of Warranties: THERE ARE NO WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, WHICH EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agents of Cedar Chemical Corporation is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein.

Cedar Chemical Corporation disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT CEDAR CHEMICAL CORPORATION'S ELECTION, THE REPLACEMENT OF PRODUCT.

Cedar Chemical Corporation  
5100 Poplar Avenue  
Suite 2414  
Memphis, Tennessee 38137

Rev. 3/5/02