

Pm. 22

56212-2

Page 19

DMX-7 PLUS

LECITHIN

FOR THE INHIBITION OF MOLD IN GRAINS

Active Ingredient: Propionic Acid.....	39.2%
Inert Ingredients:.....	60.8%
TOTAL	100 %

PRODUCT WEIGHT: 8.9 Pounds Per Gallon

KEEP OUT OF REACH OF CHILDREN

CAUTION: **PRECAUTIONARY STATEMENTS** **HAZARDOUS TO HUMANS & DOMESTIC ANIMALS**

Harmful if inhaled or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Statement of Practical Treatment


- If in Eyes:* Flush eyes with plenty of water. Call a physician if irritation persists.
- If Swallowed:* Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.
- If on Skin:* Wash with plenty of soap and water. Get medical attention.
- If Inhaled:* Remove victim to fresh air. If not breathing, give artificial respiration preferably mouth to mouth. Get medical attention.

See side panel for additional precautions.

NET CONTENTS: 500 POUNDS

EPA REGISTRATION NO. 56212-2

Delst, Inc.
540 E. Jamie Ave.
La Habra, California 91731



BEST COPY AVAILABLE

199

Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark.

Do not contaminate water by disposal of equipment washwaters.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

May be used on any grain intended for use as food or feed. Do not use on grain intended for use as seed or for malting purposes.

STORAGE AND DISPOSAL

Pesticide Disposal: Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent) then offer for recycling or reconditioning, or puncture and dispose of in a sanitary land fill. Or if plastic, incinerate, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Pesticide Storage: Store in a cool dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food and feed. Store in original container and out of reach of children, preferably in a locked storage area. Handle and open for any reason or cause, carefully sweep material into pile. Refer to precautionary statement on label for hazards associated with the handling of this material. Do not walk through spilled material. Dispose of pesticides as directed above. In spill incidents, keep unauthorized people away.

Do not store treated grain in the same area with untreated grain or forage.

Determine moisture content of freshly harvested grain and treat within 4 hours of harvest.

Apply at rate specified below depending on moisture content and type or length of storage.

Do not treat in the rain. Allow rain to soak into the pile and test for moisture content before continuing treatment.

FOR THE INHIBITION OF MOLD IN GRAINS:

Thoroughly mix DMX-7 PLUS and grain by spraying DMX-7 PLUS on grain in auger during movement into storage. If spray failure occurs, immediately stop auger to assure all grain will be treated. Establish auger throughput and set flowmeter to apply proper rate of DMX-7 PLUS.

APPLY THIS PRODUCT ONLY AT THE RECOMMENDED RATES.

Flush out application and handling equipment with water after DMX-7 application is completed.

WEATHERPROOF STORAGE OF GRAINS:

Follow general directions as stated above. Protect grain from outside moisture in metal or wooden bins, converted cribs, Quonset huts, or other roofed structures. Waterproof floor.

USE THE TREATMENT RATES AS GIVEN BELOW:

Recommended rates of DMX-7 PLUS per ton of corn, sorghum, wheat, oats, or barley in covered storage.

Moisture Content of Grain, Percent by Weight		
14%	16%	18%
4 lbs. or 58 liq. oz.	5 lbs. or 72 liq. oz.	6 lbs. or 86 liq. oz.

1 gallon - 128 liquid ounces

Will inhibit mold growth for approximately 2 to 4 weeks, depending upon humidity, air temperature and air temperature fluctuations within the specified storage facilities. The shorter period of protection will apply during periods of high humidity, high temperature and/or wide temperature fluctuation.

WARRANTY:

- The manufacturer guarantees and warrants (a) that the active ingredient content and the total net contents are as stated within lawful limits, and (b) that the directions are based upon responsible experts evaluation of the individual ingredients of the material herein.
- The manufacturer neither makes nor authorizes any agent or representative to make any other warranties, including warranties of fitness for a particular purpose or merchantability, guarantee or representation, expressed or implied, concerning this material. Timing and method of application, weather and crop condition, and other influencing factors are beyond control of the manufacturer, so this product is sold only on the basis that buyer assumes all risks of use or handling which result in loss or damage and which are beyond manufacturer's control. No claim of any kind, whether not based on negligence, shall be in greater amount than the purchase price of the material in respect to which such claim is made. In no event shall manufacturer or seller be liable for special, indirect or consequential damages resulting from the use or handling of this material.
- No modification of this warranty and disclaimer is authorized, except by specific reference to them in writing by an employee of the manufacturer.

BEST COPY AVAILABLE