

[Front Panel]

RESTRICTED USE PESTICIDE

Due to Inhalation Hazards to Humans and Hazards to Nontarget Species

For retail sale to, and use only by Certified Applicators, or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certification.

**Zinc Phosphide on Wheat
For Mouse Control**

For use in orchards, groves, nurseries, vineyards, ornamentals, and nonbearing fruit trees; reforestation seeding areas; and alfalfa, barley, wheat, dry beans, potatoes, and sugar beets to control voles and dry life-footed mice.

Active Ingredient:
Zinc Phosphide 1.82%
Inert Ingredients: 98.18%
TOTAL: 100.00%

ACCEPTED
JUL 10 2006
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 56228-3

KEEP OUT OF REACH OF CHILDREN [See minimum type size requirements.]

WARNING

FIRST AID

Have label with you when obtaining treatment advice.

Any person who applies zinc phosphide products and experiences signs or symptoms such as nausea, abdominal pain, tightness in the chest, or weakness, should be seen by a physician immediately.

If swallowed	<ul style="list-style-type: none"> •Immediately call a poison control center, doctor, or 1-800-858-7378 or transport the person to the nearest hospital. •DO NOT DRINK WATER. •Do not give anything by mouth or make person vomit unless told to do so by the poison control center or doctor.
If on skin or clothing	<ul style="list-style-type: none"> •Take off contaminated clothing. •Rinse skin immediately with plenty of water for 15-20 minutes. •Call a poison control center, doctor, or 1-800-858-7378 immediately for treatment advice.
If inhaled	<ul style="list-style-type: none"> •Move person to fresh air. •If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. •Call a poison control center, doctor, or 1-800-858-7378 immediately for treatment advice.
If in eyes	<ul style="list-style-type: none"> •Hold eye open and rinse slowly and gently with water for 15-20 minutes. •Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. •Call a poison control center, doctor, or 1-800-858-7378 immediately for treatment advice.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage. See additional PRECAUTIONARY STATEMENTS on Left Panel.

[Left Panel]

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

May be fatal if inhaled. Harmful if absorbed through the skin or swallowed. Causes eye irritation. Do not get in eyes, on skin or on clothing. Do not breathe dust. Keep away from humans, domestic animals and pets. Avoid contamination of feed or foodstuffs. Avoid contact with acids. Wash all utensils, spoons, or measuring devices thoroughly after using.

Personal Protective Equipment (PPE)

All applicators/mixers/loaders must wear:

- long sleeve shirt and long pants;
- shoes plus socks;
- chemical-resistant gloves (any waterproof material, Category A; and
- a dust/mist filtering respirator (MSHA/NIOSH) approval number prefix TC-21C or a NIOSH-approved respirator with an R, P, or HE filter.

All applicators and any person who and retrieves carcasses and unused bait must wear:

- gloves.

User Safety Requirements:

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions are provided for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Wash hands thoroughly after applying bait and before eating, drinking, chewing gum, using tobacco or using the toilet.

User Safety Recommendations:

Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

[Right Panel]

ENVIRONMENTAL HAZARDS

This product is toxic to wildlife and fish. Birds and other wildlife feeding on toxic bait may be killed. Use with care when applying in areas frequented by wildlife or adjacent to any body of water. Do not apply where runoff is likely to occur. Do not apply this product directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply this product in areas inhabited by livestock or where a hazard exists to rare or endangered species. Dogs and other predatory and scavenging mammals might be poisoned if they feed upon animals that have eaten this bait.

ENDANGERED SPECIES

This product may have effects on Federally listed threatened or endangered species or their critical habitat in some counties. It is a violation of Federal law to kill, harm or harass listed animal species without authorization. The killing of a member of an endangered species may result in a fine and/or imprisonment under the Endangered Species act. To limit the potential for such impacts when using this product, consult and follow the "Endangered Species Considerations" provided on this label and in the EPA Endangered Species Bulletin for the County or Parish in which you are applying this product. To determine whether you County or Parish has a Bulletin, consult <http://www.epa.gov/espp> before each season's use of this product.

United States Department of Agriculture (USDA)
Animal and Plant Health Inspection Service
Riverdale, MD20737-1237

EPA Reg. No. 56228-3
EPA Est. No. 56228-ID-1

Net Contents

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its label.

READ THIS LABEL: Read entire label and follow all use directions, use precautions, and use restrictions which pertain to the bait application to be made.

GENERAL USE RESTRICTIONS (For All Uses):

Do not apply this product by any method or use not specified on this label. Do not apply bait on roads, over water, or, except where specifically indicated in these directions, where plants are grown for food or feed. Wear all required personal protective equipment (PPE) when mixing, handling, and applying bait. See "PRECAUTIONARY STATEMENTS".

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only persons associated with the applications may be present in the area when the bait is being applied and such persons must wear required PPE. Keep all other persons out of the treated area during application.

ORCHARDS, GROVES, NURSERIES, VINEYARDS, ORNAMENTAL PLANTINGS, AND NON-BEARING FRUIT TREES (to control Voles and White-footed Mice)

USE RESTRICTIONS: Use this product to control meadow voles (*Microtus pennsylvanicus*), prairie voles (*M. ochrogaster*), pine voles (*M. pinetorum*), and white-footed mice (*Peromyscus leucopus*) in orchards, groves, horticultural nurseries, vineyards, ornamental plantings, and plantings of nonbearing fruit trees. Do not apply bait to bare ground.

Special Use Restrictions for Orchards and Groves:

Applications (except aerial) must be made at a rate 6 to 10 pounds of 2% zinc phosphide bait per acre. Aerial applications must be made at 4 to 9.5 pounds of 2% zinc phosphide bait per acre. Applications can be made after harvest or anytime during the dormant season, but before tree growth begins in the spring. Do not broadcast over non-orchard, nongrove crops. Do not graze animals on treated areas.

Special Use Restrictions for Vineyards:

Baits may be applied only during the non-bearing season. Do not apply bait over the growing crop. Do not apply bait by air. Do not graze animals on treated areas. **Ground broadcast applications** must not exceed 10 pounds of bait per acre.

APPLICATION DIRECTIONS:

Hand Baiting: Near the base of each infested tree, place one teaspoonful quantities of bait at 2 to 4 locations, either on surface or at the mouth of each of the holes leading to underground burrow systems. Cover bait artificially (e.g., mats, boards) or by pulling overhanging grass back into place. Do not disturb the runway system. Apply 3 to 5 pounds of 2% zinc phosphide bait per acre of infested trees.

TRAILBUILDER BAITING: Drop teaspoonful quantities of bait at 4 to 5 foot intervals in the artificial trail made by the machine just inside the drop line on both sides of the trees. Apply 2 to 3 pounds of 2% zinc phosphide bait per acre.

GROUND BROADCAST BAITING: Under infested trees broadcast evenly by cyclone seeder or by hand. Concentrate in areas with the heaviest vegetative cover. Apply at a rate of 6 to 10 pounds of 2% zinc phosphide per acre.

AERIAL APPLICATION: Broadcast bait at a rate of 6 to 10 pounds of 2% zinc phosphide per acre. Bait applications should be made immediately after harvest and before leaf fall, when the grass is not yet matted. Do not make broadcast application of bait over growing crops when the bait may lodge on the plant. Do not apply to bare ground.

RANGELAND AND REFORESTATION SEEDING AREAS (to control voles and white-footed mice)

USE RESTRICTIONS: Use this product to control meadow voles (*Microtus pennsylvanicus*), prairie voles (*M. ochrogaster*), pine voles (*M. pinetorum*), and white-footed mice (*Peromyscus leucopus*) in rangeland and reforestation area. Apply only to areas with less than 50% ground cover. For **hand bait:** Apply at a rate of 1 teaspoon per mound or burrow at a maximum of one application per year. Do not graze animals in treated areas. Do not apply where plants are grown for food or feed or in areas inhabited by livestock. For **broadcast treatment:** Apply at the rate of 6 to 10 pounds of 2% zinc phosphide per acre in the fall before the seeds fall or seeding efforts. Treatment may be necessary in spring to protect emergent seedlings.

APPLICATION DIRECTIONS

BAITING: For hand baiting, place 1 teaspoon of bait in surface runways or in burrow entrances. If a trailbuilder is used, apply at the rate of 1 teaspoon of bait at 4 to 5 foot intervals. For broadcast baiting, apply in 20 foot swaths evenly by hand or with a cyclone seeder at a rate of 5.5 to 9 pounds of 2% zinc phosphide bait per acre. For aerial application, broadcast bait at the rate of 6 to 10 pounds of 2% zinc phosphide per acre per treated swath acre.

ALFALFA, BARLEY, DRY BEANS, POTATOES, SUGAR BEETS, AND WHEAT (to control Voles and White-Footed Mice)

USE RESTRICTIONS: Use this product to control meadow voles (*Microtus pennsylvanicus*), prairie voles (*M. ochrogaster*), pine voles (*M. pinetorum*), and white-footed mice (*Peromyscus leucopus*) on alfalfa, barley, dry beans, potatoes, sugar beets, and wheat.

PREBAITING: Prebaiting with untreated wheat may lead to more consistent acceptance of the toxic bait. Prebaits should be applied, at rates identical to toxic baiting, beginning 2 to 3 days before toxic baits are to be used.

APPLICATION DIRECTIONS:

ALFALFA:

Use Restrictions: All applications must occur shortly after a cutting of the hay, and/or prior to new growth attaining a length of 2 inches. Alfalfa forage from treated areas must not be harvested until it reaches maturity. Do not apply this product to alfalfa within 30 days of harvest. This use is restricted to California, Idaho, Oregon, and state of Washington.

BROADCAST BAITING: This product may be broadcast by air or ground driven dispensing devices. Apply Zinc Phosphide bait at a rate up to 10 lbs. (0.2 lb active) per acre. A second application may be after 10 days. A maximum of 20 lbs. (0.4 lb active) per acre may be applied per year. Do not apply by air when wind velocity exceeds 10 mph. Do not apply in piles or permit piles to be formed by equipment.

BARLEY

USE RESTRICTIONS: Do not apply bait post-harvest or pre-plant. Do not apply this product to barley within 50 days of harvest. This use is restricted to Idaho, Oregon and the state of Washington.

BROADCAST BAITING: This product may be broadcast by air or ground driven dispensing devices. Apply Zinc Phosphide bait at a rate up to 6 lbs. (0.12 lb active) per acre. Two applications may be made at a minimum interval of 25 days. A maximum of 12 lbs (0.24 lb active) per acre may be applied per year. All applications must be made prior to the boot stage.

POTATOES

USE RESTRICTIONS: Do not apply bait post-harvest or pre-plant. Do not apply this product to potatoes within 30 days of harvest. This use is restricted to Idaho, Oregon and the state of Washington.

BROADCAST BAITING: This product may be broadcast by air or ground driven dispensing devices. Apply Zinc Phosphide bait at a rate up to 10 lbs. (0.2 lb active) per acre, or two applications of Zinc Phosphide Pellets at the rate of up to 5 lbs. (0.1 lb active) per acre. The maximum application rate per year is 10 lbs. (0.2 lb active).

SUGAR BEETS

USE RESTRICTIONS: Do not apply bait post-harvest or pre-plant. Do not apply this product to sugar beets within 30 days of harvest. This use is restricted to Idaho, Oregon and the state of Washington.

BROADCAST BAITING: This product may be broadcast by air or ground driven dispensing devices. Apply Zinc Phosphide bait at a rate up to 10 lbs. (0.2 lb active) per acre at planting. A second application may be made at this rate. The maximum application rate per year is 20 lbs. (0.4 lb active).

WHEAT

USE RESTRICTIONS: Do not apply bait post-harvest or pre-plant. Do not apply this product to wheat within 50 days of harvest. This use is restricted to Idaho, Oregon and the state of Washington.

BROADCAST BAITING: This product may be broadcast by air or ground driven dispensing devices. Apply Zinc Phosphide bait at a rate up to 6 lbs. (0.12 lb active) per acre. A second applications may be made within 25 days. The maximum application rate per year is 12 lbs. (0.24 lb active).

DRY BEANS

USE RESTRICTIONS: Do not apply by air. Do not apply by air. Do not apply this product to dry beans within 30 days of harvest. This use is restricted to Idaho, Oregon and the state of Washington.

BROADCAST BAITING: This product may be broadcast by hand or ground driven dispensing devices at a rate of 6 lbs. (0.12 lb active) per acre. No more than one application per growing season is permitted.

NON-CROP AREAS (RIGHTS-OF-WAYS, PARKS, HIGHWAY MEDIANS, NONRESIDENTIAL LAWNS, TURF, GOLF COURSES, AIRPORT GRASSES) to control Voles and White-Footed Mice

USE RESTRICTIONS: Use this product to control meadow voles (*Microtus pennsylvanicus*), prairie voles (*M. ochrogaster*), pine voles (*M. pinetorum*), and white-footed mice (*Peromyscus leucopus*) in non-crop areas (rights-of-ways, nonresidential lawns, parks, turf, golf courses, airport grasses, highway medians). Do not apply bait to bare ground. Do not apply bait by air.

PREBAITING: Prebaiting with untreated wheat may lead to more consistent acceptance of toxic bait. Prebaits should be applied, at rates appropriate for toxic baiting, beginning 2 or 3 days before toxic baits are to be used.

INS.

APPLICATION DIRECTIONS:

Hand Baiting: Place one (4 grams) of bait in surface runways or at the mouths of holes leading to underground burrow systems. Apply bait at rats of 3 to 5 pounds (0.06 to 0.1 pounds of active ingredient) per acre.

Trail-Builder Baiting (grain or sunflower seed baits only): Follow manufacturer's instructions for type of equipment used. Set equipment to drop one teaspoon (4 grams) at 4- to 5-foot intervals in the artificial trail made by the machine. Apply 2 to 3 pounds of 2% zinc phosphide (0.04 to 0.06 pounds of active ingredient) per acre.

Ground Broadcast Baiting (grain or sunflower seed baits only):

Broadcast bait evenly by cyclone seeder or by gloved hand. Apply 6 to 10 pounds of 2% zinc phosphide bait (0.12 to 0.2 pounds of active ingredient) per acre.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store only in original container, in a dry place inaccessible to children, pets, and domestic animals.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spilled bait, or rinsate is a violation of Federal law. If these wastes cannot be disposed of according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

ENDANGERED SPECIES CONSIDERATIONS

Whooping Crane (*Grus americana*): Do not use this product 30 days prior to arrival and 30 days after the whooping crane leaves its known critical habitat.

Attwater's Greater Prairie Chicken (*Tympanuchus cupido attwateri*): Do not use this product in critical habitat of the Attwater's greater prairie chicken in the following Texas counties: Aransas, Austin, Brazoria, Colorado, De Witt, Ft. Bend, Galveston, Goliad, Harris, Refugio, Victoria, Waller, and Wharton.

Yellow-Shouldered Blackbird (*Agelaius xanthomus*) and Puerto Rican Plain Pigeon (*Columba inornata wetmorei*): Do not use in critical habitat except in tamper-proof bait boxes: within 6.3 miles of central Aquirre Lago Cidra, Ceiba, San German; 9.4 miles of La Esperanza; south of highway 2 from city at Mayaquez to the city of Ponce, and all of Mona Island, Puerto Rico.

Utah Prairie Dog (*Cynomys parvidens*): Do not use this product in critical habitat of the Utah prairie dog. (Utah).

Salt Marsh Harvest Mouse (*Reithrodontomys raviventris*): Do not use this product in critical habitat within 0.5 mile of salt marsh vegetation and/or brackish water wetlands which are located: 1) near or adjacent to San Pablo Bay and San Francisco Bay, or 2) in the Sacramento River below or adjacent to the confluence of the Sacramento River and the San Joaquin River. (California)

Morro Bay Kangaroo Rat (*Dipodomys heermanni morroensis*): Do not use this product in critical habitat within 2.5 miles of Baywood Park which is located on Morro Bay. (California)

Northern Idaho Ground Squirrel (*Spermophilus brunneus brunneus*): Do not use this product within 0.5 mile of the critical habitats of the Northern Idaho Ground Squirrel in Adams and Valley counties, Idaho.