



U S ENVIRONMENTAL PROTECTION AGENCY
 Office of Pesticide Programs
 Registration Division (7505P)
 1200 Pennsylvania Ave N W
 Washington D C 20460

EPA Reg Number
 56228-3

Date of Issuance
 SEP 27 2012

NOTICE OF PESTICIDE
 Registration
 Reregistration
 (under FIFRA as amended)

Terms of Issuance
 Unconditional
 Zinc Phosphide on Wheat

Name and Address of Registrant (include ZIP Code)

U S Department of Agriculture
 Animal and Plant Health Inspection Service
 4700 River road Unit 152
 Riverdale MD 20737

Decision 324965
 676 5-924479
 1275

Attention Ms Stephanie Stephens

Note Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce In any correspondence on this product always refer to the above EPA registration number

On the basis of information furnished by the registrant the above named pesticide is hereby registered/reregistered under the Federal Insecticide Fungicide and Rodenticide Act Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency In order to protect health and the environment the Administrator on his motion may at any time suspend or cancel the registration of a pesticide in accordance with the Act The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others

EPA received your label amendment of September 27 2012 EPA grants this request under the authority of section 3(c)(5) of the Federal Insecticide Fungicide and Rodenticide Act as amended

Signature of Approving Official

John Hebert
 Product Manager (07)
 Insecticide-Rodenticide Branch
 Registration Division (7505P)

Date

SEP 27 2012

Page 2

Notice of Reregistration (continued)
EPA Reg No 56228 3

With this accepted labeling the product meets the risk management goals of the Reregistration Eligibility Decision (RED) for Zinc Phosphide Therefore EPA reregisters the product listed above This action is taken under the authority of section 4(g)(2)(c) of the Federal Insecticide Fungicide and Rodenticide Act as amended Reregistration under this section does not eliminate the need for continual reassessment of pesticides

Confidential Statement of Formula

The acceptable Confidential Statements of Formula (CSF) for this product are dated September 27 2012 (Basic Formulation and Alternative) All other versions are obsolete

Existing Stocks Provisions

Stocks of product with existing labels may be used for twelve (12) months

Per 40 CFR 156 10(a)(6) submit one copy of your final printed labeling before you release the product for shipment

Follow the enclosed Minimum Type Size Requirements for Final Printed Label

If these conditions are not complied with the registration will be subject to cancellation in accordance with FIFRA section 6(e) Your release for shipment of the product constitutes acceptance of these conditions

We have enclosed a copy of your label stamped Accepted for your records

If you have any questions or comments regarding this letter please contact Mr Dan Peacock by phone (703- 305-5407) or e-mail (peacock dan@epa gov)

- Enclosures 1 Stamped Label
- 2 Minimum Type Sizes for Final Printed Label

056228 00003 20120927

09/25/2012 FINAL clean

RESTRICTED USE PESTICIDE DUE TO HAZARDS TO NONTARGET SPECIES

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator s Certification

ZINC PHOSPHIDE ON WHEAT

For use in dormant pome and stone fruit orchards non bearing nursery stock conifer/Christmas tree groves ornamentals and nonbearing fruit trees rangelands and reforestation seeding areas vineyards airports highway medians turf farms nonresidential lawns golf courses parks and alfalfa barley potatoes sugar beets and beans (dry) to control voles and mice

ACCEPTED SEP 27 2012

Active Ingredient
Zinc Phosphide
Inert Ingredients
Total

Under the Federal Insecticide Fungicide,
and Rodenticide Act as amended, for the
pesticide registered under

EPA Reg No 56228-3

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed absorbed through the skin or inhaled Causes moderate eye irritation Do not get in eyes on skin or on clothing Avoid breathing dust Keep away from humans domestic animals and pets Avoid contamination of feed or foodstuffs Avoid contact with acids Wash all utensils spoons or measuring devices thoroughly after use

PERSONAL PROTECTIVE EQUIPMENT (PPE)

All handlers including loaders and applicators must wear long sleeved shirts long pants shoes socks and waterproof gloves

In addition persons loading pellets or baits into aircraft or mechanical ground equipment and persons loading/applying with hand pushed or hand held equipment such as a push type spreader or cyclone spreader must wear a dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC 21C or a NIOSH approved respirator with an R P or HE filter) and protective eyewear

Any person who retrieves carcasses or unused bait following application of this product must wear waterproof gloves

User Safety Requirements

Follow manufacturer s instructions for cleaning/maintaining PPE If no such instructions for washables are available use detergent and hot water Wash the outside of gloves before removing or turn gloves inside out when removing and dispose of properly Keep and wash PPE separately from other laundry Remove PPE immediately after handling this product As soon as possible wash hands thoroughly after applying bait and before eating drinking chewing gum using tobacco or using the toilet and change into clean clothing

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

HAVE LABEL WITH YOU WHEN SEEKING TREATMENT ADVICE (1 800 858 7378)

If you experience signs and symptoms such as nausea abdominal pain tightness in chest or weakness see a physician immediately For information on health concerns medical emergencies or pesticide incidents call 911 or the National Pesticide Information Center at 1 800 858 7378

IF SWALLOWED	Call 911 a Poison Control Center doctor or 1 800 858 7378 immediately for treatment advice or transport the person to the nearest hospital Do not give any liquid to the patient Do not administer anything by mouth Do not induce vomiting unless told to do so by the poison control center or doctor
IF ON SKIN OR CLOTHING	Take off contaminated clothing Rinse skin immediately with plenty of water for 15 20 minutes Call a Poison Control Center doctor or 1 800 858 7378 immediately for treatment advice
IF INHALED	Move person to fresh air If person is not breathing call 911 or an ambulance then give artificial respiration preferably mouth to mouth if possible Call a Poison Control Center doctor or 1 800 858 7378 immediately for treatment advice
IF IN EYES	Hold eye open and rinse slowly and gently with water for 15 20 minutes Remove contact lenses if present after the first 5 minutes then continue rinsing eye Call a Poison Control Center doctor or 1 800 858 7378 immediately for treatment advice

NOTE TO PHYSICIAN

Contains the phosphine producing active ingredient zinc phosphide Probable mucosal damage may contraindicate the use of gastric lavage Sodium bicarbonate can be given orally to neutralize the stomach acidity The stomach and intestinal tract can be evacuated oxygen administered and cardiac and circulatory stimulants given See Left Panel for additional precautionary statements

TREATMENT FOR PET POISONING

If animal eats bait call a veterinarian at once

NOTE TO VETERINARIAN

For animals ingesting bait and/or showing poisoning signs induce vomiting by using hydrogen peroxide

ENVIRONMENTAL HAZARDS

This product is extremely toxic to birds fish and other wildlife Wildlife feeding on treated bait may be killed Dogs cats and other predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark Do not apply where runoff is likely to occur Do not contaminate water by cleaning of equipment or disposal of wastes

ENDANGERED SPECIES

Notice The use of this product may pose a hazard to Federally designated endangered/ threatened species It is a Federal offense to use any pesticide in a manner that results in the death of a member of an endangered species Consult the nearest U S Fish and Wildlife Service regional office or the appropriate state agency for current information on habitats occupied by endangered species

See the last page of this label for ENDANGERED SPECIES CONSIDERATIONS

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its label

READ THIS LABEL Read entire label and follow all use directions use precautions and use restrictions Use only for the sites pests and application methods described on this label

Use Restrictions – Non Structural Agricultural Sites

Do not apply this product in a way that will contact workers or other persons either directly or through drift

Only protected handlers may be in the area during application

Keep all other persons out of the treated area during this application

Limitations in All Agricultural Sites

Do not apply on roads near residential areas or over water

Do not broadcast over crops unless use directions specifically permit aerial application

Do not apply by air when wind velocity exceeds 10 mph

Do not apply this product in areas inhabited by livestock

When possible do not apply when precipitation is expected

PREBAITING (RECOMMENDED BUT NOT REQUIRED FOR ALL USES UNLESS OTHERWISE SPECIFIED) Prebaiting with untreated wheat may improve acceptance of this zinc phosphide bait

Prebait should be applied at rates appropriate for treated bait 2 or 3 days before toxic baits are used

FRUIT AND NUT TREE ORCHARDS (DORMANT)

USE RESTRICTIONS For control of meadow voles (*Microtus pennsylvanicus*) prairie voles (*M ochrogaster*) pine voles (*M pinetorum*) long tailed voles (*M longicaudus*) California voles (*M californicus*) Oregon voles (*M oregoni*) mountain voles (*M montanus*) Townsend s voles (*M townsendii*) white footed mice (*Peromyscus leucopus*) and deer mice (*P maniculatus*) in pome fruit (apple pear) and stone fruit (peach cherry apricot plum prune nectarine)

Apply when no rain or snow is expected for three consecutive days Apply only after fall harvest (including drops) before new spring growth

PREBAITING (Recommended but not required) To increase acceptance of treated baits apply 1 teaspoon (4 grams) of untreated wheat per tree 1 or 2 days prior to scheduled application of this product

BAITING

HAND BAITING

Place 1 teaspoon (4 grams) of bait at 2 4 locations near bases of each infested tree either on surface trails or at mouth of holes leading to underground burrow systems Do not allow bait to be exposed on bare ground Do not disturb the runway system Apply at a rate up to 5 lbs/A (0 09 lb a i /A)

TRAILBUILDER

Set equipment to drop 1 teaspoon of bait (4 grams) at 4 5 foot intervals in the artificial trail made by the machine just inside the drop line on both side of the trees Apply at a rate up to 3 lbs/A (0 06 lb a i /A)

GROUND BROADCAST

Broadcast evenly by cyclone seeder or by hand Concentrate in areas with heaviest vegetative cover Do not broadcast on bare ground Apply at a rate up to 10 lbs/A (0 18 lb a i /A)

AERIAL BROADCAST

Do not broadcast on bare ground Apply at a rate up to 10 lbs/A (0 18 lb a i /A)

NON BEARING NURSERY STOCK, CONIFER/CHRISTMAS TREES, ORNAMENTAL, AND NONBEARING FRUIT TREES, AND TREE PLANTATIONS AND WINDBREAKS

USE RESTRICTIONS For control of meadow voles (*Microtus pennsylvanicus*) prairie voles (*M ochrogaster*) pine voles (*M pinetorum*) long tailed voles (*M longicaudus*) California voles (*M californicus*) Oregon voles (*M oregoni*) mountain voles (*M montanus*) Townsend s voles (*M townsendii*) white footed mice (*Peromyscus leucopus*) and deer mice (*P maniculatus*) in non bearing nursery stock conifer/Christmas trees ornamental non bearing fruit trees and windbreaks

Apply when no rain or snow is expected for three consecutive days

PREBAITING (Recommended but not required) To increase acceptance of treated baits apply 1 teaspoon (4 grams) of untreated wheat per tree 1 or 2 days prior to scheduled application of this product

BAITING

HAND BAITING

Place 1 teaspoon (4 grams) of bait near bases of each infested tree at 2-4 locations either on surface trails or at mouth of holes leading to underground burrow systems Do not allow bait to be exposed on bare ground Do not disturb the runway system Apply at the rate of 1 teaspoon (4 grams) per placement a rate up to 5 lbs/A (0.09 lb a.i./A)

TRAILBUILDER

Set equipment to drop 1 teaspoon of bait (4 grams) at 4-5 foot intervals in the artificial trail made by the machine just inside the drop line on both side of the trees Apply at a rate up to 3 lbs/A (0.06 lb a.i./A)

GROUND BROADCAST

Broadcast evenly by cyclone seeder or by hand Concentrate in areas with heaviest vegetative cover Do not broadcast on bare ground Apply at a rate up to 10 lbs/A (0.18 lb a.i./A)

AERIAL BROADCAST

Do not broadcast on bare ground Apply at a rate up to 10 lbs/A (0.18 lb a.i./A)

GRAPE VINEYARDS

USE RESTRICTIONS For control of meadow voles (*Microtus pennsylvanicus*) prairie voles (*M. ochrogaster*) pine voles (*M. pinetorum*) long tailed voles (*M. longicaudus*) white footed mice (*Peromyscus leucopus*) deer mice (*P. maniculatus*) and meadow jumping mice (*Zapus hudsonius*) in vineyards Do not apply by air

BAITING

HAND BAITING

Place 1 teaspoon (4 grams) of bait near bases of each infested tree at 2-4 locations either on surface trails or at mouth of holes leading to underground burrow systems Do not allow bait to be exposed on bare ground Do not disturb the runway system Apply at the rate of 1 teaspoon (4 grams) per placement a rate up to 5 lbs/A (0.09 lb a.i./A)

TRAILBUILDER

Set equipment to drop one teaspoon quantity of bait (4 grams) at 4-5 foot intervals in the artificial trail made by the machine just inside the drop line on both side of the trees. Apply at a rate up to 3 lbs/A (0.06 lb a.i./A)

GROUND BROADCAST

Broadcast evenly by cyclone seeder or by hand. Concentrate in areas with heaviest vegetative cover. Do not broadcast on bare ground. Apply at a rate up to 10 lbs/A (0.18 lb a.i./A)

RANGELANDS AND REFORESTATION AREAS

USE RESTRICTIONS Use this product to control meadow voles (*Microtus pennsylvanicus*) prairie voles (*M. ochrogaster*) pine voles (*M. pinetorum*) long tailed voles (*M. longicaudus*) white footed mice (*Peromyscus leucopus*) and deer mice (*P. maniculatus*) in rangeland and reforestation seeding areas. Apply only to areas with less than 50% ground cover

BAITING**HAND BAITING**

Apply bait at the rate of 1 teaspoon (4 grams) either on surface trails or at mouth of holes leading to underground burrow systems. Do not allow bait to be exposed on bare ground. Do not disturb the runway system. Apply at a rate up to 3 lbs/A (0.06 lb a.i./A)

GROUND BROADCAST

Broadcast evenly in 20 foot swaths by cyclone seeder or by hand. Concentrate in areas with heaviest vegetative cover. Do not broadcast on bare ground. Apply at a rate up to 10 lbs of bait/A (0.18 lb a.i./A) in the fall before the seeds fall or seeding efforts. Treatment may be necessary in spring to protect emergent seedlings

TRAILBUILDER

Set equipment to drop 1 teaspoon quantity of bait (4 grams) at 4-5 foot intervals in the artificial trail made by the machine just inside the drop line on both side of the trees. Apply at a rate up to 3 lbs/A (0.06 lb a.i./A)

AERIAL BROADCAST

For aerial application broadcast bait at a rate up to 10 lbs of bait per treated acre (0.18 lb a.i./A)

ALFALFA

USE RESTRICTIONS For control of meadow voles (*Microtus pennsylvanicus*) prairie voles (*M. ochrogaster*) pine voles (*M. pinetorum*) long tailed voles (*M. longicaudus*) California voles (*M. californicus*) Oregon voles (*M. oregoni*) mountain voles (*M. montanus*) Townsend's voles (*M. townsendii*) white footed mice (*Peromyscus leucopus*) and deer mice (*P. maniculatus*) in alfalfa. All applications must occur shortly after a cutting of the hay and/or prior to the next growth's attaining a

length of 2 inches Alfalfa forage from treated areas must not be harvested until it reaches maturity
This use is restricted to Montana, California, Idaho, Oregon, Utah and Washington

BAITING

BROADCAST BAITING This product may be broadcast by air or ground driven dispensing devices Apply at a rate up to 10 lbs/A (0 18 lb a i /A) A second application may be made after 10 days Do not apply in piles or permit piles to be formed by equipment Apply bait at a maximum rate of 20 lbs/A (0 36 lb a i /A) per year

BARLEY

USE RESTRICTIONS For control of meadow voles (*Microtus pennsylvanicus*) prairie voles (*M ochrogaster*) pine voles (*M pinetorum*) long tailed voles (*M longicaudus*) California voles (*M californicus*) Oregon voles (*M oregoni*) mountain voles (*M montanus*) Townsend s voles (*M townsendii*) white footed mice (*Peromyscus leucopus*) and deer mice (*P maniculatus*) in barley Do not apply this product to barley within 50 days of harvest This use is restricted to Montana, Idaho, Oregon and Washington

BAITING

BROADCAST BAITING This product may be broadcast by air or ground driven dispensing devices Apply bait at a rate up to 6 lbs/A (0 11 lb a i /A) A second application may be made after 25 days Apply bait at a maximum rate of 12 lbs/A (0 22 lb a i /A) per year All applications must be made prior to the boot stage

POTATOES

USE RESTRICTIONS For control of meadow voles (*Microtus pennsylvanicus*) prairie voles (*M ochrogaster*) pine voles (*M pinetorum*) long tailed voles (*M longicaudus*) California voles (*M californicus*) Oregon voles (*M oregoni*) mountain voles (*M montanus*) Townsend s voles (*M townsendii*) white footed mice (*Peromyscus leucopus*) and deer mice (*P leucopus*) in potatoes Do not apply this product to potatoes within 30 days of harvest This use is restricted to Idaho, Oregon and Washington

BAITING

BROADCAST BAITING This product may be broadcast by air or ground driven dispensing devices Apply bait once at a rate up to 10 lbs/A (0 18 lb a i /A) or make two separate applications at a rate of up to 5 lbs/A (0 09 lb a i /A) Apply bait at a maximum rate of 10 lbs/A (0 18 lb a i /A) per year

SUGAR BEETS

USE RESTRICTIONS For control of meadow voles (*Microtus pennsylvanicus*) prairie voles (*M ochrogaster*) pine voles (*M pinetorum*) long tailed voles (*M longicaudus*) California voles (*M californicus*) Oregon voles (*M oregoni*) mountain voles (*M montanus*) Townsend s voles (*M townsendii*) white footed mice (*Peromyscus leucopus*) and deer mice (*P leucopus*) in sugar beets

Do not apply this product to sugar beets within 30 days of harvest This use is restricted to Idaho, Oregon and Washington

BAITING

BROADCAST BAITING Apply bait by ground driven dispensing devices at rates up to 10 lbs/A (0 18 lb a i /A) at planting A second application may be made at the same rate Apply bait at a maximum rate is of 20 lbs/A (0 36 lb a i /A) per year

AERIAL APPLICATION Apply bait by air at rates up to 10 lbs/A (0 18 lb a i /A) at planting A second application may be made at the same rate Apply bait at a maximum rate is of 20 lbs/A (0 36 lb a i /A) per year

WHEAT

USE RESTRICTIONS For control of meadow voles (*Microtus pennsylvanicus*) prairie voles (*M ochrogaster*) pine voles (*M pinetorum*) long tailed voles (*M longicaudus*) California voles (*M californicus*) Oregon voles (*M oregoni*) mountain voles (*M montanus*) Townsend s voles (*M townsendii*) white footed mice (*Peromyscus leucopus*) and deer mice (*Peromyscus leucopus*) in wheat Do not apply this product to wheat within 50 days of harvest This use is restricted to Montana, Idaho, Oregon and Washington

BAITING

BROADCAST BAITING Apply bait by ground driven dispensing devices at a rate of up to 6 lbs/A (0 11 lb a i /A) Two applications may be made per year separated by a minimum interval of 25 days Apply bait at a maximum rate of 12 lbs/A (0 22 lb a i /A) per year

AERIAL APPLICATION Apply bait by air at a rate of up to 6 lbs/A (0 11 lb a i /A) Two applications may be made per year separated by a minimum interval of 25 days Apply bait at a maximum rate of 12 lbs/A (0 22 lb a i /A) per year

BEANS (DRY)

USE RESTRICTIONS For control of meadow voles (*Microtus pennsylvanicus*) prairie voles (*M ochrogaster*) pine voles (*M pinetorum*) long tailed voles (*M longicaudus*) white footed mice (*Peromyscus leucopus*) and deer mice (*P leucopus*) in dry beans Do not apply by air Do not apply this product to dry beans within 30 days of harvest This use is restricted to Idaho, Oregon and Washington

BAITING

BROADCAST BAITING This product may be broadcast by hand or ground driven dispensing devices at a rate up to 6 lbs/A (0 11 lb a i /A) No more than one application per growing season is permitted

AIRPORTS (RUNWAYS AND GRASSES), HIGHWAY MEDIANS, AND TURF

USE RESTRICTIONS For control of meadow voles (*Microtus pennsylvanicus*) prairie voles (*M ochrogaster*) pine voles (*M pinetorum*) long tailed voles (*M longicaudus*) California voles (*M californicus*) Oregon voles (*M oregoni*) mountain voles (*M montanus*) Townsend s voles (*M townsendii*) white footed mice (*Peromyscus leucopus*) and deer mice (*P maniculatus*) at airports on highway medians and on turf Do not apply bait to bare ground Do not apply bait by air

BAITING

HAND BAITING Place 1 teaspoon (4 grams) of bait in surface runways or at the mouths of holes leading to underground burrow systems Allow bait to fall through to the ground surface but do not apply bait to bare ground Do not place bait in heaps or piles Apply bait at a rate up to 5 lbs/A (0 09 lb a i /A)

TRAILBUILDER BAITING Set equipment to drop 1 teaspoon (4 grams) of bait at 4 to 5 foot intervals in the artificial trail made by the machine Apply the bait at a rate up to 3 lbs/A (0 06 lb a i /A)

GROUND BROADCAST BAITING Broadcast bait evenly by hand by cyclone seeder or by mechanical spreader mounted on a tractor or 4 wheel drive all terrain vehicle (ATV) Apply treated bait at a rate up to 10 lbs/A (0 18 lb a i /A)

NONRESIDENTIAL LAWNS, GOLF COURSES AND PARKS

USE RESTRICTIONS For control of meadow voles (*Microtus pennsylvanicus*) prairie voles (*M ochrogaster*) pine voles (*M pinetorum*) long tailed voles (*M longicaudus*) California voles (*M californicus*) Oregon voles (*M oregoni*) mountain voles (*M montanus*) Townsend s voles (*M townsendii*) white footed mice (*Peromyscus leucopus*) and deer mice (*P maniculatus*) in nonresidential lawns golf courses and parks

BAITING

HAND BAITING Place bait either on surface trails or at mouth of holes leading to underground burrow systems Do not allow bait to be exposed on bare ground Do not disturb the runway system Apply at the rate of 1 teaspoon (4 grams) per placement up to 5 lbs/A (0 09 lb a i /A)

BAIT STATIONS Place bait in tamper resistant bait stations if children pets or nontarget mammals or birds may access the bait

STORAGE AND DISPOSAL

Do not contaminate water food or feed by storage or disposal

STORAGE Store only in original container in a dry place inaccessible to children pets and domestic animals Keep container closed and away from other chemicals

PESTICIDE DISPOSAL Pesticide wastes are toxic Improper disposal of excess pesticide spilled bait or rinsate is a violation of Federal law If these wastes cannot be disposed of by use according to label instructions contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance

Container Handling Nonrefillable container Do not reuse or refill this container Offer for recycling if available Completely empty the bag by shaking and tapping sides and bottom to loosen clinging particles Empty residue from bag into application equipment If container is not to be recycled then dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities

ENDANGERED SPECIES CONSIDERATIONS

Whooping Crane (*Grus americana*) Do not use this product in habitats occupied or occasionally visited by whooping cranes during the period from 30 days before the expected arrival of cranes to 30 days after the time of their usual departure

Attwater's Greater Prairie Chicken (*Tympanuchus cupido attwateri*) Do not use this product in the present occupied range of the Attwater's Greater Prairie Chicken in the following counties in Texas: Aransas, Austin, Brazoria, Colorado, DeWitt, Ft. Bend, Galveston, Goliad, Harris, Refugio, Victoria, Waller, and Wharton

Yellow Shouldered Blackbird (*Agelaius xanthomus*) and Puerto Rican Plain Pigeon (*Columba inornata wetmorei*) Do not use in the following critical habitat in Puerto Rico except in tamper proof bait boxes

- (1) within 6.3 miles (10 kilometers) of Central Aquirre, Lago Cidra, Ceiba, San German
- (2) within 9.4 miles (15 kilometers) of La Esperanza south of Highway 2 from city of Mayaguez to the city of Ponce and all Mona Island

Utah Prairie Dog (*Cynomys parvidens*) Do not use this product in critical habitat of the Utah Prairie Dog (Utah)

Salt Marsh Harvest Mouse (*Reithrodontomys raviventris*) Do not use this product in critical habitat within 0.5 miles of salt marsh vegetation and/or brackish water wetlands which are located 1) near or adjacent to San Pablo Bay and San Francisco Bay or 2) in the Sacramento River below or adjacent to confluence of the Sacramento River and the San Joaquin River (California)

Morro Bay Kangaroo Rat (*Dipodomys heermanni morroensis*) Do not use this product in critical habitat within 2.5 miles of Baywood Park which is located on Morro Bay (California)

Northern Idaho Ground Squirrel (*Spermophilus brunneus brunneus*) Do not use this product within 0.5 miles in critical habitat of the Northern Idaho Ground Squirrel in Adams and Valley counties Idaho

Oldfield mice (*Peromyscus polionotus*) threatened or endangered subspecies Do not use this product in areas of Alabama and Florida that are occupied by threatened or endangered subspecies of the oldfield mouse

**UNITED STATES DEPARTMENT OF AGRICULTURE
Animal and Plant Health Inspection Service
Riverdale MD 20737 1237**

**EPA Reg No 56228 3
EPA Est No 56228 ID 1**

**Net Contents
Batch Code No**