

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER

Keep away from humans, domestic animals, and pets. May be fatal or harmful if swallowed. Avoid eye or skin contact. Do not breathe dust. Avoid contamination of feed or food stuffs. Prevent the contact of bait with acids. Wash hands thoroughly with soap and water after use.

ENVIRONMENTAL HAZARDS

This product is toxic to wildlife and fish. Birds and other wildlife feeding in treated areas may be killed. Use with care when applying in areas frequented by wildlife or adjacent to any body of water. Keep out of lakes, ponds, or streams. Do not apply over bodies of water, in areas inhabited by livestock, or where a hazard exists to rare or endangered species. Do not contaminate water by cleaning of equipment or disposal of wastes.

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicators certification.

ZINC PHOSPHIDE CONCENTRATE FOR MOUSE CONTROL

For the control of meadow and pine voles, and white-footed mice in orchards and groves.

ACTIVE INGREDIENT:

Zinc Phosphide 63.2%

INACTIVE INGREDIENTS: 36.8%

TOTAL 100.0%

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store only in original container, in a dry, place inaccessible to children, pets and domestic animals.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spilled bait, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities.

ENDANGERED SPECIES CONSIDERATIONS

Whooping Crane (*Grus americana*)
Do not use this product 30 days prior to arrival and 30 days after the whooping crane leaves its known critical habitat.

Attwater's Greater Prairie Chicken (*Tympanuchus cupido attwateri*)
Do not use this product in critical habitat of the Attwater's greater prairie chicken in the following Texas counties: Aransas, Austin, Brazoria, Colorado, De Witt, Ft. Bend, Galveston, Goliad, Harris, Refugio, Victoria, Waller, and Wharton.

Yellow-Shouldered Blackbird (*Agelaius xanthomus*) and Puerto Rican Plain Pigeon (*Columba inornata wetmorei*)
Do not use in critical habitat except in tamper-proof bait boxes within 6.3 miles of central Aquirre Lago Cicra Ceiba, San German, 9.4 miles of La Esperanza, south of highway 2 from city at Mayaguez to the city of Ponce, and all of Mona Island, Puerto Rico.

Utah Prairie Dog (*Cynomys parvidens*)
Do not use this product in critical habitat of the Utah prairie dog (Utah).

Salt Marsh Harvest Mouse (*Reithrodontomys raviventris*)
Do not use this product in critical habitat within 0.5 mile of salt marsh vegetation and/or brackish water wetlands which are located: 1) near or adjacent to San Pablo Bay and San Francisco Bay, or 2) in the Sacramento River below or adjacent to the confluence of the Sacramento River and the San Joaquin River (California).

Morro Bay Kangaroo Rat (*Dipodomys heermanni morroensis*)
Do not use this product in critical habitat within 2.5 miles Baywood Park which is located on Morro Bay (California).

Aleutian Canada Goose (*Branta canadensis leucopareia*)
Do not use this product from October to March in the federally closed goose hunting areas in the State of California. See current duck and goose hunting game laws for current map.

KEEP OUT OF REACH OF CHILDREN DANGER-POISON



STATEMENT OF PRACTICAL TREATMENT

Any persons applying zinc phosphide products and experiencing signs or symptoms such as nausea, abdominal pain, tightness in the chest or weakness, should be seen by a physician immediately.

IF SWALLOWED: Immediately call a Poison Control Center or physician, or transport the patient to the nearest hospital. Do not drink water. Do not administer anything by mouth or make the patient vomit unless advised to do so by a physician.

IF IN EYES: Flush with plenty of water. Get medical attention if irritation persists.

IF ON SKIN: Remove contaminated clothing and wash affected areas with soap and water.

ACCEPTED UNITED STATES DEPARTMENT OF AGRICULTURE ANIMAL INDUSTRY HEALTH INSPECTION SERVICE NATIONAL TECHNICAL SUPPORT STAFF/ADC

SEP 2 1993

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. E 56228-6

Hyattsville, MD 20782
EPA Reg No. 56228-6
EPA Est. No. 56228-ID-1

Net Weight oz

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

ORCHARDS AND GROVES

USE RESTRICTIONS: For control of meadow and pine voles (*Microtus spp.*) and white-footed mice (*Peromyscus spp.*) in orchards and groves. Apply bait after harvest while the orchard or grove is in a nonbearing phase. Do not apply on bare ground or growing crops. Wear rubber gloves when handling bait. Do not graze animals in treated areas.

MIXING: Use concentrates for making baits in open areas, especially if the baiting material is wet. Do not breathe any dry bait. Use respiratory protection when mixing baits. Due to compaction of contents, can should be shaken before opening. Gradually mix one level teaspoon with 1 quart cubed apples (1/2" cubes). Bait material resembling human food must be altered in form by crushing, baling, dicing, or pelleting so that it is not readily recognizable food. Keep products away from children during bait preparation, storage, and transport.

Keep products away from irresponsible individuals, domestic animals, pets, and poultry.

BAITING: Place 1 or 2 cubes at frequent intervals in active trails under the cover of grass near the base of infested trees. Do not place the bait on bare ground. Do not retreat the same area with zinc phosphide at less than 30-day intervals. Do not exceed application rates specified on label.