PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS CANGER

Harmful if swallowed, inhaled, or absorbed through the skin. Avoid contact with eyes, skin, or clothing. Handle only with protective gloves, clothing, and face mask, or respirator. Wash hands with soap and water after handling.

ENVIRONMENTAL HAZARDS

This product is toxic to birds. Do not expose in areas accessible to waterfowl, poultry, and other non-target birds. Keep out of lakes, ponds, streams, tidal marshes, and estuaries. Do not apply where runoff is likely to occur. Do not contaminate water by the cleaning of equipment or disposal of waste.

ENDANGERED SPECIES CONSIDERATIONS

Before undertaking any control operations with the product, consult with local, State, and Federal wildlife authorities to ensure the use of this product presents no hazard to any endangered species.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

GENERAL INFORMATION: Compound DRC-1339 Concentrate is a slow acting avicide for the control of blackbirds, cowbirds, grackles, and starlings. Birds ingesting treated bait(s) die within one to three days.

USE RESTRICTIONS

Application of treated bait(a) in any problem area ab aid be made only after careful observation of bird feeding habits to locate preferred feeding sites, determine the optimum tim: of application, and evaluate hazards of the application to desirable or protected animals. DO NOT apply bait(a) in areas where there is danger of consumption by endangered species. Prebaiting may be necessary to obtain successful results. When baiting is completed, remove all unconsumed bait material and dispose of in accordance with applicable state or federal laws. Carcasses of dead or dying birds that are found should be collected and either burned or buried according to applicable laws.

NOTE: WHEN CONTROLLING BLACKBIRDS, COWBIRDS, AND GRACKLES, IT MAY BE NECESSARY TO OBTAIN A KILL-PERMIT FROM THE U.S. FISH AND WILDLIFE SERVICE AND/OR THE APPLICABLE STATE WILDLIFE AGENCY.

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicators certification.

For use only by U.S. Department of Agriculture personnel trained in bird control or persons under their direct supervision.

COMPOUND DRC-1339 CONCENTRATE

BEST AMAILALLE COPY

ACTIVE INGREDIENT:

KEEP OUT OF REACH OF CHILDREN

DANGER-POISON



STATEMENT OF PRACTICAL TREATMENT IF SWALLOWED: CALL A PHYSICIAN OR POISON CONTROL CENTER IMMEDIATELY!

IF SWALLOWED: Induce vomiting and immediately call a physician.

IF INHALED: Move patient from contaminated area and immediately call a physician.

IF ON SKIN OR IN EYES: Immediately flush eyes or skin with large quantities of water. Call a physician immediately.

UNITED STATES DEPARTMENT OF AGRICULTURE ANIMAL AND PLANT HEALTH INSPECTION SERVICE

EPA Est. No. 56228-ID-1

Net Weight

Under the Federal Insecticide, Fusgicide, and Rodenticide Act. as amended, for the pesticide registered under EPA Rea. No. 66248-10

MAR 2 5 1992

STORAGE AND DISPOSAL

Do not contaminate water, fund, or feet by storage or disposal.

STORAGE: Store only in critical container, in a dry place inaccessible to children, pets and domestic animals.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess posticide, spilled built, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hezardous Waste, representative at the nearest Er's Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment, then dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities. If burned, stay out of smoke.

DIRECTIONS FOR USE (Cont.)

Brewer's blackbird (Euphagus cyanocaphalus)
Brown-headed cowbird (Molothrus ater)
Common grackle (Quisculus quiscula)
Red-winged blackbird (Agelaius phoeniceus)
Starling (Sturnus vulgaris)

FOR CONTROL IN FEEDLOTS

FORMULATION DIRECTIONS: Dissolve 46 grams (1.6 oz) of Compound DRC-1339 Concentrate in 600 ml (1.3 pt) of warm potable water or edible oil (the amount can be adjusted according to the bait material used to obtain uniform coverage). Pour the solution over 4.5 kg (10 Lbs.) of bait material and mix or tumble slowly until mixture appears evenly distributed and dry. Bait may consist of poultry pellets, rice, corn, or milo. Do not use bait materials that have less than 2500 particles/lb or more than 25,000 particles/lb.

Dilute treated baits with 5 to 10 parts of similar untreated bait. Baits prepared in this manner should reduce the possibility of target birds consuming more than one lethal bait and should allow more birds an opportunity to consume a lethal bait.

APPLICATION IN BEEF CATTLE FEEDLOTS: With gloved I arios, place bait in feeding stations or scatter diluted bait 'hinly and uniformly at a rate of 2.5.kg/100ml (1 lb/1009ft) over dry or frozen areas in pens and alleyways before target birds arrive in the morning. Place bait in areas inaccessible to livestock. DO NOT bait pens within the feedlot which are occupied by cattle.

APPLICATION IN POULTRY AND SWINE FEEDLOTS: Place in feeding stations or hand scatter baits winly and uniformly over dry or frozen areas at a rate of 2.5 kg/100m² (1 lb/1000m²) on the PERIMETER of the lot OUTSIDE the pen areas occupied by poultry and swine. Outside the pen within the feedlots which are occupied by authors or swine.