

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER

Harmful if swallowed, inhaled, or absorbed through the skin. Avoid contact with eyes, skin, or clothing. Handle only with protective gloves, clothing, and face mask, or respirator. Wash hands with soap and water after handling.

### ENVIRONMENTAL HAZARDS

This product is toxic to birds. Do not expose in areas accessible to waterfowl, poultry, and other non-target birds. Keep out of lakes, ponds, streams, tidal marshes, and estuaries. Do not apply where runoff is likely to occur. Do not contaminate water by the cleaning of equipment or disposal of waste.

### ENDANGERED SPECIES CONSIDERATIONS

Before undertaking any control operations with the product, consult with local, State, and Federal wildlife authorities to ensure the use of this product presents no hazard to any endangered species.

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**GENERAL INFORMATION:** Compound DRC-1339 Concentrate is a slow acting avicide for the control of blackbirds, cowbirds, grackles, and starlings. Birds ingesting treated bait(s) die within one to three days.

### USE RESTRICTIONS

Application of treated bait(s) in any problem area should be made only after careful observation of bird feeding habits to locate preferred feeding sites, determine the optimum time of application, and evaluate hazards of the application to desirable or protected animals. **DO NOT apply bait(s) in areas where there is danger of consumption by endangered species.** Prebaiting may be necessary to obtain successful results. When baiting is completed, remove all unconsumed bait material and dispose of it in accordance with applicable state or federal laws. Carcasses of dead or dying birds that are found should be collected and either burned or buried according to applicable laws.

**NOTE: WHEN CONTROLLING BLACKBIRDS, COWBIRDS, AND GRACKLES, IT MAY BE NECESSARY TO OBTAIN A KILL-PERMIT FROM THE U.S. FISH AND WILDLIFE SERVICE AND/OR THE APPLICABLE STATE WILDLIFE AGENCY.**

## RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicators certification.

For use only by U.S. Department of Agriculture personnel trained in bird control or persons under their direct supervision.

## COMPOUND DRC-1339 CONCENTRATE - FEEDLOTS

For bird control in feedlots

### ACTIVE INGREDIENT:

3-Chloro-4-methylbenzenamine hydrochloride... 98.0%

INERT INGREDIENTS:..... 2.0%

TOTAL: ..... 100.0%

KEEP OUT OF REACH OF CHILDREN

DANGER--POISON



### STATEMENT OF PRACTICAL TREATMENT

**IF SWALLOWED: CALL A PHYSICIAN OR POISON CONTROL CENTER IMMEDIATELY!**

**IF SWALLOWED:** Induce vomiting and immediately call a physician.

**IF INHALED:** Move patient from contaminated area and immediately call a physician.

**IF ON SKIN OR IN EYES:** Immediately flush eyes or skin with large quantities of water. Call a physician immediately.

UNITED STATES DEPARTMENT OF AGRICULTURE  
ANIMAL AND PLANT HEALTH INSPECTION SERVICE

Hyattsville, MD 20782  
EPA Reg. No. 56228-10  
EPA Est. No. 56228-ID-1

Net Weight

**BEST AVAILABLE COPY**

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**STORAGE:** Store only in original container, in a cool, dry place, away from children, pets and domestic animals.

**PESTICIDE DISPOSAL:** Pesticide wastes are acutely toxic. Improper disposal of excess pesticide, spray mixture, or empty container is a violation of Federal law. If these wastes cannot be disposed of by the applicator, contact your nearest Environmental Control Agency, or the Hazardous Waste Division of the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Completely empty container and bottom to loosen clinging particles. Wash application equipment, then dispose of bags in incineration if allowed by State and local authorities. Do not burn.

## DIRECTIONS FOR USE

Brewer's blackbird (*Euphagus cyanocephalus*)  
Brown-headed cowbird (*Molothrus ater*)  
Common grackle (*Quiscalus quiscula*)  
Red-winged blackbird (*Agelaius phoeniceus*)  
Starling (*Sturnus vulgaris*)

FOR CONTROL IN FEEDLOTS

### FORMULATION DIRECTIONS:

**Roller Barley, Cracked Corn, Whole Poultry Pellets (1.8 - 2.0% bait):** Dissolve 1.8 to 2.0% of Compound DRC-1339 Concentrate (1.3 pt) of warm potable water at 43°F solution over 4.5 kg (10 lbs) of bait and tumble slowly until mixture appears evenly dry at ambient temperature. Screen bait materials that have less than 5000 particles/5,000 particles/lb. Mix treated bait with similar untreated bait. Use treated

**Whole Raisins (0.18 - 0.2% bait):** Dissolve 0.18 to 0.2% of Compound DRC-1339 Concentrate in warm potable water or edible oil at 43°F solution over 4.5 kg (10 lbs) of raisins and tumble slowly until mixture appears evenly dry at ambient temperature. Distribute raisins. Post the treated area with watering operation. Use treated raisins with

**Roller Whole Corn (0.9 - 1.0% bait):** Dissolve 0.9 to 1.0% of Compound DRC-1339 Concentrate in 600 ml (1.3 pt) of warm potable water (110°F). Pour the solution over 4.5 kg whole corn and mix or tumble slowly until evenly distributed. Air dry at ambient temperature. Distribute corn. Post the treated area with watering operation. Use treated corn within 7 days.

SEE REVERSE SIDE FOR ADDITIONAL DIRECTIONS FOR USE

ACCEPTED  
MAY 14 1992  
For the Federal Insecticide, Fungicide, and Rodenticide Act, issued under Reg. No. 56228-10

**Cull French Fries (0.18 - 0.2% bait):** Dissolve 46 grams (1.6 oz) of Compound DRC-1339 Concentrate - Feedlots in 600 ml (1.3 pt) of warm potable water or edible oil at 43°C (110°F). Pour the solution over 23 kg (50 Lbs.) of cull French fries and mix or tumble slowly until mixture appears evenly distributed. Air dry at ambient temperature. Dilute at no less than 1:2 with untreated cull French fries. Post the area with warning signs during the baiting operation. Use treated French fries within 24 hours.

**Note:** For any bait prepared from this product, it is important to dilute treated bait particles with untreated particles at the appropriate ratio (see directions above). Diluting baits should reduce the possibility of any target bird consuming more than one lethal bait and should allow more birds an opportunity to consume a lethal bait.

**APPLICATION IN BEEF CATTLE FEEDLOTS:** With gloved hands, place bait in feeding stations or scatter diluted bait thinly and uniformly at a rate of 2.5 kg/100m<sup>2</sup> (1 lb/1000ft<sup>2</sup>) over dry or frozen areas in pens and alleyways before target birds arrive in the morning. Place bait in areas inaccessible to livestock. **DO NOT** bait pens within the feedlot which are occupied by cattle.

**APPLICATION IN POULTRY, SWINE, AND DAIRY CATTLE FEEDLOTS:** Place in feeding stations or hand scatter baits thinly and uniformly over dry or frozen areas at a rate of 2.5 kg/100m<sup>2</sup> (1 lb/1000ft<sup>2</sup>) on the PERIMETER of the lot **OUTSIDE** the pen areas occupied by poultry, swine, or dairy cattle. **DO NOT** bait pens within the feedlots which are occupied by poultry, swine, or dairy cattle.

**BEST AVAILABLE COPY**