

PRECAUTIONARY STATEMENTS

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Keep away from humans, domestic animals, and pets. May be fatal or harmful if swallowed. Avoid eye or skin contact. Do not breathe dust. Avoid contamination of feed or food stuffs. Prevent the contact of bait with acids. Wash hands thoroughly with soap and water after use. Use rubber gloves if contact with skin occur when mixing or placing bait. Wash all utensils, spoons, or measuring devices thoroughly.

ENVIRONMENTAL HAZARDS

This product is toxic to wildlife and fish. Birds and other wildlife feeding in treated areas may be killed. Use with care when applying in areas frequented by wildlife or adjacent to any body of water. Keep out of lakes, ponds, or streams. This product shall not be applied over bodies of water, in areas inhabited by livestock, or where a hazard exists to rare or endangered species. Do not contaminate water by cleaning of equipment or disposal of wastes.

ENDANGERED SPECIES CONSIDERATIONS

Whooping Crane (*Grus americana*)
Do not use this product 30 days prior to arrival and 30 days after the whooping crane leaves its known critical habitat.

Utah Prairie Dog (*Cynomys parvidens*)
Do not use this product in critical habitat of the Utah prairie dog (Utah)

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicators certification.

**ZINC PHOSPHIDE
ON STEAM-ROLLED OATS
FOR PRAIRIE DOG CONTROL**

For the control of black-tailed prairie dogs, white-tailed prairie dogs, and Gunnison's prairie dogs on rangelands in the following states: North Dakota, South Dakota, Nebraska, Kansas, Oklahoma, Texas, New Mexico, Arizona, Colorado, Montana, Utah, and Wyoming.

ACTIVE INGREDIENT:

Zinc Phosphide 2.0%

INACTIVE INGREDIENTS: 98.0%

TOTAL 100.0%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: CALL A PHYSICIAN OR POISON CONTROL CENTER IMMEDIATELY!

Any person applying zinc phosphide products and experiencing signs or symptoms such as nausea, abdominal pain, tightness in the chest or weakness, should be seen by a physician immediately.

If swallowed, drink 1-2 glasses of water and induce vomiting by touching the back of the throat with the finger. Avoid the use of all oils. Have person lie down and keep warm. Do not induce vomiting or give anything by mouth to an unconscious person.

If on skin or eyes, flush with plenty of water.

UNITED STATES DEPARTMENT OF AGRICULTURE
ANIMAL AND PLANT HEALTH INSPECTION SERVICE
NATIONAL TECHNICAL SUPPORT STAFF/ADC

Hyattsville, MD 20782
EPA Est. No. 56228-ID-1
EPA Reg. No. 56228-14

Net Weight lbs.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store only in original container, in a dry place inaccessible to children, pets and domestic animals.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spilled bait, or residue is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS: For control of black-tailed (*Cynomys ludovicianus*), white-tailed (*C. leucurus*), and Gunnison's prairie dogs (*C. gunnisoni*), on rangeland in Western United States (North Dakota, South Dakota, Nebraska, Kansas, Oklahoma, Texas, New Mexico, Arizona, Colorado, Montana, Utah, and Wyoming) where non-target species (especially black footed ferrets) will not be endangered.

PREBAITING: Prebait with 4 grams (1 teaspoon) of untreated steam-rolled oats per mound, 1-2 days prior to baiting, to increase acceptance of treated baits by prairie dogs.

BAITING: Apply bait only after all or most of prebait is eaten and only to areas where untreated bait was consumed. Establish observation period during prebaiting. Apply bait by hand as a six-inch bait spot on edge of each mound or in adjacent feeding areas. Application rate should not exceed 4 grams (1 teaspoon) per bait spot. Treat during late summer or fall period (July-January). Do not apply more than 1 bait application during this period. Dispose of spilled or unwanted bait by burial.

ACCEPTED

AUG 31 1983

Under the Animal and Plant Health Inspection Service
Rodenticide Act, as amended, for the
pesticide registered under
EPA Reg. No. 56228-14

