



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

October 20, 2015

Jim Warren
USDA APHIS Environmental and Risk Analysis Services, Unit 149
4700 River Road
Riverdale, MD 20737

Subject: Label Amendment –Per Registration Review for Compound DRC-1339
Product Name: Compound DRC-1339 Concentrate-Livestock Nest & Fodder
Depredations
EPA Registration Number: 56228-29
Application Date: July 10, 2015
Decision Number: 509984

Dear Mr. Warren:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance

Page 2 of 2
EPA Reg. No. 56228-29
Decision No. 509984

with FIFRA section 6. If you have any questions, please contact Gene Benbow by phone at 703-347-0235, or via email at Benbow.Gene@epa.gov.

Enclosure

Sincerely,

A handwritten signature in black ink, appearing to read "Mark Suarez", with a stylized flourish at the end.

Mark Suarez, Product Manager 07
Invertebrate & Vertebrate Branch 3
Registration Division (7505P)
Office of Pesticide Programs

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS **DANGER**

Acute Hazards: Fatal if inhaled. Corrosive. Causes irreversible eye damage and skin burns. May be fatal if swallowed. Harmful if absorbed through skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some people.

Hazard Avoidance: Do not get in eyes, on skin, or on clothing. Do not breathe dust. Wear protective clothing, eyewear, and respiratory protection as listed under "**PERSONAL PROTECTIVE EQUIPMENT.**" Wash thoroughly with soap and water after handling and before eating or smoking. Remove contaminated clothing and wash before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Handlers who mix packages containing 1 lb (0.45 kg) or more of this product must wear:

- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant gloves (such as waterproof or rubber gloves)
- Chemical-resistant footwear plus socks
- Protective eyewear (goggles or face shield)
- A NIOSH approved particulate respirator with any N, R, or P filter with NIOSH approval number prefix TC-84A

Handlers who mix packages containing less than 1 lb (0.45 kg) of this product must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves (such as waterproof or rubber gloves)
- Protective eyewear (goggles or face shield)

Applicators who handle bait must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves (such as waterproof or rubber gloves)

User Safety Requirements:

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions are provided for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

RESTRICTED USE PESTICIDE

Due to High Acute Inhalation Toxicity and Eye and Skin Corrosiveness to Humans; High Acute Toxicity to Nontarget Birds and Aquatic Invertebrates; and the Need for Highly Specialized Applicator Training.

For retail sale to and use only by USDA APHIS Certified Applicators trained in bird control or by persons under their direct supervision.

COMPOUND DRC-1339 CONCENTRATE – Livestock, Nest & Fodder Depredations

For control of crows, ravens, and magpies that prey on newborn livestock, that prey on eggs or the young of Federally-designated Threatened or Endangered Species or of other species designated to be in need of special protection or that damage and feed on the contents of silage/fodder bags.

Active Ingredient:
DRC-1339, 3-chloro-p-toluidine hydrochloride:97.0%
Other Ingredients: 3.0%
TOTAL:100.0%

KEEP OUT OF REACH OF CHILDREN

DANGER-PELIGRO POISON



If swallowed	<ul style="list-style-type: none"> - Call a poison control center or doctor immediately for treatment advice. - Have person sip a glass of water if able to swallow. - Do not induce vomiting unless told to do so by the poison control center or doctor.
--------------	---

If on skin or clothing	<ul style="list-style-type: none"> - Take off contaminated clothing. - Rinse skin immediately with plenty of water for 15-20 minutes. - Call a poison control center or doctor immediately for advice.
------------------------	---

If inhaled	<ul style="list-style-type: none"> - Move person to fresh air. - If person is not breathing, call an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. - Call a poison control center or doctor immediately for advice.
------------	---

If in eyes	<ul style="list-style-type: none"> - Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. - Call a poison control center or doctor immediately for advice.
------------	--

NOTE TO PHYSICIAN AND VETERINARIAN: Probable mucosal damage may contraindicate the use of gastric lavage. See additional "**PRECAUTIONARY STATEMENTS**" on Left Panel. If pet eats bait, call a veterinarian at once.

ENVIRONMENTAL HAZARDS:

This product is very highly toxic to birds and aquatic invertebrates. **Do not** use in any manner that may endanger nontarget and protected bird species. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not apply when runoff is likely to occur. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. **Do not** contaminate water by the cleaning of equipment or disposal of waste.

ENDANGERED SPECIES CONSIDERATIONS:

Before undertaking any control operations with the product, consult with local, State, and Federal Wildlife authorities to ensure the use of this product presents no hazard to any Threatened or Endangered Species. **DO NOT** apply treated baits where there is a danger that Threatened or Endangered Species will consume baits unless special precautions are taken to limit such exposures.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL:

Read the entire label. This product must be used strictly in accordance with this label's precautionary statements and use directions, as well as with all applicable State and Federal laws and regulations.

Before using this product, contact the U.S. Fish and Wildlife Service and the applicable State wildlife agency and obtain all necessary kill or collecting permits. Use only for the sites, pests, and application methods described on this label.

PRODUCT INFORMATION:

This product contains a slow-acting avicide which kills target bird species (see list below) in 1 to 3 days. As many types of nontarget birds are potentially vulnerable to DRC-1339, it is necessary to use care and to follow the requirements of this label to minimize impacts to nontarget species.

(See next page for additional "DIRECTIONS FOR USE")

UNITED STATES DEPARTMENT OF AGRICULTURE ANIMAL AND PLANT HEALTH INSPECTION SERVICE

**4700 River Road, Unit 149
Riverdale, MD 20737**

**EPA Reg. No. 56228-29
California Reg. No 56228-29-AA
EPA Est. No. 56228-ID-1
Net Contents: _____
Batch Code No.: _____**

DIRECTIONS FOR USE, continued

USE RESTRICTIONS:

Baits made from Compound DRC-1339-Livestock, Nest & Fodder Depredations may only be used to control the following species: common raven (*Corvus corax*), Chihuahuan raven (*Corvus cryptoleucus*), American crow (*Corvus brachyrhynchos*), Black-billed magpie (*Pica hudsonia*), and Fish crow (*Corvus ossifragus*). This product may be used to prepare egg or meat-cube baits to control the target species listed above in the following use sites:

- Rangeland and pastureland areas where ravens or crows prey upon newborn livestock;

- Refuges or other areas where ravens or crows prey upon the eggs and/or young of Federally-designated Threatened or Endangered Species, or upon the eggs and young of other species which Federal or State wildlife agencies have determined to be in need of protection from nest predators due to documented declines in numbers and/or in nesting success; or

- Within 25 feet (7.6 m) of silage/fodder bags that have been damaged or are likely to be damaged by crows, ravens, or black-billed magpies.

Baits must be prepared and applied as specified on this label. DO NOT apply baits made from this product by air or by use of any mechanical equipment designed to broadcast baits or other pesticides. Users of this product must follow all limitations indicated on this label regarding the placement and monitoring of treated baits.

Before baits made from this product are applied, sites that are to be treated must be observed for evidence of nontarget activity **and must be prebaited** (see specific instructions for these activities). **DO NOT** apply treated baits where there is a danger that Threatened or Endangered Species will consume baits unless special precautions are taken to limit such exposures. Such precautions shall include observation of baited sites and use of hazing tactics to frighten away Threatened or Endangered Species that otherwise might feed upon baits.

DIRECTIONS FOR USE, continued

USE RESTRICTIONS, continued:

DO NOT apply treated baits within 50 feet (15.2 m) of permanent manmade or natural bodies of water, unless baited sites are under constant observation while baits are exposed.

DO NOT exceed a maximum application rate of 0.083 lbs of active ingredient per treated acre (0.93 g active ingredient/100 m²). That rate is equivalent to 18 treated-egg baits in at least 5 bait sets applied over an area of 400 ft² (37.2 m²) surrounding an animal carcass draw station.

DO NOT store, apply, or even temporarily place treated bait in locations accessible to children, pets, domestic animals, or nontarget wildlife.

DO NOT apply bait in a way that will contact workers or other persons.

DO NOT use treated baits as food or feed.

DO NOT apply baits made from this product in any way that could contaminate human food or animal feed.

ENTRY RESTRICTIONS:

Only protected applicators may be in the area during bait application. Keep pets and livestock, and persons other than authorized handlers away from the bait at all times, and exclude all unauthorized persons from application sites during prebaiting and baiting. For example, post signage near, in the vicinity of, or at main entrances or commonly used access points to prebaiting and baiting sites that warns persons not to pick up or handle any baits and to keep pets and livestock away from bait.

PRETREATMENT OBSERVATIONS:

Prior to application, carefully observe target birds' feeding habits to locate their preferred feeding sites, determine the optimum time of application, and evaluate potential hazards of the application to nontarget and protected species.

PREBAITING:

Prebaiting with untreated bait materials (or use of a draw station) is necessary to promote feeding by target species and to assess potential for exposure of nontarget species. Apply prebait using the same procedures that are prescribed below for the type of bait ("**EGG BAITS**" or "**MEAT BAITS**") that is to be used for toxic baiting.

Observe baited areas (from blinds) early in prebaiting period to determine whether nontarget species are approaching baits. Haze away Threatened or Endangered and nontarget species that might consume baits. Remove baits if such nontarget species continue to approach them.

DIRECTIONS FOR USE, continued

BAIT PREPARATION:

MEAT BAITS:

MEAT BAIT PREPARATION:

1. Mix 0.027 oz (0.75 g) of this product with 0.18 oz (5.0 g) of powdered sugar.
2. Pour or sprinkle concentrate-sugar mixture over 200 meat cubes that measure about 0.5 in (1.3 cm) on each side.
3. Mix or tumble bait slowly until all meat cubes appear to be evenly covered.

MEAT BAIT APPLICATION:

NOTE: During application, wear all PPE as listed under "PERSONAL PROTECTIVE EQUIPMENT."

Control of crows, magpies, and ravens with meat baits prepared from this product is limited to the sites indicated above under "**USE RESTRICTIONS.**" Wear rubber gloves while handling baits. Place no more than 75 meat cube baits at each baited site. Place 5 to 10 baits in clusters over an area not to exceed 1000 ft² (93 m²) where control of ravens, magpies, and/or crows is to be affected. **Draw stations** (fresh, unpoisoned animal carcasses) may be needed to attract ravens, magpies, and/or crows to the locations selected for bait exposure. If draw stations are used, place meat baits on or within a few feet of the animal carcasses.

WHILE TREATED MEAT BAITS ARE EXPOSED, BAITED AREAS MUST BE OBSERVED CONTINUOUSLY FROM A DISTANCE OF NO MORE THAN 1000 YARDS (914 m) TO DETECT APPROACHES BY THREATENED OR ENDANGERED SPECIES AND OTHER NONTARGET OR PROTECTED ANIMALS LIKELY TO EAT BAITS.

Because of wariness of target bird species, it may be necessary to observe baits from behind natural or specially-constructed blinds. Haze away Threatened or Endangered and nontarget species that might consume baits. Remove baits if such nontarget species continue to approach them.

Unconsumed bait cubes must be retrieved daily, at the conclusion of each observation period and no later than one hour after sunset. Dispose of retrieved baits in accordance with applicable State and Federal laws.

DIRECTIONS FOR USE, continued

EGG BAITS:

EGG BAIT PREPARATION:

Dissolve 0.07 oz (2 g) of the product in 0.2 pint (100 ml) of warm potable water at 110 °F (43.3 °C) to make an approximately 2% solution; or dissolve 0.14 oz (4 g) of the product in 0.2 pint (100 ml) of warm potable water at 110 °F (43.3 °C) to make an approximately 4% solution; or in other proportions to produce a 2% or 4% solution.

Using an 18-gauge hypodermic needle or similarly-sized implement, make an entry hole in the end of each hard-boiled chicken, turkey, or duck egg to be used. Using a syringe and a 20-gauge hypodermic needle, slowly inject 0.002 pints (1 ml) of the 2% solution (or 0.001 pints or 0.5 ml of the 4% solution) into the yolk of each egg.

Make only enough solution to treat the desired number of eggs. Mark treated eggs with small skull and crossbones or the word POISON.

EGG BAIT APPLICATION:

NOTE: During application, wear all PPE as listed under "PERSONAL PROTECTIVE EQUIPMENT."

Control of crows, magpies, and ravens with egg baits prepared from this product is limited to the sites indicated above under "USE RESTRICTIONS". Place all egg baits to be used at one baited site within 25 ft (7.6 m) of the center of the site or within 25 ft (7.6 m) of any silage/fodder bags that are to be protected. Place 1-4 eggs in each bait set, and do not use more than a total of 18 eggs per baited site. If a draw station (fresh, unpoisoned animal carcass) is used, all bait sets must be located at least 10 ft (3 m) from the carcass. Wherever practical, bait sets should be made in "dummy" nests created by making small depressions in the ground. Dummy nests may be partially hidden by vegetation or other debris. In other situations, eggs may be placed on elevated wooden platforms 1 to 2 ft² (0.1 to 0.2 m²) in area. Eggs placed on platforms must be restrained by wire to prevent them from falling off platforms or being removed by birds. Apply 2-3 eggs per platform.

DO NOT USE MORE EGGS THAN ARE NEEDED TO EFFECT CONTROL, as ravens and crows tend to cache surplus food.

Observe baited areas (from blinds) early in baiting period to determine whether nontarget species are approaching egg baits. Haze away Threatened or Endangered and nontarget species that might consume baits. Remove baits if such nontarget species continue to approach them.

Rebait with additional treated eggs when more than 50% of the treated eggs offered have been removed by ravens, magpies, or crows. When replacing baits, take care not to frighten target birds actively removing or feeding upon eggs. Retrieve unconsumed treated eggs within 7 days of exposure. Old treated eggs and treated eggs not eaten by the time control operations cease must be disposed of in accordance with applicable State and Federal laws.

POSTTREATMENT CLEAN-UP (Meat and Egg Baits):

NOTE: During clean-up, wear long-sleeved shirt and long pants and chemical-resistant gloves (such as waterproof or rubber gloves). To further reduce the potential for exposure, use appropriate implements such as scoops or other tools to collect carcasses or uneaten bait.

Collect unconsumed and leftover meat daily, and unconsumed and leftover egg baits, dying birds, and carcasses within 7 days of treatment. Dispose of such baits and carcasses by burning or burial, as authorized by applicable laws and ordinances.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store only in original container, in a dry place inaccessible to children, pets, and domestic animals.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spilled bait, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Completely empty bags by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. If bags are not to be recycled, dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities. If burned, stay out of smoke.

ACCEPTED

10/20/2015

Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under
EPA Reg. No. 56228-29

56228-29