


FRONT PANEL**CHECKMATE® CM**

Active Ingredients
 (E, E)-8,10-Dodecadien-1-ol 12.36%
 Inert Ingredients 87.64%
 TOTAL 100.00%
 Net Weight: 180 mg active ingredients
 per dispenser.



A BIOCHEMICAL PHEROMONE FOR
 MATING DISRUPTION OF
 CODLING MOTH (*Cydia pomonella*)
 HICKORY SHUCKWORM (*Cydia caryana*)

Keep Out Of Reach Of Children

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes skin irritation. Avoid contact with eyes, skin or clothing. Avoid contact of pheromone reservoir inside dispenser with skin or clothing. Wash thoroughly with soap and water after handling.

FIRST AID

IF ON SKIN: Take off contaminated clothing.
 Rinse skin immediately with plenty of water for 15-20 minutes.
 Call a Poison Control center or doctor for treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes.
 Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
 Call a Poison Control Center or doctor for treatment advice.

Have the product container or label with you when calling a Poison Control Center or doctor, or going for treatment.

ENVIRONMENTAL HAZARDS

Keep out of lakes, ponds or streams. Do not contaminate water by disposal of wastes.

Manufactured by:
CONSEP®

Consep, Inc.

213 SW Columbia Street
 Bend, Oregon 97702
 1/866-326-6737
 FAX (541) 312-8298

EPA Registration No. 56336-13
 EPA Establishment No. 56336-OR-1
 Net Contents: xxx Dispensers

BACK PANEL

WARRANTY AND LIMITATION OF DAMAGES

Consep, Inc. warrants that this material conforms to the chemical description on the label. Consep, Inc. neither makes, nor authorizes any agent or representative to make, any other warranty of fitness or of merchantability, guarantee or representation, express or implied, concerning this material. Consep, Inc.'s maximum liability for breach of this warranty shall not exceed the purchase price of this product. Buyer and user acknowledge and assume all risks and liabilities resulting from the handling, storage and use of this material.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling

CheckMate® CM is a behavior modifying biochemical product (pheromone) which functions by mating disruption. It is to be used in pest management programs for control of codling moth (*Cydia pomonella*) and hickory shuckworm (*Cydia caryana*).

METHOD OF APPLICATION: For use on apples, pears, quinces, and other pome fruits; peaches, nectarines, plums, prunes, and other stone fruits the recommended rate of application of **CheckMate CM** is 120 to 160 dispensers per acre (300 to 400 dispensers per hectare). For use on walnuts, pecans, and other tree nuts the recommended rate of application of **CheckMate CM** is 150 to 200 dispensers per acre (370 to 500 per hectare). Do not exceed a total of 150 grams a.i./acre/year.

Use gloves when handling dispensers. Provide adequate ventilation when handling dispensers. Do not handle dispensers in an enclosed area. Hang dispensers from thumb-sized tree limb to secure placement. Place dispensers in the upper one-third of the tree or higher.

Wash hands after completion. Do not handle other types of insect lures before or after lure placement without thoroughly washing hands. Set codling moth traps before handling **CheckMate CM**. Monitor treatments carefully, especially for secondary pests such as leafrollers.

Reseal containers immediately after use, store unused product only in refrigerators or freezers not used for food storage. Do not use **CheckMate CM** from damaged, punctured or unsealed containers.

RECOMMENDED APPLICATION: For best results apply at the time or shortly before codling moths or hickory shuckworm moths begin to emerge in the spring (biofix), or when the moths are first detected in pheromone traps such as **BioLure®**. This product only affects adult male moths and will have no effect on female moths, eggs or larvae. If application occurs following biofix or during the growing season, the orchard must be treated with insecticide treatments that will effectively control hatching larvae until egg laying by previously mated female moths has ceased to occur. In moderate to high insect pressure situations, supplemental insecticide applications during the season may be necessary to provide adequate protection to developing fruit. These supplemental insecticide applications should be made based on trap monitoring, field scouting, and appropriate degree-day models for the local growing area and must be timed to control emerging larvae. Follow the *BioLure Technical Bulletin* for trap placement. Monitor insect infestation with traps and by visual inspection of trees and fruit. Re-apply **CheckMate CM** as needed, based on monitoring results and field scouting.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal

STORAGE: Store container in cool place until used. **PESTICIDE DISPOSAL:** Wastes resulting from use of this product may be disposed of on-site or at an approved disposal facility. **CONTAINER DISPOSAL:** Do not reuse container. Discard in trash.