

Reg # 56387-1

PM-13

194

BARRE-NATIONAL INC.

PYRINYL II LIQUID - 4 FL OZ

ACCEPTED
 MAY 19 1992
 Product Name: Pyrinyl II Liquid
 EPA Reg. No.: 56387-1
 56387-1

**READ EYE
CAUTION
STATEMENT
BEFORE
USE**

INDICATIONS:
For the treatment of infestations of head lice, pubic (crab) lice, and body lice

STORAGE AND DISPOSAL:
Do not transport or store below 32°F (0°C) Do not reuse empty container. Wrap and put in trash.

See bottom of carton for lot number and expiration date.

EPA EST. 56387-MD-1
EPA REG NO 56387-1
Rev 10/91K1

Barre

NDC 0472-1023-94



**KILLS LICE
& REMOVES THEIR EGGS**
(Head, pubic (crab), and body lice)

ACTIVE INGREDIENTS:

Pyrethrins	0.3%
Piperonyl Butoxide Technical (equivalent to 2.4% (butylcarbityl) (6-propylpiperonyl) ether and 0.6% related compounds)	3.0%

INERT INGREDIENTS: 96.7%
TOTAL 100.0%
 * Contains petroleum distillate

**FINE-TOOTHED
EGG REMOVAL
COMB ENCLOSED**

6505-01-186-8437



CAUTION:
 KEEP OUT OF REACH OF CHILDREN
 FOR EXTERNAL USE ONLY
 4 FL OZ (118 mL)

BEST AVAILABLE COPY

(Head, pubic (crab), and body lice)
**& REMOVES THEIR EGGS
KILLS LICE**

PYRINYL II LIQUID

Barre

INSTRUCCIONES EN ESPAÑOL ADJUNTAS.

Barre® PYRINYL II LIQUID

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. Shake well. Apply to affected area until all the hair is thoroughly wet with product.
2. Allow product to remain on area for 10 minutes but no longer. Wash area thoroughly with warm water and soap or shampoo.
3. A fine-toothed comb may be used to help remove dead lice or their eggs (nits) from hair.
4. A second treatment must be made in 7 to 10 days to kill any newly hatched lice. Use no more than two times in 24 hours.

See accompanying bilingual package insert for additional important information.

In order to prevent reinfestation with body lice, all clothing must be sterilized concurrent with the application of this preparation.

Manufactured by
BARRE-NATIONAL INC.
Baltimore, MD 21207 USA

**READ EYE
CAUTION
STATEMENT
BEFORE
USE**

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Use with caution on persons allergic to ragweed. For external use only. Harmful if swallowed. Do not inhale. Do not use near the eyes or permit contact with mucous membranes. If product gets into the eyes, immediately flush with water. Consult a physician if irritation of eyebrows and eyelashes occurs. If skin irritation or infection is present or develops, discontinue use and consult a physician. Avoid contamination of feed or foodstuffs.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of throat with a finger. Repeat until vomit is clear. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

If Inhaled: Remove victim to fresh air. Apply artificial respiration if indicated.

If On Skin: Remove contaminated clothing and wash affected areas with soap and water.

If In Eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

BEST AVAILABLE COPY

394

BARRE-NATIONAL INC.

PYRINYL II LIQUID - 4 FL OZ

Barre

NDC 0472-1023-94
657 5-01-106-8437

PYRINYL II LIQUID

Lice Treatment
KILLS LICE
(head, pubic (crab), and body lice)

ACTIVE INGREDIENTS

Pyrethrin 0.3%
Piperonyl Butoxide Technical 3.0%
(equivalent to 2.4% (2-allylcyclohexyl) (5-propylpiperonyl) ether
and 0.6% related compounds)

NET WEIGHT

TOTAL

* Contains petroleum distillate

98.7%

100.0%

98.7%

100.0%

CAUTION:
KEEP OUT OF REACH OF CHILDREN

Manufactured by
BARRE-NATIONAL INC.
Baltimore, MD 21207 USA

4 FL OZ (118 mL)

FOR EXTERNAL USE ONLY

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. Shake well. Apply to affected area prior to the hair is thoroughly wet with product.
2. Allow product to remain on area for 10 minutes but no longer. Wash area thoroughly with warm water and soap or shampoo.
3. A fine toothed comb may be used to help remove head lice or their eggs (nits) from hair.
4. A second treatment must be made in 7 to 10 days to kill any newly hatched lice. Use no more than two times in 24 hours.

In order to prevent reinfestation with body lice, all clothing must be sterilized concurrently with the application of this preparation.

PRECAUTIONS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION: Use with caution on persons allergic to pyrethroids. For external use only. Harmful if swallowed. Do not inhale. Do not use near the eyes or permit contact with mucous membranes. If product gets into the eyes, immediately flush with water and call a physician if irritation of eyes and eyelashes occurs. If skin irritation or infection is present or develops, discontinue use and consult a physician. Avoid consumption of food or foodstuffs.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Call a physician or Poison Control Center immediately. Drink or eat two glasses of water and induce vomiting by touching the back of throat with finger. Repeat until vomit is clear. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

If Irritated: Remove victim to fresh air. Apply artificial respiration if indicated.

If On Skin: Remove contaminated clothing and wash affected areas with soap and water.

If In Eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

STORAGE AND DISPOSAL: Do not transport or store below 32°F (0°C). Do not re-use empty container. WRAP and put in trash.

EPA EST. 58387-MD-1 EPA REG. NO. 58387-1 Rev. 10/91/1K1

BEST AVAILABLE COPY

FORM NO. 1023

PYRINYL II LIQUID - 4 FL. OZ.

TREATING LICE INFESTATION WITH

LICE TREATMENT KIT

KILLS LICE

& REMOVES THEIR EGGS

EPA Reg. No. 56387-1

This booklet will tell you about human lice and how to treat infestations effectively with this product.

Lice are tiny parasitic insects the size of a pinhead. People at all social levels can become infested with lice; no one is immune. The 3 types of lice which infest humans are:

Head Lice: Head lice live on the scalp and lay small white eggs (nits) on the hair shaft close to the scalp. The nits are most easily found on the nape of the neck or behind the ears. All personal headgear, scarfs, coats and bed linen should be disinfected by machine washing in hot water and drying, using the hot cycle of a dryer for at least 20 minutes. Personal articles of clothing or bedding that cannot be washed may be dry-cleaned, sealed in a plastic bag for a period of about 2 weeks, or sprayed with a product specifically designed for this purpose. Personal combs and brushes may be disinfected by soaking in hot water (above 130°F) for 5 to 10 minutes. Thorough vacuuming of rooms inhabited by infected patients is recommended.

Pubic (Crab) Lice: Pubic lice may be transmitted by sexual contact; therefore sexual partners should be treated simultaneously to avoid reinfestation. The lice are very small and look almost like brown or grey dots on the skin. Pubic lice usually cause intense itching and lay small white eggs (nits) on the hair shaft generally close to the skin surface. In hairy individuals, pubic lice may be present on the short hairs of the thighs and trunk, underarms, and occasionally on the beard or mustache. Underwear should be disinfected by machine washing in hot water; then drying, using the hot cycle for at least 20 minutes.

Body Lice: Body lice and their eggs are generally found in the seams of clothing, particularly in the waistline and armpit area. They move to the skin to feed, then return to the seams of the clothing where they lay their eggs. Clothing worn and not laundered before treatment should be disinfected by the same procedure as described for head lice, except that sealing clothing in a plastic bag is not recommended for body lice because the nits (eggs) from these lice can remain dormant for a period of up to 30 days.

Lice are easily transmitted from person to person by shared personal articles such as hats, hair ribbons, combs, towels and bedding, and by physical contact.

Infestations of lice are easily treated today, especially with this product. This package, with its special comb, is a complete product designed to kill and remove lice and their eggs rapidly and effectively. (Its principal active ingredient is derived from a species of the chrysanthemum flower.)

TREATMENT OF LICE INFESTATION

STEP 1: Shampoo with Liquid

- Shake Well. Apply undiluted product to dry hair and scalp or any other infested areas until entirely wet. Do not use on eyelids or eyebrows.
- Allow product to remain on the area for 10 minutes but no longer.
- Rinse with warm water and your regular shampoo or soap.

This treatment should kill virtually 100% of the adult lice but not all their eggs. All eggs must be removed to prevent

reinfestation. To ensure that all eggs are removed, it is very important to properly use the special comb provided.

STEP 2: Comb Hair Properly

- While hair is still wet, comb with your regular comb to remove snarls and tangles. Part hair into four sections.
- Starting at the top of the head from any one of the four parted sections, lift a one-inch wide tuft of hair up and away from the scalp.
- Take the special nit removal comb in your other hand and place the teeth of the comb as close to the scalp as possible. Comb slowly away from the scalp to the end of the hair, being careful to comb the entire length of each one-inch section of hair.
- Clip or pin back nit-free strands. Continue combing and pinning until the entire section is free of nits and lice. Wipe nits from the comb frequently with a tissue.
- Comb remaining hair sections as directed above. Always comb away from the top of the head to prevent reinfestation of combed hair. This may take some time if hair is long and thick. If hair dries during combing, wet with water.
- When entire head is combed, rinse thoroughly with water.
- After hair is dry, recheck entire head for stray nits and comb out any remaining nits.

STEP 3: Prevent Reinfestation

- Soak comb in either the lice treatment liquid or hot soapy water (130°F) for 10 minutes.
- Repeat treatment with this product in 7-10 days to kill any hatched lice which may have been missed during the initial treatment. Do not exceed two consecutive applications within 24 hours.

Since there is no immunity from lice, personal cleanliness and the avoidance of infested persons and their bedding and clothes will aid in preventing infestation. These additional steps are important in order to minimize the chance of possible reinfestation.

- Inspect all family members daily for at least two weeks, and if they become infested, treat with this product as directed.
- Wash all personal clothing, nightwear and bedding of any infested person in hot water, at least 130°F, or by dry cleaning.
- Thoroughly soak all personal articles such as combs, brushes etc., in lice treatment liquid or hot soap/water (at least 130°F) for ten minutes.
- Tell children not to use any borrowed combs or brushes, nor to wear anyone else's clothes.
- For carpets, bedding, furniture, and other objects which become infested with lice, but which are not easily laundered or dry cleaned, ask your pharmacist for Lice Control Spray (not for use on humans or animals).

This product should not be swallowed or inhaled. It should be kept out of the eyes. If it should come in contact with the eyes, immediately flush eyes with water. This product may be used on children, but only with adult supervision.

Manufactured by
BARRE-NATIONAL INC.
Baltimore, MD 21207 USA

FORM NO. 1023

Rev 10/91

BEST AVAILABLE COPY