HOSPITAL SPRAY DISINFECTANT



DEC 1:6 2005

Under the Federal Insecticide, Fungicide, and Rodenticide, Act as amended, for the pesticide, registered under EPA Reg. No. 5639

Destroys Odors at Their Source

Fast Acting for Multiple Microorganisms

Kills Bacteria, Viruses[†], and Fungi on Hard Surfaces:

Mycobacterium tuberculosis (TB); HIV-1 (AIDS) Hepatitis: Virus (HBV); Hepatitis C Virus (HCV); Trichophyton mentagrophytes (Athlete's Foot Fungus)

In 2 Minutes:

Staphylococcus aureus; Pseudomonas aeruginosa; Salmonella choleraesuis; Streptococcus pyogenes; Clostridium difficile (vegetative)

EPA Reg. No. 56392-4

	-0	
A	T 11	
Active	Ingredic	ents.
Β.		

o-Phenylphenol								. 00.13%
Ethanol								. 69.10%
Other Ingredients								
TOTAL			٠	Ý.				100.00%

Contains sodium nitrite.

Contains no CFCs or other ozone depleting substances. Federal regulations prohibit CFC propellants in aerosols.

KEEP OUT OF REACH OF CHILDREN

READ COMPLETE PRECAUTIONS AND **DIRECTIONS ON REAR PANELS**


NET WT. 14 oz.

CITREX® HOSPITAL SPRAY DISINFECTANT

KILLS GERMS: Spray as directed to kill germs on environmental surfaces such as stainless steel, chrome, plastic, porcelain and mirrors. FOR USE IN MEDICAL FACILI-TIES: Hospitals, Clinics, Nursing Homes, Physician Offices and Dental Clinics, SUGGESTED USES: Door knobs, telephones, supply carts, empty waste cans and laundry hampers, bathroom basins, urinals and mirrors, bed frames and springs. wheelchairs and walkers, DESTROYS ODORS: This effective product can also be used to eliminate odors at their source by killing germs that cause odor. Non-germicidal sprays cannot do this. PREVENTS MOLD AND MILDEW.

THIS PRODUCT IS NOT TO BE USED AS A TERMI-NAL STERILANT/HIGH LEVEL DISINFECTANT ON ANY SURFACE OR INSTRUMENT THAT (1) IS INTRODUCED DIRECTLY INTO THE HUMAN BODY, EITHER INTO OR IN CONTACT WITH THE **BLOODSTREAM OR NORMALLY STERILE AREAS OF** THE BODY. OR (2) CONTACTS INTACT MUCOUS MEMBRANES BUT WHICH DOES NOT ORDINARILY PENETRATE THE BLOOD BARRIER OR OTHERWISE ENTER NORMALLY STERILE AREAS OF THE BODY. THIS PRODUCT MAY BE USED TO PRECLEAN OR DECONTAMINATE CRITICAL OR SEMI-CRITICAL MEDICAL DEVICES PRIOR TO STERILIZATION OR HIGH LEVEL DISINFECTION.

STORAGE AND DISPOSAL: Do not contaminate water, food, or feed by storage and disposal. Storage: Do not use or store near heat or open flame. Store out of reach of children. Disposal: This container may be recycled in the few but growing number of communities where (steel) aerosol can



PRECAUTIONARY STATEMENTS: Hazards to Humans and Domestic Animals. CAUTION: Do Not Spray in Eyes. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear safety glasses. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. FIRST AID: IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Have the product with you when calling a Poison Control Center or doctor or going for treatment. In case of emergency, call a doctor or National Poison Control Center at 800-222-1222. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting. Before using on painted or plastic surfaces, spot test on an inconspicuous area. Not recommended for use on acrylic plastics or vinyl.



DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. TO CLEAN: Shake well. Spray lightly and immediately wipe dry with cloth or paper towel. Effective as a light duty cleaning agent to remove fingerprints and hand smudges. Use on nonporous surfaces such as stainless steel, chrome, mirrors, TO DISINFECT: Shake well. Hold can upright 6-8" from surface. Spray non-porous surface until thoroughly wet and allow to remain wet for appropriate time, depending on the organism (See front panel). Wipe dry or allow to air dry. Gross soil must be removed prior to disinfecting. For Mycobacterium tuberculosis (TB), spray this product on a pre-cleaned surface until thoroughly wet. Allow to remain wet for 1 minute at room temperature for TB. Kills HIV, HBV, and HCV on precleaned environmental surfaces/objects previously soiled with blood/body fluids in healthcare settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood/body fluids, and in which the surfaces/objects likely to be soiled with blood/body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus, Type 1 (HTV-1)(associated with AIDS), Hepatitis B Virus (HBV), and Hepatitis C Virus (HCV). Special Instructions For Cleaning and Decontamination Against HTV-1 (AIDS Virus), HBV, and HCV on Surfaces/Objects Soiled with Blood/Body Fluids: Personal Protection: When handling items soiled with blood/body fluids, appropriate personal protective equipment such as disposable plastic gloves, gowns, masks or eye coverings should be considered. Cleaning Procedure: Blood/body fluids must be thoroughly cleaned from surfaces and objects prior to disinfecting. Contact Time: This product kills HIV-1 (AIDS Virus), HBV, and HCV in 1 minute at room temperature (68-77°F, 20-25°C). Disposal of Infectious Materials: Blood/body fluids and cleaning materials should be autoclaved and disposed of according to federal, state and local regulations for infectious waste disposal. TO DESTROY ODORS: Shake well. Hold upright. Spray 1-2 seconds. May be used to eliminate odors retained in fabrics, draperies or upholstered furniture. TO PREVENT MOLD AND MILDEW: Shake well. Clean surface prior to application. Spray surface until thoroughly wet and repeat weekly to prevent regrowth. Provides residual bacteriostatic action in the presence of moisture.

©2005 Caltech Industries, Inc. Midland, MI 48642 1-800-234-7700 ®Trademark of Caltech Industries, Inc.

REORDER NO. 49200 DOT 2P

