Practical Treatment

Atropine is the emergency antidote for **Vapona** Insecticide poisoning

If swallowed Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger, or, if available, by administering syrup of ipecac Do not induce vomiting or give anything by mouth to an unconscious person Call a physician or Poison Control Center

If on skin Immediately remove the patient from the vicinity of the insecticide, remove all contaminated clothing, and wash skin with soap and running water

If in eyes Wash immediately with running water

If Warning Symptoms Appear Keep patient prone and quiet Start artificial respiration immediately if patient is not breathing Transport the patient immediately to the nearest hospital (See **Warning Symptoms** below)

Note to Physician

Warning Symptoms Symptoms include weakness headache, tightness in chest blurred vision, non reactive pinpoint pupils salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps

Treatment Atropine is the specific therapeutic antagonist of choice against parasympathetic nervous stimulation If there are signs of parasympathetic stimulation, atropine sulfate should be injected at 10 minute intervals, in doses of 1 to 2 milligrams until complete atropinization has occurred

Morphine is contraindicated

Pralidoxime chloride (2 PAM chloride) may also be used as an effective antidote in addition to and while maintaining full atropinization In adults, an initial dose of 1 gram of 2 PAM should be injected, preferably as an infusion in 250 cc of saline over a 15 to 30 minute period If this is not practical, 2 PAM may be administered slowly by intravenous injection as a 5 percent solution in water over not less than two minutes After about an hour, a second dose of 1 gram of 2 PAM will be indicated if muscle weakness has not been relieved For infants and children the dose of 2 PAM is 0 25 grams

Clear chest by postural drainage

Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood test

Warranty and Limitation of Damages

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and Buyer assumes the risk of any use contrary to such directions **Seller makes no other express or implied warranty, including any other express or implied warranty of Fitness or of Merchantability and no agent of Seller is authorized to do so except in writing and with specific reference to this warranty** In no event shall Seller's liability for any breach of warranty exceed the purchase price of the material as to which claim is made

Net Contents: 1 Gallons (3.8 Liters)

FERMENTA ANIMAL HEALTH

Vapona Insecticide Emulsifiable

For Fly and Mosquito Control on Dairy and Beef Cattle & in and Around Animal Buildings.

Active Ingredients:	By Weight
Dichlorvos 2,2-Dichlorovinyl dimethyl phosphate	21.82*
Related Compounds	1.65*
Inert Ingredients**	76.53**
Total	100.00

*Equivalent to 23.41 % Vapona Insecticide

**Inert Ingredients include petroleum hydrocarbons

Keep Out of Reach of Children

DANGER! POISON

See side panel for Precautionary Statements

Must be diluted prior to use.

Fermenta Animal Health Company
Kansas City, MO 64120 1350
EPA Reg No 564914

06-23-1

Not Reviewed for compliance with