

300  
17  
420983

AUG 25 1992

Mr. Robert Shoup  
Fermenta Animal Health Co.  
10150 N. Executive Hills Blvd.  
Kansas City, MO 64153-2314

Dear Mr. Shoup:

Subject: Label Amendment  
RAVAP E.C. Livestock, Poultry & Premise Insecticide  
EPA Registration No. 56493-42  
Your Submission Dated May 4, 1992

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable, provided that you make the labeling change listed below before you release the product for shipment bearing the amended labeling:

Under Directions for Use for Lactating Dairy Cattle add "Care should be taken that the spray does not come in direct contact with the lactating dairy cow's teats unless they are washed with an approved cleansing solution and dried before milking. Apply the spray 20 minutes before beginning to milk, or after milking has been completed."

Submit five copies of your final printed labeling before you release the product for shipment.

A stamped copy of the labeling is enclosed for your records.

Sincerely,

George T. LaRocca  
Product Manager 13  
Insecticide-Rodenticide Branch  
Registration Division (H7505C)

CONCURRENCES

SYMBOL							
SURNAME	LaRocca						
DATE	8-17-92						

# Ravap E.C.

## Livestock, Poultry & Premise Insecticide Spray

ACCEPTED  
with COMMENTS  
to EPA Letter Dated

AUG 15 1973

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

56493-42

TAKE TIME  
OBSERVE LABEL  
DIRECTIONS



### Recommended for Agricultural Use

Contains Rebon\* and Vapona\* Insecticides

Active Ingredients	By Weight
Tetrachloro-phos 2-Chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate	23.0%*
Dichlorod 2,2-Dichlorovinyl dimethyl phosphate	5.2%**
Related Compounds	0.4%***
Inert Ingredients	71.3%***
Total:	100.0%

\*Rebon Insecticide \*\*Equivalent to 5.7% W Vapona Insecticide  
\*\*\*Inert ingredients include Xylene

This product contains the toxic inert ingredient phenol

**Keep Out of Reach of Children**

**DANGER:** See other panels for additional  
precautionary statements

Store Above 32°F.

Do not store above 120°F for  
an extended period of time.

Net Contents: 1 Gallon (3.8 Liters)

Fermenta Animal Health Company  
Kansas City, MO 64153

EPA Reg. No. 56493-42  
EPA Est. No. 602 MD-1



### Statement of Practical Treatment

If swallowed- Call a physician or Poison Control Center immediately. If possible, vomiting should be induced under medical supervision. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled- Remove victim to fresh air. Apply artificial respiration if necessary.

If on skin- Remove contaminated clothing and wash skin immediately with soap and water.

If in eyes- Flush eyes with plenty of water. Call a physician immediately.

Note to physician- If signs of cholinesterase inhibition are observed, atropine by injection is antidotal and may be administered in conjunction with 2-PAM (Pralidoxime Chloride). Product may cause aspiration pneumonia. Gastric lavage may be indicated if product was taken internally.

### Precautionary Statements

#### Hazards to Humans and Domestic Animals

##### Danger

Corrosive, causes eye damage, causes skin irritation. Do not get in eyes (including animals), on skin or on clothing. Wear goggles or face shield and rubber gloves; long-sleeved shirt and pants; shoes and socks when handling.

Harmful if inhaled. Avoid breathing spray mist.

Harmful or fatal if swallowed. Wash thoroughly after using and before smoking or eating. Avoid contamination of food.

#### Environmental Hazards

This product is toxic to fish, birds, and other wildlife. Drift and runoff may be hazardous to aquatic organisms in adjacent areas. Do not apply directly to water or wetlands. Do not contaminate water when disposing of equipment washwater.

#### Physical and Chemical Hazards

Do not use, pour, spill or store near heat or open flame. Do not use with thermal foggers or heat-generating devices.

DSL 841 B 9110

3/3

### Directions for Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. For Agricultural Use Only.

This product is suitable for use in conventional power or low pressure knapsack sprayers and in livestock backrubbers. Follow the Use Directions for the proper dilution needed for specific insect control. Must be diluted in water except for backrubber use.

#### Livestock Use Directions

Use	Insect	Dilution Ratio in Water	Remarks
Livestock Spray (Beef Cattle)	Horn Flies, Lice, including Tail Lice, and Face Flies (aid in control)	1 gal. in 75 gal. or 5 oz. in 3 gal.	Dilute in water as indicated and spray directly on the animal only to the point of runoff. Use between 1/2 to 1 gallon of diluted spray solution per animal depending on size and hair coat. Do not treat more often than every 10 days. Do not apply to calves under six months of age. Brahman and Brahman-cross cattle should not be treated as they may show hypersensitivity to organophosphate pesticides. Do not apply in combination with other dermal organophosphate pesticides (e.g., Ivermectin). There is no withholding period from last application to slaughter.
	Horn Flies, Lone Star Ticks	1 gal. in 200 gal. or 2 oz. in 3 gal.	Apply as above. For severe tick infestations, dilution may be increased to 1 gallon in 50 gallon water.
Livestock Spray (Lactating Dairy Cattle)	Horn Flies, Lice, Lone Star Ticks, and aids in control of Face Flies	1 gal. in 200 gal. or 2 oz. in 3 gal.	Dilute in water as indicated. Direct spray to cover thoroughly with up to 1/2 gallon of the dilution per animal. Repeat as necessary. Do not apply to calves under six months of age. No milk discard is required.

Beck Page 1

#### Backrubber Use Directions

Use	Insect	Dilution Ratio in Oil	Remarks
Backrubber or Face Rubber (Beef Cattle and Dairy Cattle)	Horn Flies, Face Flies (aid in control)	1 gal. in 25 gal. or 5 oz. in 1 gal.	Mix with #2 diesel oil or any approved backrubber base oil. Pour diluted solution into oil reservoir of mechanical rubbing devices or pour one gallon per 20 linear feet on burlap or rope backrubbers. Keep backrubber or face rubber charged.

#### Poultry Use Directions

Use	Insect	Dilution Ratio in Water	Remarks
Caged Chickens	Lice & Mites	1 gal. in 50 gal. or 2 oz. in 2 gal.	Apply 1 gal. of dilution/100 birds under high pressure (no less than 100-125 psi) to the vent and fluff areas from below. Repeat when necessary; however, not more often than every 14 days. For individual bird treatment, apply 1 ounce of dilution per bird.
Chickens on Litter	Lice & Mites	1 gal. in 50 gal. or 5 oz. in 2 gal.	Apply 1-2 gal. of dilution/1000 sq. ft. evenly with penetration of litter surface. Also apply thoroughly to walls, roosts, cracks and crevices. Spray birds as above. Treat roosters carefully to avoid reinfestation of breeding flocks.
Roost Paint	Lice & Mites	1 gal. in 25 gal. or 5 oz. in 1 gal.	Apply 1 pt. of dilution/100 ft. of roost area with brush or spray.
Poultry Buildings	Fowl Tick (Blue Bug)	1 gal. in 25 gal. or 5 oz. in 1 gal.	Apply 1 gal. of dilution/100-150 sq. ft. to thoroughly cover walls, ceilings, floors, cracks and crevices using high pressure spray.

#### Premise Use Directions for Poultry and Livestock Facilities

Use	Insect	Dilution Ratio in Water	Remarks
Residual Surface Spray	Flies, Litter Beetles	1 gal. in 25 gal. or 5 oz. in 1 gal.	Apply 1 gallon of dilution/500-1000 sq. ft. Thoroughly cover walls, ceilings or other areas where pests rest or congregate in dairy barns, poultry houses, swine buildings, livestock sheds and other farm buildings. Extreme infestations may necessitate increasing the diluted spray to one gallon per 12 1/2 gal. or 10 oz. per 1 gal. of water.
Larvicide	Maggots (fly larvae)	1 gal. in 25 gal. or 5 oz. in 1 gal.	Apply 1 gal. of dilution/100 sq. ft. of droppings as a coarse spray. Repeat at 7-10 day intervals until droppings begin to come up, then treat only "hot spots" (small areas found to have large number of maggots). Do not spray manure where runoff to soil or water can occur. Do not spray animals directly with this concentration.

### Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage-** Store in original container only. Store at temperatures above 32° F. Do not store above 120° F for an extended period of time. Do not store diluted product.

**Pesticide Disposal-** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your state pesticide or environmental control agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**Container Disposal-** Triple rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or by other approved State and local procedures.

Beck Page 1

### Warranty and Limitation of Damages

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and Buyer assumes the risk of any use contrary to such directions. Seller makes no other express or implied warranty, including any other express or implied warranty of Fitness or of Merchantability, and no agent of Seller is authorized to do so except in writing and with specific reference to this warranty. In no event shall Seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

Fermenta Animal Health Company • Kansas City, MO 64153-2314

**BEST AVAILABLE COPY**