

TH 10 56493-77

10

56493-79

32 Weeks

932770

EctoGard Fogger contains a combination of three insecticides that attack the flea life cycle from three directions & controls ticks, cockroaches, crickets, silverfish, earwigs, woodrats, confused flour beetles, ants, spiders & flies. 1) The first action is provided by the effective, quick killing activity of pyrethrins. Pyrethrins will kill insects they come in direct contact with but will not provide residual activity. 2) Permethrin is the second active ingredient. It takes over as the pyrethrins lose effectiveness by providing residual control of fleas in carpet. 3) The third & most important ingredient is the new insect growth regulator, fenoxycarb. It inhibits development of the immature stages of the flea by 34 weeks, preventing them from reaching the biting adult stage.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION: Harmful if inhaled or absorbed through skin. Causes eye injury. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap & water after handling. Remove contaminated clothing & wash before reuse. Do not contaminate food or feedstuffs. Do not use in edible product areas of food processing plants, restaurants, or other areas where food is commercially prepared, processed or stored. Remove pets, birds and/or cover fish aquariums before use.

STATEMENT OF PRACTICAL TREATMENT

IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated.
IF ON SKIN: Remove contaminated clothing & wash affected areas with soap & water.
IF IN EYES: Flush eyes with plenty of water. Get medical attention, if irritation persists.

PHYSICAL AND CHEMICAL HAZARDS

Flammable: Contents under pressure. Keep away from heat, sparks & open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

EctoGard™

with Fenoxycarb™ Insect Growth Regulator

Fogger

KILLS BOTH ADULT & PRE-ADULT FLEAS

PLUS BREAKS FLEA LIFE CYCLE BY PREVENTING DEVELOPMENT INTO ADULTS

One Treatment Kills Fleas For 34 Weeks

ACTIVE INGREDIENTS

Fenoxycarb (ethyl 2-(4 phenoxycarbonyl) ethylcarbamate)	100%
Pyrethrins	0.05%
Permethrin (3 phenoxycarbonyl methyl (±) cis trans 3 (2,2 dichloroethoxy) 2,2 dimethylcyclopropane carboxylate)	0.50%
Related Compounds	0.04%
Picroyl Butoxide, technical	0.10%
N acetyl bicycloheptene dicarboximide	0.10%
INERT INGREDIENTS	97.91%
	100.00%

KEEP OUT OF REACH OF CHILDREN

See side panel for additional precautionary statements

TechAmerica™

NET WT. 8 OZ. (170.1 g)

Formosa Animal Health Company
PO Box 991158
Kansas City, MO 64148-1158 A

EPA Reg. No. 56493-79
EPA Est. No. 7554-1158
EPA 1137

EctoGard™ and Fenoxycarb™ are trademarks of Formosa Animal Health Co.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

For the most effective results, the entire dwelling should be treated. Use one canister for each 8,000 cu. ft. (or 750 sq. ft.) Use additional units for remote rooms or where free flow of the mist is not assured.

PREPARATION: Cover exposed foods, dishes & food handling equipment. Open cabinets & closet doors to areas to be treated. Turn off fans & air conditioners. Extinguish all open flames except pilot light. Remove pets & cover aquariums or fish bowls with paper, or remove from area. Be sure to leave pet bedding as this is a primary hiding place for fleas & must be treated to get best results. No need to discard pet bedding after treatment. Close outside doors & windows.

TO START FOGGER: Place fogger on table, chair or stand, in center of area, at least six feet from pilot light. Place several sheets of newspaper covering the area directly under the can. Hold can at arm's length, with top of can pointing away from face & eyes. Push down actuator tab until it clicks—this will start fogging action. Set canister in an upright position on table, etc., and leave area of building at once.

DO NOT RE-ENTER BUILDING FOR TWO HOURS. AFTER TWO HOURS, open all outside doors & windows, turn on air conditioners & fans & let treated area air out for 30 minutes before reoccupying. If additional units are used for remote rooms or where free flow of mist is not assured, increase airing out time accordingly. Thoroughly clean all covered food processing areas before using.

STORAGE AND DISPOSAL

Store in a cool area away from children. Fogging cap, wrap in newspaper & discard container in trash. Do not incinerate or puncture.

BEST AVAILABLE COPY