

P11 10 56493-84

1 of 1

FRONT PANEL

ELIMINATOR™ FOGGER  
with Tenocide™ Insect Growth Regulator

KILLS BOTH ADULT & PRE-ADULT FLEAS  
plus breaks flea life cycle by preventing  
development into adults  
plus kills ticks, cockroaches and flies

One Treatment Kills Fleas For ~~21 Weeks~~ 32 weeks

**ACCEPTED**  
**17 APR 1992**  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 56493-84

ACTIVE INGREDIENTS:

Fenoxycarb [ethyl[2-(4-phenoxyphenoxy)ethyl]carbamate]	1.00%
Permethrin [(3-phenoxyphenyl) methyl(+) cis-trans-3-(2,2-dichloroethenyl) 2,2-dimethylcyclopropane- carboxylate]	0.50%
Related Compounds	0.04%
Piperonyl Butoxide, technical*	0.10%
N-octyl bicycloheptene dicarboximide+	0.40%

INERT INGREDIENTS

97.96%  
100.00%

\*Equivalent to 0.08% (butylcarbityl) (6-propylpiperonyl)  
ether and 0.02% related compounds.

+MGK 264, Insecticide Synergist

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

See back panel for additional precautionary statements

Net Wt. 6 oz. (170.1 g)

Distributed by:  
Fermenta Animal Health Company  
P. O. Box 901350  
Kansas City, MO 64190-1350 U.S.A.

EPA Reg. No. 56493-84

EPA Est. No. 7056-TX-1

Eliminator and Tenocide are trademarks of Fermenta Animal Health Company.

## BACK PANEL

Eliminator Fogger contains a unique combination of ingredients that attack the flea life cycle plus controls ticks, cockroaches, crickets, silverfish, earwigs, sowbugs, confused flour beetles, ants, spiders and flies. 1) Permethrin provides insecticidal activity by knocking down and controlling fleas in carpet. 2) Tenocide, a new insect growth regulator, is the most important active ingredient in the Eliminator Fogger. Tenocide inhibits development of the immature stages of the flea for at least ~~21 weeks~~, preventing them from reaching the biting adult stage.

7  
32 WEEKS

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if inhaled or absorbed through skin. Causes eye injury. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap & water after handling. Remove contaminated clothing & wash before reuse. Do not contaminate food or foodstuffs. Do not use in edible product areas of food processing plants, restaurants, or other areas where food is commercially prepared, processed or stored. Remove pets, birds and/or cover fish aquariums before use.

#### STATEMENT OF PRACTICAL TREATMENT

IF INHALED remove victim to fresh air. Apply artificial respiration if indicated.

IF ON SKIN remove contaminated clothing & wash affected areas with soap & water.

IF IN EYES flush eyes with plenty of water. Get medical attention, if irritation persists.

#### PHYSICAL AND CHEMICAL HAZARDS

Flammable. Contents under pressure. Keep away from heat, sparks & open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

#### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

For the most effective results, the entire dwelling should be treated. Use one canister for each 6,000 cu. ft. (or 750 sq. ft.). Use additional units for remote rooms or where free flow of the mist is not assured.

**PREPARATION:** Cover exposed foods, dishes & food handling equipment. Open cabinets & closet doors to areas to be treated. Turn off fans & air conditioners. Extinguish all open flames except pilot light. Remove pets & cover aquariums or fish bowls with paper, or remove from area. Be sure to leave pet bedding as this is a primary hiding place for fleas & must be treated to get best results. No need to discard pet bedding after treatment. Close outside doors & windows.

**TO START FOGGER:** Place fogger on table, chair or stand in center of area, at least six feet from pilot light. Place several thicknesses of newspaper covering the area directly under the can. Hold can at arm's length, with top of can pointing away from face & eyes. Push down actuator tab until it locks--this will start fogging action. Set canister in an upright position on table, etc. and leave area or building at once.

**DO NOT RE-ENTER BUILDING FOR TWO HOURS.** AFTER TWO HOURS, open all outside doors & windows, turn on air-conditioners & fans & let treated area air out for 30 minutes before reoccupying. If additional units are used for remote rooms or where free flow of mist is not assured, increase airing-out time accordingly. Thoroughly clean all covered food processing areas before using.

#### STORAGE AND DISPOSAL

Store in a cool area away from children. Replace cap, wrap in newspaper & discard container in trash. Do not incinerate or puncture.