

5. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.



William H. Miller
Product Manager (16)
Insecticide-Rodenticide Branch
Registration Division (TS-767C)

Enclosures

TRAP-N-A-SAK™
WITH DIPHACINONE
A Pelletized Rodenticide

READY MIXED, EASY TO USE
USE INDOORS AND OUTDOORS



ACCEPTED
with COMMENTS
in EPA Letter Dated:

NOV 1960

Under the Fungicide, Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

56637-1

ACTIVE INGREDIENT

Diphacinone [2-(Diphenylacetyl)-
1,3-indandione].....

0.005%

INERT INGREDIENTS.....

99.995%

TOTAL.....

100.000%

Patent Pending

EPA Reg. No.

EPA Est. No. 2393-WI-1

KEEP OUT OF REACH OF CHILDREN (12 pt.)

CAUTION (18 pt.)

See Other Panel For Additional Precautions

Manufactured For

BLOOMFIELD FEED MILL
BOX 487
BLOOMFIELD, NE 68718

NET WEIGHT

1 LB

Left Panel

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION

Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.

NOTE TO PHYSICIAN - If ingested, administer Vitamin K1 intramuscularly or orally, as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARD
Keep Out of Lakes, Streams or Ponds.


DIRECTIONS FOR USE
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DISPOSAL

Do not reuse empty container.
Wrap container and put in trash.

USE RESTRICTIONS - For control of Norway rats, Roof rats and House mice in and around homes, industrial and agricultural buildings, and similar man-made structures.

Treated baits must be placed in tamper-proof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.



Right Panel

SELECTION OF TREATMENT AREAS - Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Protect bait from rain or snow. Remove as much food as possible.

APPLICATION DIRECTIONS

RATS: Apply 4 to 16 ounces of bait per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

MICE: Apply 1/4 to 1/2 ounce (1-2 level tablespoons) of bait at 8 to 12 foot intervals. Larger placements (up to 2 ounces) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

RATS & MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

NOTICE OF WARRANTY

SELLER AND MANUFACTURER MAKE NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, EXPRESS OR IMPLIED, CONCERNING THIS PRODUCT OR ITS USE, WHICH EXTEND BEYOND THE STATEMENTS ON THIS LABEL.

