

10/24/89

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

WARNING

May be fatal if swallowed. May be absorbed through skin. Do not breathe spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling and before eating or smoking. Avoid contamination of food and feedstuffs. Do not use on household pets or humans. Do not permit children or pets to go onto sprayed areas until the spray has completely dried. Keep out of reach of domestic animals. Do not contaminate ornamental landscapes. DO NOT USE IN THE HOME.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. PROTECT BEES: This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are actively visiting the crop, cover crop, or weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

LAWNS: Overseed, Merion, Fescue, St. Augustine, Bermuda, Centennial, Bermuda. To control Chinch Bugs*, Sod Webworms, Bermuda Mites, Ants, Ticks, Earwigs, Lawn Billbugs, apply 2 lbs. (1 fl. oz.) in 3 gals. of water per 125 sq. ft. or 8 fl. oz. in 24 gals. of water per 1,000 sq. ft. Thoroughly wet down grass a few hours before applying. For Lawn Billbugs, water lawn thoroughly after treatment. *In the North, if Chinch Bugs are the only insect problem, a lower rate of application can be used, 2 lbs. in 3 gals. of water per 250 sq. ft. To control Rhodes-Grass Scale Crawlers, apply 2 lbs. (1 fl. oz.) in 3 gals. of water per 100 sq. ft. or 10 fl. oz. in 30 gals. of water per 1,000 sq. ft. Apply when Scale Crawlers first emerge.

FIRE ANTS: To aid in the control of fire ants in lawns and other recreational areas, apply 1 fl. oz. (2 lbs.) in a minimum of 1 gal. of water as a drench. Apply over and 6 inches around each mound. Apply gently to avoid disturbing ants. Reapply as necessary. Larger quantities of solution can be prepared for multiple mound treatments. Use equipment capable of delivering the diluted product as a gentle rain at the rate of 1 gal. per mound. High pressure sprays may disturb the ants and cause migration, reducing product effectiveness. For best results, apply in cool weather, 65-90°F, or in early morning or late evening hours. Treat new mounds as they appear.

ROSES AND FLOWERS: Rose, Azalea, Chrysanthemums, Gladiolus, Carnations. To control Aphids, Mites, Leafhoppers, Thrips apply 2 lbs. (1 fl. oz.) per 3 gals. of water. Apply as a thorough foliar spray. Spray underside of leaves. Wet plants to the dripping point.

TREES AND SHRUBS: Oak-To control Aphids and Scale Crawlers, Maple-To control Aphids, Locust-To control Aphids, Holly and Boxwood-To control Mites, Scale Crawlers and Leaf Miners, Pine-To control Aphids and Scale Crawlers, Spruce-To control Aphids, Scale Crawlers, and Bagworms, Willow-To control Aphids, Scale Crawlers, and Leaf Miners. Apply 2 lbs. (1 fl. oz.) in 3 gals. of water. Apply as a thorough foliar spray. Wet leaves and branches to the dripping point.

BEST AVAILABLE COPY

DIAZINON 2-EC

For Use On Lawns, Flowers,
Ornamentals, Vegetables,
Fruits, Outdoor Areas.

THIS PRODUCT MUST NOT BE USED
ON GOLF COURSES AND SOD FARMS

ACTIVE INGREDIENTS:

O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl)
phosphorothioate 75.0%

INERT INGREDIENTS: 25.0%

Total 100.0%

* Diazinon is a registered trademark of CIBA-GEIGY CORPORATION.

This product contains 2 lbs. of Diazinon® per gallon.

KEEP OUT OF REACH OF CHILDREN

WARNING

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. If possible, vomiting should be induced under medical supervision. Drink one or two glasses of water and induce vomiting by touching the back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated.

IF ON SKIN: Remove contaminated clothing and wash affected areas with soap and water.

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

Manufactured for Security Products
Company of Delaware, Inc.

Atlanta, GA

EPA Reg. No. 56644-14

EPA Est. No. 769-GA-1

Not Reviewed. Registrant claims to be in accordance with
Reregistration of Pesticide Product Guidance for Phase 2
Response, Page 2.12, Active to Inert Change in Status.

FRUITS: Apples-To control Leafhoppers, Mites, Aphids, Leaf Rollers and Scale Crawlers. Pear-To control Mites, Aphids, Leaf Rollers, and Scale Crawlers. Oranges-To control Mites, Aphids, and Scale Crawlers. Plums-To control Mites, Aphids, Leaf Rollers and Scale Crawlers. Grapes-To control Mites, Aphids, Leaf Rollers and Scale Crawlers. Cherries-To control Mites, Aphids, and Scale Crawlers. Strawberries-To control Mites, Aphids and Leaf Rollers. Apply 2 lbs. (1 fl. oz.) in 3 gals. of water. Apply as a full cover spray. Start spraying after blossoming and repeat at 10 to 14 day intervals if needed. Do not spray fruit within 21 days of picking.

VEGETABLES

SECURITY DIAZINON 2-EC may be applied to the following vegetables. Follow the time interval (in days) between last application and harvest indicated by the number in () following each vegetable. A (0) indicates product may be applied through day of harvest.

Cabbage (7)	Peas (0)	Radish (10)
Carrots (10)	Pole Beans (7)	Red Beets (14)
Corn (0)	Potatoes (35)	Snap Beans (7)
Cucumber (7)	Lettuce (10)	Tomatoes (1)
Cauliflower (3)	Peppers (5)	Turnips (10)

FOLIAR APPLICATION: To control Aphids, Leaf Miner, Bean Beetles, Cucumber Beetle, Diamond-back Moth Larvae, and Imported Cabbageworm on the vegetables listed above, apply 2 lbs. (1 fl. oz.) in 3 gals. of water per 1,000 sq. ft. as a foliar spray. Generally applications at 7-10 day intervals may be required.

PREPLANT SOIL APPLICATION: To control Cutworms (surface, subterranean) on Cabbage, Carrots, Corn, Lettuce, Peas, Pole Beans, Radishes, Red Beets, Snap Beans, Tomatoes and Turnips, apply 6 fl. oz. per 1,000 square feet in sufficient water to achieve thorough coverage. Apply broadcast just before planting. Immediately work into soil 2-3 inches for surface cutworms and 3-6 inches for subterranean cutworms. To control Mole Crickets on Corn, Carrots, Peas, Pole Beans, Red Beets and Snap Beans, apply 1 fl. oz. per 1,000 square feet in sufficient water to achieve thorough coverage. Apply broadcast just before planting. Immediately work into soil 1-2 inches. To control Root Maggots on Cabbage, apply 4 fl. oz. per 1,000 square feet in sufficient water to achieve thorough coverage. Apply broadcast just before planting. Immediately work into soil 3-4 inches before seeding, either in seed bed or field; or before transplanting. To control Wireworms on Cabbage, Corn, Carrots, Lettuce, Peas, Snap Beans and Tomatoes, apply 6 fl. oz. per 1,000 sq. ft. in sufficient water to achieve thorough coverage. Apply broadcast just before planting. Immediately work into soil 4-8 inches.

OUTSIDE NUISANCE PESTS

Flies, Ants, Cockroaches, Bedbug Bugs, Clover Mites, Spiders, Ticks, Chiggers, Earwigs. Apply 8 fl. oz. per 3 gals. of water. Thoroughly spray window frames, doors, sills, screens, porch and patio walls and ceiling, foundations, walks, garbage cans and into cracks and other places where insects may hide. Do not treat plants with this dosage rate.

DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Do not reuse container. Wrap container and put in trash collection.

NOTICE: Seller makes no implied warranty of merchantability nor any other warranties which extend beyond the description on this label.