



# 5% Sevin® Garden Dust

### ACTIVE INGREDIENT:

Carbaryl (1-naphthyl N-methylcarbamate) 5.0%

INERT INGREDIENTS: 95.0%

EPA REG. NO. 769-418 EPA EST. 769-GA-1 TOTAL 100.0%

\* Trademark of Union Carbide Corp. for 1-naphthyl N-methylcarbamate

## KEEP OUT OF REACH OF CHILDREN CAUTION

(See additional precautionary statements on side panel)

### STATEMENT OF PRACTICAL TREATMENT

**IF SWALLOWED:** Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention. **IF INHALED:** Remove victim to fresh air. **IF ON SKIN:** Wash with soap and water. Remove contaminated clothing. **IF IN EYES:** Flush eyes with water.

**NOTE TO PHYSICIAN:** Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal. Do not use 2-PAM, opiates or cholinesterase-inhibiting drugs.

### PRECAUTIONARY STATEMENTS

**CAUTION:** May be harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling.

### ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of waste. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are present in the treatment area.

### DIRECTIONS FOR USE

This application of Federal law to use this product in a manner inconsistent with its labeling. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl label.

**CAUTION:** Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity of several days' duration following application.

**PREHARVEST USE LIMITATION:** This product can be used up to a certain number of days before harvest of crops listed except on strawberries allow one day between application and harvest, and on broccoli, Brussels sprouts, cabbage, cauliflower, cowpeas, head lettuce, garden beets (roots), kohlrabi, radishes, rutabaga, and turnips allow 3 days between application and harvest, and on blackberries, raspberries, dewberries, loganberries and raspberries allow 7 days and between application to poultry and game birds (when so treated) and slaughter of birds, allow 7 days on collards, garden beets, kale, leaf lettuce, mustard greens, spinach and turnip tops allow 14 days between application and harvest.

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under 769-418

skeletonizer, apply at the rate of 20 to 40 pounds per acre. Apply just before first brood leaf to permit larvae emerge from rolls and as needed for leafhoppers. To control grape berry moth, Japanese beetle, June beetle and red-banded leafroller, apply at the rate of 40 pounds per acre. **STRAWBERRIES:** To control meadow spittlebug, strawberry leafroller and strawberry weevil, apply at the rate of 20 to 40 pounds per acre.

### ORNAMENTALS

To control scale aphid, bagworm, birch leafminer, blister beetles, boxelder bug, boxwood leafminer, elm leaf aphid, elm leaf beetle, flea beetles, Japanese beetle, lacebugs, leafhoppers, leaf rollers, oak leafminer, orange tortrix, periodical cicada, plant bugs, psyllids, rose slug, rose-worm, scale, spider mite, the crawler stage, tent caterpillars, willow leaf beetle and exiled trips, apply thoroughly to the infested plants at first sign of damage and repeat at 7 day intervals or as necessary for best results. To obtain scale control apply in spring and early summer when new growth is present.

### LAWNS

To control sod webworm, cutworm and sod webworm, apply at the rate of 4 to 6 pounds per 1000 sq. feet. For best results, apply after rain or watering, and repeat at least 2 days after application. Repeat in 2 to 3 weeks if necessary.

### POULTRY INSECT CONTROL

**CHICKENS, TURKEYS, GOOSE, DEER, GAME BIRDS AND PIGEONS:** **DUST-BATH TREATMENT:** Apply 1/2 pound per 40 square feet to control Northern fowl mite, lice, ticks, fleas and bed bugs. Use a spray and dust combination for best results. Avoid contamination of nests, eggs, feed and watering troughs.

**HAND APPLICATION TO BIRDS:** Use one pound to treat 100 birds by means of a hand duster or hand bottle or hand duster to control Northern fowl mite, lice, ticks, fleas. Direct dust to vent and fluff areas. Repeat at weekly intervals. Hand application to birds for chicken mite and lice is also useful. For better treatment for control of these pests, follow the instructions for dust-bath treatment. It is especially important to treat necks and feet.

**DUST-BATH TREATMENT:** Use 1/2 pound per box for each 50 birds to control Northern fowl mite, lice, ticks, fleas and fleas. This is a supplement to other measures for lice, ticks and flea control. On ranges, two natural lawns must be substituted for each dust-bath box.

### PET INSECT CONTROL

**DOGS AND CATS:** Use 1/2 ounce of dust liberally over the entire body of the animal. Rub the dust into the skin. Comb out dead ticks a few days after treatment. Repeat at weekly intervals if needed. Avoid application to the face of the animal. Do not use on puppies and kittens.

**POSSIBLE ADVERSE EFFECTS:** Use dust liberally on the face and ears of dogs and cats. Do not use on the face of the animal. Do not use on the face of the animal. Do not use on the face of the animal.

### STORAGE AND DISPOSAL

Store in original container. Store in a cool, dry place. Do not use if container is damaged. Do not use if container is damaged. Do not use if container is damaged.

NET WEIGHT 4 POUNDS

SECURITY CHEMICAL COMPANY

MANUFACTURERS ATLANTA, GEORGIA

U-123933

boysenberries, dewberries, loganberries and raspberries allow 7 days and between application to poultry and game birds (when so treated) and slaughter of birds, allow 7 days on collards, garden beets, kale, leaf lettuce, mustard greens, spinach and turnip tops allow 14 days between application and harvest.

**APPLICATION:** Begin applications when damage is first observed and repeat on a regular 5 to 7 day schedule as necessary for good insect control.

### VEGETABLES

**BEANS:** To control Mexican bean beetle, leaf beetles, cucumber beetles, leafhoppers, flea beetles, Japanese beetle, and cutworms, apply at the rate of 20 pounds per acre or 1 to 2 ounces to 50 feet of row. Where Mexican bean beetle is the only problem, a rate of 10 pounds per acre is sufficient. To control lygus bug, stink bug and corn earworm borer, apply 40 pounds per acre.

**CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CABBAGE, KCHLRABI:** To control flea beetles and harlequin bugs, apply at the rate of 10 to 20 pounds per acre. To control armyworm, imported cutworm, European imported cabbageworm, diamondback moth larvae, apply 20 to 40 pounds per acre or 1 to 2 ounces to 50 feet of row. For control of cabbage loopers.

**COLLARDS, GARDEN BEETS, KALE, LETTUCE, MUSTARD GREENS, RADISHES, RUTABAGAS, SPINACH, TURNIPS, and turnip tops:** To control harlequin bug and leafhoppers, apply at the rate of 10 to 20 pounds per acre. To control armyworms, imported cutworms, diamondback moth larvae, corn earworm, tarnished plant hoppers and stink bugs, apply 20 to 40 pounds per acre or 1 to 2 ounces to 50 feet of row. For control of corn earworm after lettuce heads begin to form, use 40 pounds per acre in a 5 to 7 day schedule. Will not control cabbage looper.

**CUCUMBERS, SQUASH, CANTALOUPE, PUMPKINS, MELONS:** To control pickleworm, melonworm, cucumber beetle, squash bug, and beetles, apply at the rate of 20 pounds per acre or 1 to 2 ounces to 50 feet of row.

**CORN:** To control corn earworm, European cutworm, Japanese beetle, sap beetles, and Japanese beetles, apply at the rate of 20 to 40 pounds per acre or 2 ounces to 50 feet of row. For control of European cutworm, apply to entire plant and repeat as necessary. For control of stink bug, and ears, apply at 2 to 3 day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending upon severity of infestation.

**CARROTS:** To control flea beetles, leafhoppers, harlequin bug and stink bugs, apply at the rate of 20 to 40 pounds per acre or 1 to 2 ounces to 50 feet of row.

**CORN:** To control corn earworm and stink bugs, apply at the rate of 20 to 40 pounds per acre or 1 to 2 ounces to 50 feet of row.

**TOMATO, EGGPLANT, PEPPER:** To control European cutworm, European corn borer and stink bug, apply at the rate of 20 to 40 pounds per acre.

**POTATO:** To control Colorado potato beetle, apply at the rate of 20 pounds per acre. For control of leafhoppers, diamondback moth larvae, European corn borer and stink bug, apply at the rate of 20 to 40 pounds per acre.

### SMALL FRUIT

**BLACKBERRIES, BOYSENBERRIES, DEWBERRIES, RASPBERRIES:** To control Japanese beetle, European cutworm, raspberry aphid, apply at the rate of 40 pounds per acre.

**BLUEBERRIES:** To control blueberry maggot, European cutworm, European fruitworms and Japanese beetles, apply at the rate of 20 to 40 pounds per acre. Apply 3 weeks before harvest and repeat if necessary.

→ Transferred to 56644-15  
BEST AVAILABLE COPY



# 5% Sevin Garden Dust

**ACTIVE INGREDIENT** 5.0%  
5% Sevin (Cyfluthrin) 5.0%  
**INERT INGREDIENTS** 95.0%  
EPA REG. NO. 274-418 EPA EST. NO. 2411 TOTAL 100.0%  
\*Trademark of Union Carbide Corp. for Cyfluthrin, N-methylcarbamate.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

(See additional precautionary statements on side panel)

### STATEMENT OF PRACTICAL TREATMENT

**IF SWALLOWED** - If swallowed, do not induce vomiting. Give 1-2 glasses of water with 1-2 teaspoons of table salt or give 1-2 glasses of water with 1-2 teaspoons of table salt. **IF INHALED** - If inhaled, do not breathe dust. **IF ON SKIN** - Wash with soap and water. **IF IN EYES** - Flush eyes with water.

### PRECAUTIONARY STATEMENTS

**CAUTION** - May be harmful to fish, aquatic life, and birds. Do not apply directly to water.

### ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic life. Do not apply directly to water or to areas that contaminate water by leaching, runoff, or drift. Do not apply directly to fish, birds, or bees. Do not apply to flowering crops or wild flowers. Do not apply to lawns or drift to blooming crops or wild flowers.

### DIRECTIONS FOR USE

This product is for use on lawns, ornamentals, and garden plants. Apply to foliage and soil. Do not apply to water.

**CAUTION** - Do not apply to lawns, ornamentals, and garden plants. Do not apply to water.

**PREHARVEST USE LIMITATION** - Do not apply to strawberries, blueberries, raspberries, blackberries, and on broccoli, Brussels sprouts, cabbage, cauliflower, kohlrabi, and garden beans. Do not apply to garden beans, peas, and soybeans. Do not apply to garden beans, peas, and soybeans. Do not apply to garden beans, peas, and soybeans.

**ACCEPTED**  
JUN 28 1985  
EPA REG. NO. 274-418

### ORNAMENTALS

### LAWNS

### POULTRY INSECT CONTROL

### PET INSECT CONTROL

### STORAGE AND DISPOSAL

**SECURITY CHEMICAL COMPANY**  
MAF, FORTREIS ATLANTA, GEORGIA

boysenberries, dewberries, loganberries and raspberries allow 7 days, and between application to poultry and game birds or premise treatment and slaughter of birds allow 7 days. On collards, garden beets (tops), kale, leaf lettuce, mustard greens, spinach and turnips (tops) allow 14 days between application and harvest.

**APPLICATION** - Begin applications when insect or damage is first observed and repeat on a regular 5 to 7 day schedule or as necessary to good insect control.

### VEGETABLES

**BEANS** - To control Mexican bean beetle, bean leaf beetle, cucumber beetles, leafhoppers, flea beetles, Japanese beetles and corn earworm, apply at the rate of 20 pounds per acre or 1 ounce per 50 feet of row. Where Mexican bean beetle is the only problem, a rate of 10 pounds per acre is sufficient. To control lygus bug, stink bug and Lima bean pod borer, apply 40 pounds per acre.

**CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, KOHLRABI** - To control flea beetles and harlequin bugs, apply at the rate of 10 to 20 pounds per acre. To control corn earworm, armyworm, imported cabbageworm, diamond back moth larvae, apply 20 to 40 pounds per acre or 1 to 2 ounces to 50 feet of row. Will not control cabbage loopers.

**COLLARDS, GARDEN BEETS, KALE, LETTUCE, MUSTARD GREENS, RADISHES, RUTABAGAS, SPINACH, TURNIPS** - To control flea beetles, harlequin bug and leafhoppers, apply at the rate of 10 to 20 pounds per acre. To control armyworms, imported cabbageworm, diamond back moth larvae, corn earworm, tarnished plant bug and stink bugs, apply 20 to 40 pounds per acre or 1 to 2 ounces to 50 feet of row. For best control of corn earworm on lettuce heads begin to form, use 40 pounds per acre in a 7 day schedule. Will not control cabbage looper.

**CUCUMBERS, SQUASH, CANTALOUPE, PUMPKINS, MELONS** - To control pickleworm, armyworms, cucumber beetles, squash bug, flea beetles, apply at the rate of 20 pounds per acre or 1 ounce to 50 feet of row.

**CORN** - To control corn earworm, European corn borer, fall armyworm, sugarcane beetle, and Japanese beetles, apply at the rate of 30 to 40 pounds per acre or 2 ounces to 50 feet of row. For budworm damage in the whorl, apply to entire plant and repeat as necessary. For insects attacking silks and ears, apply at 1 to 3 day intervals starting when first silks appear and continue until silks begin to dry. Three or more applications may be required depending on severity of infestation.

**CARROTS** - To control flea beetles, leafhoppers, spiderbugs, lygus and stink bug, apply at the rate of 20 to 40 pounds per acre or 1 to 2 ounces to 50 feet of row.

**CORN** - To control armyworm and stink bugs, apply at the rate of 20 to 40 pounds per acre or 1 to 2 ounces to 50 feet of row.

**TOMATO, EGGPLANT, PEPPER** - To control tomato fruitworm, fall armyworm, flea beetles and leafhoppers, apply at the rate of 20 pounds per acre or 1 ounce to 50 feet of row. For control of tomato hornworm, European corn borer and stink bugs, apply at the rate of 20 to 40 pounds per acre.

**POTATO** - To control Colorado potato beetle, apply at the rate of 10 pounds per acre. For control of leafhoppers and flea beetles, apply at 20 pounds and for lygus bugs, apply at 40 pounds per acre.

### SMALL FRUIT

**BLACKBERRIES, BOYSENBERRIES, DEWBERRIES, LOGANBERRIES, RASPBERRIES** - To control Japanese beetles, leafrollers, leafhoppers and raspberry aphid, apply at the rate of 40 pounds per acre.

**BLUEBERRIES** - To control blueberry maggot, cherry and cranberry fruitworms and Japanese beetles, apply at the rate of 30 to 40 pounds per acre. Apply 3 weeks before harvest and repeat 10 days later if necessary.