SECURITY DIPEL® DUST BIOLOGICAL INSECTICIDE

ACTIVE INGREDIENT:

Bacillus thuringiensis subsp. kurstaki

TOTAL 100.000%

Potency: 240 International Units per mg. (at least 0.38 billion viable spores per gram). Equivalent to potency of 0.11 billion International Units per pound of this product. Potency units should not be used to adjust use rated beyond those specified

in the Directions for Use Section.

 ullet Dipel is a registered trademark of Abbott Laboratories, North Chicago, Illinois.

KEEP OUT OF REACH OF CHILDREN CAUTION

See Back Panel for Additional Precautionary Statements

Manufactured For: Security Products Co. of Delaware, Inc.

Minneapolis, Minnesota 55459

EPA Reg. No. 56644-19 EPA Reg. No. 1386-0H-1

)

NET CONTENTS: 4 Pounds

ACCEPTED

MAY 1 0 1993

Under the Federal Insecticide, Fungicide and Rodenticide Act. or amond id, for the pesticide registered under EPA Reg. No. 56644-19

BEST AVAILABLE COPT

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

ï

CAUTION: Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.

STATEMENT OF PRACTICAL TREATMENT

If in eyes, flush with plenty of water. If on skin, wash with soap and water. Remove contaminated clothing and launder before reuse. If inhaled, remove victim to fresh air. Apply respiration if indicated.

FOR EMERGENCY MEDICAL INFORMATION Call 1-800-228-5635, Extension 138.

ENVIRONMENTAL HAZARDS

Do not contaminate water when disposing of equipment washwaters.

)

)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DIPEL DUST is a biological insecticide intended for use against chewing larvae (caterpillars) of certain lepidopterous species. Apply DIPEL DUST when pests are first noticed and repeat as necessary. Thorough plant coverage is necessary for control.

BROCCOLI, CABBAGE, CAULIFLOWER, COLLARDS, KALE, MUSTARD GREENS, TURNIP GREENS: To control Cabbage Looper and Imported Cabbage Worm, apply 2 to 3 ounces per 50 foot of row.

CELERY, LETTUCE, MELONS, POTATOES, SPINACH, TOMATOES, CUCUMBERS: To control Cabbage Looper, apply 2 to 3 ounces per 50 foot of row.

TOBACCO AND TOMATOES: To control Hornworm, apply 1 to 2 ounces per 50 foot of row.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store this product in original container in a cool, dry place inaccessible to children and pets. Store at temperatures below 90°F.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Do not reuse empty container. Securely wrap original container in several layers of newspaper and discard in trash.

BEST AVAILABLE COPY

Warranty and Limitation of Damages

ķ

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and Buyer assumes the risk of any use contrary to such directions. Seller makes no other express or implied warranty, including any other express or implied warranty of Fitness or of Merchantability, and no agent of Seller is authorized to do so except in writing with a specific reference to this warranty. In no event shall Seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

/28

