

ACCEPTED

DEC 16 1991

for the
5644-84




DIAZINON

Yard & Vegetable Garden Granules

BEST AVAILABLE COPY

NOT TO BE USED ON GOLF COURSES OR SOD FARMS



DIAZINON

Yard & Vegetable Garden Granules

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of throat with finger. Repeat until vomit field is clear. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing. IF INHALED: Remove victim to fresh air and apply respiration if indicated. IF ON SKIN: Wash with soap and water. IF IN EYES: Flush with water for 15 minutes and call a physician immediately. HAVE TO PHYSICIAN: Remove patient to hospital. DIAZINON is also ocular and may be administered in conjunction with appropriate buffer.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing dust and contact with eyes, skin, or clothing. Wash thoroughly after handling and before eating or smoking. May cause contact dermatitis following repeated contact with skin of susceptible individuals. Avoid repeated contact with skin. If sensitization reactions result, consult a physician. Do not store near food or feed products. Food utensils such as spoons or measuring cups must not be used for food purposes after use in measuring pesticides. Do not contaminate ornamental fish ponds. Do not allow children or pets on treated areas until granules have been watered into the soil and the grass or soil is dry.

ENVIRONMENTAL HAZARDS

This pesticide is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lands in Monroe County, New York, between November 1 and May 26. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping granules. Where irrigation (watering) is recommended in the directions, or if watered, La. ducks or geese, can be expected in the treated area after treatment (except spot treatment), apply a minimum of 1/4 inch of water immediately after applying this product. Stop irrigation before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes, and estuaries. Do not apply directly to water or wetlands (marshes, bays, marshes, and potholes). Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

IN LAWNS SECURITY: DIAZINON YARD AND VEGETABLE GARDEN GRANULES KILLS:

Ants	Chinch Bugs	Fleas	Springtails
Armyworms	Clover Mites	Lawn Moths	White Grubs of
Bermudagrass Mites	Crickets	Leatherjackets	Japanese Beetles
Billbugs	Cutworms	Millipedes	European Chafers
Brown Dog Ticks	Copper Wasps	Sod Webworms	Southern Chafers
Chiggers	Cowbugs	Sowbugs	

IN GARDENS SECURITY: DIAZINON YARD AND VEGETABLE GARDEN GRANULES KILLS:

Cutworms	Onion Maggots	Vinegar Flies
More Crickets	Root Maggots	Wireworms

OVERALL LAWN APPLICATION

SECURITY® Diazinon Yard & Vegetable Garden Granules controls the following lawn insects on both grass and sedge/foam lawns: Ants, Armyworms, Bermudagrass Mites, Brown Dog Ticks, Chiggers, Chinch Bugs, Clover Mites, Crickets, Cutworms, Earwigs, Fleas, Lawn Moths (Sod Webworms), Leatherjackets, Millipedes, Sowbugs, Springtails (Collembola).

Apply SECURITY® Diazinon Yard & Vegetable Garden Granules with a fertilizer spreader. See spreader setting chart on bag. Apply when grass is dry. Water lawn lightly after application. Repeat as necessary for application with other types of equipment, apply at the rate of 2 lbs. per 1,000 sq. ft. (an area 20' x 50') or 3.2 oz. per 100 sq. ft. (an area 10' x 10').

IMPORTED FIRE ANTS: To aid in the control of imported fire ants in lawns and other recreational areas, apply 1/2 cup over and 2 ft. around each fire ant mound. Immediately after application, water in thoroughly. A minimum of 1 gallon of water should be applied to each mound area. Repeat as necessary. Apply gently to avoid disturbing ants. Use of high pressure watering equipment may disturb the ants and cause migration, reducing product effectiveness.

NOTE: For best results, apply in cool weather, 65-80°F, or in early morning or late evening hours. Test new mounds as they appear.

BILLBUGS: WHITE GRUBS OF JAPANESE BEETLE, EUROPEAN CHAFER, SOUTHERN CHAFER, ATLANTIC SPRAETULUS (Horn beetle): Apply 2 lbs. per 1,000 sq. ft. of lawn. Water grass thoroughly after application. For billbug control, apply when activity is first observed or when chafers or horn grubs indicate damage from these insects. Repeat as necessary. For grub control, except during beetle, treat anytime between late July and early October. For dung beetle grub control, apply in spring or early summer when damage is first noticed. Water grass thoroughly with 1/4-1/2 in. of water after application. Test again in late summer or early fall if a second generation occurs.

HYPERODES WEEVILS (New York only): Apply 2 lbs. per 1,000 sq. ft. of grass. Water grass thoroughly after application. Treat problem areas in mid-April and again in mid-May.

NOTE: The spreader settings listed below are approximate. Accurately weigh a few pounds into your spreader and apply to a measured area. Then adjust spreader setting if necessary, in order to assure correct rate of application.

LAWN SPREADER SETTING NO.

(on dialer 2 lbs. per 1,000 sq. ft.)

Spreader Make & Model	Rate or (Dialer) Lawn
Control Empire	
Model No. 1002-0	6 1/2
Control Empire Lawn	
Spreader No. 1022-0	6 1/2
Control Model 81	
Rate	4 1/2
Control Model 105	6 1/2
Orion King Spreader	
Model 5000	4 1/2
Orion Broadcast Spreader	
Model 4000	4 1/2
J. C. Pomeroy	
Model 8000	7 1/4
J. C. Pomeroy	
Rate Spreader	5
Scott Model 35	4
Scott Chalkman	
Model 671 10100	6

SPOT TREATMENT: Apply 1 1/2 teaspoons SECURITY® Diazinon Yard & Vegetable Garden Granules over and around each ant hill. Digger Wasps: Apply 1 1/2 teaspoons of SECURITY® Diazinon Yard & Vegetable Garden Granules in and around each nest opening. Apply at dusk when wasps are not active.

BAND TREATMENT AROUND HOUSE FOUNDATION

The following pests are commonly found around or near foundations of houses as well as in lawns. Control of these pests may enter the house.

Ants	Chinch Bugs	Millipedes	Springtails
Clover mites	Earwigs	Sowbugs (Pillbugs)	(Collembola)

Apply a 5-foot band of SECURITY® Diazinon Yard & Vegetable Garden Granules next to the house foundation. Distribute granules uniformly using a rate of 3.2 oz. for each 100 sq. ft. of area (5' x 20') treated. Repeat as necessary.

GARDEN INSECT CONTROL

SECURITY® Diazinon Yard & Vegetable Garden Granules controls certain garden insects on crops listed below. Apply just prior to planting and immediately mix thoroughly into soil to the specified depth. For time beans, snap beans, and peas, use on succulent varieties only. Do not use on commercially-grown potatoes which will be hand-harvested.

Cucumbers (surface and subterranean):	Peas	Summer Squash
Brussels	Peas	Sweet Corn
Cabbage	Peppers	Sweet Chard
Cauliflower	Peppers	Tomatoes
Corn	Peppers	Watermelons
Corn	Peppers	Winter Squash
Corn	Peppers	
Corn	Peppers	

Apply 7 to 14 oz. per 500 sq. ft. and mix into soil to a depth of 2-3 inches for surface cucumbers, and 3-4 inches for subterranean cucumbers. For potatoes, mix into soil 4-6 inches deep.

Male Crickets:	Corn	Onions	Summer Squash
Brussels	Corn <td>Onions<td>Sweet Potatoes</td></td>	Onions <td>Sweet Potatoes</td>	Sweet Potatoes
Cabbage	Cucumbers <td>Peas<td>Sweet Chard</td></td>	Peas <td>Sweet Chard</td>	Sweet Chard
Cauliflower	Endive (Escarole) <td>Peppers<td>Tomatoes</td></td>	Peppers <td>Tomatoes</td>	Tomatoes
Corn	Kale <td>Peppers<td>Watermelons</td></td>	Peppers <td>Watermelons</td>	Watermelons
Corn	Lettuce <td>Peppers<td>Winter Squash</td></td>	Peppers <td>Winter Squash</td>	Winter Squash
Corn	Lima Beans <td>Peppers<td></td></td>	Peppers <td></td>	
Corn	Muskmelon <td>Peppers<td></td></td>	Peppers <td></td>	

Apply 3 1/2 oz. per 500 sq. ft. and mix into soil to a depth of 1-2 inches.

Wireworms:	Corn	Onions	Summer Squash
Brussels	Corn <td>Onions<td>Sweet Potatoes</td></td>	Onions <td>Sweet Potatoes</td>	Sweet Potatoes
Cabbage	Cucumbers <td>Peas<td>Sweet Chard</td></td>	Peas <td>Sweet Chard</td>	Sweet Chard
Cauliflower	Endive (Escarole) <td>Peppers<td>Tomatoes</td></td>	Peppers <td>Tomatoes</td>	Tomatoes
Corn	Kale <td>Peppers<td>Watermelons</td></td>	Peppers <td>Watermelons</td>	Watermelons
Corn	Lettuce <td>Peppers<td>Winter Squash</td></td>	Peppers <td>Winter Squash</td>	Winter Squash
Corn	Lima Beans <td>Peppers<td></td></td>	Peppers <td></td>	
Corn	Muskmelon <td>Peppers<td></td></td>	Peppers <td></td>	

Apply 7 to 14 oz. per 500 sq. ft. and mix into soil to a depth of 4-8 inches.

BROCCOLI, CABBAGE, CAULIFLOWER: ROOT MAGGOT: Apply 7 to 11 oz. per 500 sq. ft. over entire seedbed before planting or transplanting and mix into the soil to a depth of 3 to 4 inches.

ONIONS: CHIRN MAGGOTS: Apply 7 - 14 oz. per 500 sq. ft. Broadcast just before planting and mix into the top 3-4 inches of soil.

NOTE: Diazinon may not control nematophagous - resistant onion maggots.

SWEET POTATOE (South Carolina only): WIREWORMS: Apply 11 oz. per 500 sq. ft. over the top of sweet potato plant foliage when roots begin to enlarge.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Store pesticide in original container and place in a locked storage area. Do not reuse container. Security wrap original container in several layers of newspaper and discard in trash.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

EPA Est. 769-GA-1(A) 59144-GA-1(B) 6720-FL-1(C) 51036-GA-1(D) 19713-GA-1(E) 1386-OH-1(F) Establishment location is indicated by alphabet letter following lot number.

Manufactured For
The Security Products Co.
of Delaware, Inc.
Minneapolis, MN 55459



DIAZINON MIXTURE, DRY



NA2783, RQ