



Net Weight: 12 1/2 lbs. or
5.67 Kilograms

PM 12 54,44-35

Bug Off Dust

Contains SEVIN® and DIPEL® Insecticides

ACTIVE INGREDIENTS

Bacillus thuringiensis, Berliner Potency of 240 International Units per mg. (at least 0.38 billion viable spores per g.) 0.048%

Carbaryl (1-naphthyl N-methyl carbamate) 5.000%

INERT INGREDIENTS:

94.952%

EPA Reg No. 769-480

TOTAL 100.000%

EPA Est. 769-GA-1

* SEVIN is a registered Trademark of Union Carbide Corp. for carbaryl insecticide.

* DIPEL is a registered trademark of Abbott Laboratories.

* Equivalent to potency of 0.11 billion International Units per pound of this product.

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

CAUTION Harmful if swallowed, inhaled or absorbed through the skin. Do not breathe dust. Do not take internally. Avoid skin contact. Avoid getting in eyes. Wash hands before eating. Wash all contaminated clothing with soap and water before reuse. Avoid contamination of foodstuffs, feed, troughs, and water receptacles.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

ENVIRONMENTAL HAZARDS

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DISPOSAL

Do not reuse empty container. Wrap container and put in trash collection.

NOTICE: The seller makes no implied warranty of merchantability nor any other warranties which extend beyond the description on this label.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

This product is a combination of Bacillus Thuringiensis, a biological insecticide, specifically for control of chewing caterpillars and Carbaryl, a broad spectrum insecticide, and will control the

following insect pests on vegetables in the home garden:

- | | |
|---------------------|--------------------------|
| Mexican Bean Beetle | Imported Cabbageworm |
| Bean Leaf Beetle | Diamond-Back Moth Larvae |
| Cucumber Beetle | Cabbage Looper |
| Leafhoppers | Tarnished Plant Bug |
| Flea Beetle | Pickleworm |
| Japanese Beetle | Mellonworms |
| Lygus Bug | Squash Bug |
| Stink Bug | European Corn Borer |
| Lima Bean Pod Borer | Fall Armyworm |
| Corn Earworm | Spittlebug |
| Sap Beetle | Tomato Fruitworm |
| Harlequin Bug | Tomato Hornworm |
| Armyworms | Colorado Potato Beetle |

FOLIAGE APPLICATION: Apply dust at the rate of 2 oz. per 50 feet of row (40 lbs. per acre) at the first sign of insects using a good mechanical or hand-powered duster. Repeat as necessary for control. For best results, dust in early morning or late evening when temperature is low and air is still. Apply dust so that all parts of plants are contacted, especially upper and lower leaf surfaces. The number in () following each crop indicates days required between last application and harvest; (0) indicates it may be used up to and including day of harvest. This product may be used on the following vegetables:

- | | |
|---------------------------------|--------------------------------|
| Beans (0) | Spinach (14) |
| Cabbage (3) | Turkeys (root 3, tops 14)..... |
| Broccoli (3) | Cucumbers (0) |
| Brussels Sprouts (3) | Squash (0)..... |
| Caul. flower (3) | Cantaloupe (0) |
| Kohlrabi (3) | Pumpkins (0)..... |
| Collards (14) | Melons (0)..... |
| Garden Beets (roots 3, tops 14) | Corn (0) |
| Kale (14) | Carrots (0) |
| Lettuce (head 3, leaf 14) | Okra (0) |
| Mustard Greens (14) | Tomato (0)..... |
| Radishes (3) | Eggplants (0) |
| Rutabaga (3) | Peppers (0) |
| | Potatoes (0) |

769-480
Worthington Chemical Works, Inc.
Manufacturers Ft. Valley, Ga.
JUN 10 1980
Under the Federal Insecticide, Fungicide, and Rodenticide Act