

SECURITY

**Dursban
Lawn Insect Spray**

Can also be used on Ornamentals

ACTIVE INGREDIENTS:

Chlorpyrifos (0,0-diethyl 0-(3,5,6 trichloro-2-pyridyl) phosphorothioate).....6.15%

INERT INGREDIENTS:.....93.85%
TOTAL 100.00%

EPA Reg. No. 56644-39
EPA Est. 769-GA-1

Dursban is a registered trademark of the Dow Chemical Co. This product contains 1/2 pound chlorpyrifos per gallon.

NET CONTENTS
1 PINT

Graphics

Sod Webworm

Chinch Bug


Ant

**KEEP OUT
OF REACH
OF CHILDREN
WARNING**

(See additional precautionary statements in attached pamphlet.)

**Manufacturers
Security Lawn &
Garden Products Co.
Atlanta, Georgia**

Not Reviewed. Registrant claims to be in accordance with Reregistration of Pesticide Product Guidance for Phase 2 Response, Page 2.12, Active to Inert Change in Status.



FOLD OUT LABEL

OPEN ALONG PERFORATION

SECURITY

**Dursban
Lawn Insect Spray**

NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

**NET CONTENTS
1 PINT**

**Manufacturers
Security Lawn &
Garden Products Co.
Atlanta, Georgia**

FOLD OUT LABEL

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician immediately. DO NOT induce vomiting. **IF INHALED:** Remove victim to fresh air. Apply respiration if indicated. **IF ON SKIN:** Wash immediately with soap and water. **IF IN EYES:** Flush eyes for at least 15 minutes with water. Call physician immediately.

NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.


PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

WARNING: Keep out of reach of children. May be fatal if swallowed. May be absorbed through skin. May be injurious to the eyes and skin. Do not take internally. Do not get in eyes or on skin. Do not smoke while using. Wash thoroughly after handling. Wash contaminated clothing before reuse. Flush contaminated eyes with plenty of water and get medical attention. Avoid breathing spray, mist or vapors. Do not spray vegetable gardens or other plants used for food purposes. Keep away from food, feedstuffs and domestic water supplies. Keep container closed. Keep children and pets off treated areas until spray has dried.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated.



FOLD OUT LABEL - cont'd.

This product is highly toxic to bees exposed to direct treatment or residues on plants. Protective information may be obtained from your Cooperative Agricultural Extension Service. Apply this product only as specified on this label.

PHYSICAL OR CHEMICAL HAZARDS

Combustible. Do not use, pour, spill or store near heat or open flame. Handle concentrate in ventilated area.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

FOR USE ON TURF


Use SECURITY Dursban Lawn Insect Spray to control chinch bugs, sod webworms (lawn moths), ants, brown dog ticks, crickets, cutworms, earwigs and grasshoppers infesting home lawns and other ornamental and recreational turf areas. Mix SECURITY Dursban Lawn Insect Spray with water according to directions given in the following table and apply as a coarse (large droplet) spray using a low pressure sprayer such as a knapsack type, hose-attached applicator or other similar equipment. If it is impractical to use the quantity of water listed in the table, apply in at least one gallon of water per 1000 square feet and sprinkle with the recommended amount of water after application. For best results, lawn should be moist at the time of application and care should be taken to obtain complete and uniform coverage of pest infested areas. Spray when pests first appear, retreat when needed.

DOSAGE AND DILUTIONS FOR USE ON TURF

Area to be treated	Amount of Dursban Spray	Dilute with water to give
500 square feet	3 fluid ounces	15 gallons
1,000 square feet	6 fluid ounces	30 gallons
5,350 square feet	1 quart	150 gallons
10,700 square feet	2 quarts	325 gallons
One Acre	2 gallons	1,300 gallons

1 fluid ounce = 2 tablespoons = 6 teaspoons

To control certain white grubs (larvae of European chafer and Japanese beetle), use two to four times the amount of SECURITY Dursban Spray shown in the TURF table., i.e., 12 to 24 fluid ounces per 1000 square feet or 4 to 8 gallons per acre. Apply in enough water to obtain uniform coverage of the area treated followed immediately by enough watering to wash the insecticide into the turf. For best results, treat when grubs are young and actively feeding near the soil surface, usually during last July and August. Consult your local Agricultural Extension Service specialist for proper time to treat in your area. To control hyercdes (turfgrass) weevils use twice the amount of SECURITY Dursban Spray shown in the TURF table i.e., 4 gallons diluted in 1300 gallons of water per acre. Spray suspected problem areas in mid-April and again in mid-May. Consult your local Agricultural Extension Service



FOLD OUT LABEL - cont'd.

specialist for best time to treat in your area.

NOTE: To help prevent infestation of buildings by ants, crickets, earwigs and sowbugs (pillbugs) treat a band of soil 6 to 10 feet wide around and adjacent to the building, also the building foundation to a height to 2 to 3 feet, where pests are active and may find entrance. Use the same dosage and dilution recommended for control of these pests on turf as shown in the TURF table. Apply as a coarse spray to thoroughly and uniformly wet the band area.

FOR USE ON ORNAMENTAL PLANTS: Use SECURITY Dursban Spray to control aphids, bagworms, cutworms, grasshoppers, leafhoppers, mealybugs, mites, spittlebugs, thrips (exposed) and whiteflies on flowers, shrubs, vines, shade and flowering trees and evergreens found to be infested with these pests. Dilute with water according to directions given in the following table and apply using a suitable low pressure sprayer such as a knapsack type or hose-attached applicator in a manner to provide complete and uniform coverage of infested plants. Avoid over-spraying to the point of excessive runoff. Treat when pests appear and repeat at 7 to 10 day intervals if needed. NOTE: Do not use on azaleas, camellias, poinsettias, rose bushes or variegated ivy because of possible injury to these plants.

DOSAGE AND DILUTIONS FOR USE ON ORNAMENTAL PLANTS

Pests Controlled	Amount of Dursban in water to give		
	3 gals.	10 gals.	100 gals.
Aphids, Bagworms Grasshoppers, Mites Spittlebugs and Whiteflies	2 fl. oz.	6.4 fl. oz.	2 qts.*
Cutworms, Mealybugs, Leafhoppers and Thrips (Exposed)	4 fl. oz.	12.8 fl. oz.	1 gal.**

*Equivalent to 4 ounces of chlopyrifos per 100 gallons.

**Equivalent to 8 ounces of chlopyrifos per 100 gallons.

NOTE: Mosquitoes coming to rest on areas treated for control of turf and ornamental plant pests will be killed for varying periods of time after treatment, depending on exposure of treated areas to weathering conditions.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage disposal. Store pesticide in original container and place in a locked storage area. Do not reuse container. Rinse thoroughly and securely wrap original container in several layers of newspaper and discard in trash.

NOTICE: The seller makes no implied warranty of merchantability nor any other warranties which extend beyond the description on this label.

