

1055
pm21
56644-42

SECURITY
FUNGI-GARD

Broad Spectrum Liquid Fungicide
For Use On Vegetables,
Roses, Flowers and Lawns

Contains Chlorothalonil

ACTIVE INGREDIENT:	
Chlorothalonil (tetrachloroisophthalonitrile)*.....	11.24%
INERT INGREDIENTS:.....	88.76%
TOTAL	100.00%

This product contains the Toxic Inert Ingredient
Paraformaldehyde.


*Covered under U.S. Patent No. 3,290,353 & 3,331,735.
This product contains 1 lb. chlorothalonil per gallon.

KEEP OUT OF REACH OF CHILDREN

WARNING
NET CONTENTS: 1 FLUID PINT

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician Immediately. IF INHALED: Remove
victim to fresh air. Apply artificial respiration if
indicated. IF ON SKIN: Remove contaminated clothing and wash
affected areas with soap and water. IF IN EYES: Flush eyes with
plenty of water. Call a physician immediately.
NOTE TO PHYSICIAN: Persons having allergic reaction respond to
treatment with antihistamines or steroid creams and/or systemic
steroids.



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Corrosive, causes severe eye damage. Do not get in eyes; wear goggles or eye shield when handling this product. Keep out of reach of children. May be harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with skin, eyes and clothing. Chlorothalonil is a skin sensitiser. Do not breathe spray mist. Wash hands, arms, and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse. Do not contaminate feed or foodstuffs. Keep children and pets off treated areas until completely dry. Apply this product only as specified on this label.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, aquatic invertebrates, and marine/estuarine organisms. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. NOTE TO USER: This product may produce temporary allergic side effects characterized by redness of eyes, mild bronchial irritation and redness or rash on exposed skin areas. Persons having allergic reaction should contact a physician. Wear long-sleeved shirt, long pants, and gloves while mixing, loading and applying this product.

SHAKE WELL BEFORE USING!

LAWN AND ORNAMENTAL TURF

SECURITY Fungi-Gard will control many turf diseases when used according to the listed rates. Begin application when disease threatens or when conditions are favorable for disease development. Do not mow or water turf for 24 hours following treatment. Apply SECURITY Fungi-Gard in adequate amount of water to provide complete coverage. This may vary from 5 to 15 gallons per 1000 square feet. Under severe disease conditions, use the curative rates and spray on a 7 day schedule. Do not graze treated areas. Do not feed clippings to livestock.

Disease	Spray Interval	Rate Per 1000 Square Feet		Copper
		Preventative*	Curative**	
Spot	7-10 days	1 1/2 to 2 1/4 pts.	2 1/4-2 3/4 pts.	
Curcularia Leaf Spot	7-10 days	3/4 to 1 1/2 pts.	2 1/4-2 3/4 pts.	
Dollar Spot	7-14 days	3/4 to 1 1/2 pts.	1 1/2-2 3/4 pts.	
Gray Leaf Spot	7-10 days	3/4 to 1 1/2 pts.	2 1/4-2 3/4 pts.	
Helminthosporium Leaf Spot and Melting Out	7-10 days	3/4 to 1 1/2 pts.	1 1/2-2 3/4 pts.	
Large Brown Patch	7-10 days	3/4 to 1 1/2 pts.	2 1/2-2 3/4 pts.	
Red Thread	7-10 days	3/4 to 1 1/2 pts.	2 1/4-2 3/4 pts.	
Stem Rust of Bluegrass	7-14 days	1 1/2 to 2 1/4 pts.	2 1/4-2 3/4 pts.	

*Recommended rates for preventing disease establishment. Use lower rates when disease conditions are light to moderate, higher indicated rates when conditions are severe. **Rates for use on a 7 day schedule when disease is present. Higher indicated rates should be applied under severe conditions.

ORNAMENTALS

SECURITY Fungi-Gard will control a number of diseases of ornamentals. Apply 3 tablespoons per gallon of water (except Roses, apply 2 tablespoons per gallon) beginning at first sign of disease and repeat at 7 to 14 day intervals throughout the season. Make applications on a 7 day schedule during periods of wet weather or severe disease conditions. Application should be thorough to obtain adequate coverage of the foliage, Almond (Flowering): Monilinia blossom and branch blight; Ash, leaf spots; Azalea*** (Rhododendron SPP.): Ovinia flower blight and Phytophthora dieback; Carnation alternaria leaf spot and Botrytis blight; Cherry Lurel: shot-hole leaf spot; Chrysanthemum and Daisy: Ascochyta ray blight, gray mold and septoria leaf spot; Crabapple: Cedar apple rust, Sphaeropsis leaf spot and scab; Dogwood: Septoria leaf spot; Firethorn: Scab; Geranium: Botrytis blight and rust; Gladiolus: Botrytis Blight and Curvularia leaf spot; Hawthron: Rust; Hollyhock: Rust; Iris: Botrytis blight and leaf spot; Japanese Androneda (Pieris spp.): Phytophthora die-back; Leather Leaf-Fern (Florida Only): Asochyta blight, Cercospora leaf spot Cylindrocladium leaf spot and Rhizoctonia blight; Lily: Botrytis blight; Mahonia: Rust; Mountain Laurel: Leaf spot; Oyster Plant (Rhoeo Discolor) (Florida Only): Tan leaf spot; Petunia***: Botrytis and phytophthora blight; Philodendron (Florida Only): Phytophthora blight and Dactylaria leaf spot; Photinia (Red Tips): Fabraea (Entomosporium) leaf spot; Privet: Cercospora leaf spot; Purple Leaf Sand Cherry: Monilinia blossom and branch blight; Quince (Flowering): Monilinia blossom and branch blight; Rose: Apply 2 tablespoons per gallon of water for control of black spot and blossom blight; Viburnum: Powdery mildew; Zinnia: Powdery mildew;***Discoloration of blooms has been noted, at least on certain varieties.

VEGETABLES

In the garden, apply SECURITY Fungi-gard when condtions favor disease development. Follow the recommended application timing and intervals for the vegetables listed. Do not apply to any crop not listed on this label. Rates are expressed in tablespoons per gallon of water. Consult the chart below for the amount of dilute spray to apply for specific row spacing. All rates are based on a broadcast rate of 150 gallons per acre. During early stages of growth, proportionately less spray will be required. Use lower rates when disease conditions are light to moderate, higher indicated rate, when conditions are severe. Always spray for thorough coverage. Follow the time interval between last application and harvest indicated by the number in 9) following each vegetable. A(0) indicates product can be used through day of harvest. Do not graze treated areas or feed treated plant parts or ensilage to livestock. AMOUNT OF DILUTE SPRAY PER 100 FEET OF ROW WHEN SPACING IS: 12 inches, 1/3 gal; 24 inches, 2/3 gal.; 36 inches, 1 gal.; 48 inches, 1 1/3 gal. SNAP BEANS (7): To control Rust, apply 3 2/3 tablespoons per gallon of water. Begin application during early bloom or when disease first threatens. Repeat weekly or as necessary to maintain control. CABBAGE, CAULIFLOWER, BROCCOLI, BRUSSELS SPROUTS (0): To control Alternaria leaf spot (Southern states only) and Downy mildew, apply 2 tablespoons per gallon of water.

Begin application after transplants are set in the field or shortly after emergence of field seeded crop or when conditions favor disease development. repeat at 7 to 10 day intervals or as necessary to maintain control.

CANTALOUPE, HONEYDEW MELON, MUSKMELON, WATERMELON, PUMPKIN, SQUASH (0): To control Anthracnose and Downy Mildew, apply 2 1/3 tablespoons per gallon of water. Start when disease threatens. gummy Stem blight, Leaf Blight, Scab, Cercospora Leaf Spot and Powdery Mildew (on cantaloupes, muskmelons, honeydew, except in Southwestern states), apply 2 1/3 to 3 2/3 tablespoons per gallon of water. Begin when plants are first true leaf stage or when conditions are favorable for disease development. Repeat at 7 day intervals. Under severe disease conditions, shorten spray interval.

CARROTS (0): To control Early Blight and Late Blight, apply 2 to 2 1/3 tablespoons per gallon of water. Start when disease threatens. Repeat at 7 to 10 day intervals or as necessary to maintain control.

CUCUMBERS(0): To control Target Spot (Florida Only), Anthracnose and Downy Mildew, apply 2 to 2 1/3 tablespoons per gallon of water. To control Powdery Mildew (except Southwestern states), Gummy Stem Blight, Leaf Blight and Scab, apply 2 1/3 to 3 2/3 tablespoons per gallon of water. For suppression of Fruit Rot, apply 3 2/3 tablespoons per gallon of water. Begin applications when plants are in first true leaf stage or when conditions are favorable for disease development. Repeat applications at 7 day intervals. Under severe disease conditions, shorten spray intervals.

ONIONS (7 days for dry bulbs onions, 14 days for green onions): To control Botrytis Leaf Blight (Blast), apply 2 to 3 2/3 tablespoons per gallon of water. Begin application before disease appears and repeat at 7 to 10 day intervals. Under severe conditions, use higher rate and shorten spray interval. Do not apply on sweet Spanish onions. Do not apply more than 3 times per season on green onions.

PEANUTS (14): To control Cercospora Leaf Spots, apply 1 1/4 to 1 4/5 tablespoons per gallon of water. To control leaf rust, apply 1 4/5 tablespoons per gallon of water. Begin application when disease first appears. Repeat at 10 to 14 day intervals or as necessary to maintain control. Under severe conditions, use the higher rate.

POTATOES(0): To control Early Blight, Late Blight and Botrytis Vine Rote, apply 1 3/4 to 1 4/5 tablespoons per gallon of water. Begin applications when plants are 6 inches high or when disease threatens. repeat at 7 to 10 day intervals or as necessary to maintain control. Under severe disease conditions use the higher rate and shorten spray intervals.

TOMATOES(0): To control Early blight, Late Blight, Gray Leaf Spot and Gray Leaf Mold, apply 2 to 2 1/3 tablespoons per gallon of water. To control Anthracnose, apply 2 1/3 to 3 2/3 tablespoons per gallon. to control Rhizoctonia Fruit Rot and Botrytis Gray Mold, apply 3 2/3 tablespoons per gallon of water. Begin when disease threatens and repeat 7 to 10 day intervals. Under severe disease conditions, shorten spray intervals.

SOF5

STORAGE AND DISPOSAL STATEMENTS

STORAGE: Do not contaminate water, food or feed by storage or disposal. Store pesticide in original container and place in a locked storage area. Do not reuse container.

DISPOSAL: Rinse thoroughly and securely wrap original container in several layers of newspaper and discard in trash.

NOTICE: The seller makes no implied warranty of merchantability nor any other warranties which extend beyond the description on this label.

Distributed by:

SECURITY PRODUCTS COMPANY OF DELAWARE, INC.
City of Commerce, CA and Fort Valley, GA

EPA REG. NO. 56644-42

EPA EST. NO. 769-GA-1