

**WARRANTY STATEMENT**


Our recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal. STORAGE: Store product in original container in a locked storage area. CONTAINER DISPOSAL: Do not reuse empty container. Wrap container and put in trash.

**NET CONTENTS: 1 FLUID PINT**

Distributed By  
The Security Products Company  
of Delaware, Inc.  
City of Commerce, CA 90040  
Fort Valley, GA 31030



SECURITY

Malathion & Oil  
Citrus Spray

Controls Scale Insects,  
Spider Mites,  
Mealybugs & Whiteflies

ACTIVE INGREDIENTS:

Malathion (o,o-dimethyl phosphorodithioate  
mercaptosuccinate)..... 5.0%

INERT INGREDIENTS: ..... 95.0%


TOTAL 100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

(See additional precautionary statements on side panel.)

Not Reviewed. Registrant claims to be in accordance with  
Reregistration of Pesticide Product Guidance for Phase 2  
Response, Page 2.12, Active to Inert Change in Status.



## STATEMENT OF PRACTICAL TREATMENT

**IF SWALLOWED:** Call a physician or Poison Control Center. **IF INHALED:** Remove victim to fresh air. Apply respiration if indicated. **IF ON SKIN:** Wash immediately with soap and water. **IF IN EYES:** Flush eyes with water and get medical attention. **NOTE TO PHYSICIAN:** Malathion may cause cholinesterase inhibition. Atropine is antidotal.

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

**CAUTION.** Harmful if swallowed. Avoid breathing spray mist. Avoid contact with skin and eyes; wash thoroughly after using. Avoid contamination of feed and foodstuffs.

## ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish, aquatic invertebrates, and aquatic life stages of amphibians. Do not apply directly to water or wetlands (i.e., swamps, bogs, marshes, and potholes). Drift and runoff may be hazardous to aquatic organisms in areas near the application site. Do not contaminate water when disposing of equipment washwaters. **PROTECT BEES:** This product is highly toxic to bees exposed to direct treatment or residues remaining on treated areas. Do not apply when bees are actively visiting the crop, cover crop or weeds blooming in treatment area. Application should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

## PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

## DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

## USE PRECAUTIONS

Do not use this product on plants that are wilting or near wilting from lack of moisture. Do not apply with sulfur or lime sulfur. Do not apply within 3 weeks before or after application of sulfur or lime sulphur. Avoid spraying during periods of excessively high temperatures (above 90°F). Do not exceed 3 applications a year. **NOTE:** Do not apply to hibiscus, papaya, sensitive ferns such as Boston, Maiden Hair and Pteris and certain species of crassula.

GENERAL INFORMATION

Shake contents before using. Mix thoroughly with proper amount of water in spray tank and keep mixture well agitated during spraying. Do not allow mixture to stand. Insects are more difficult to control as they become older and larger. Unless otherwise noted, spray when insects first appear and repeat as needed. Spray upper and lower leaf surfaces of all leaves and branches thoroughly. Where snow scale is a problem, it is important that trunk limbs and twigs are thoroughly sprayed.

CITRUS (Grapefruit, Lemons, Limes, Oranges, Tangerines, Tangelos, Kumquats): For the control of Florida red scale, purple scale, snow scale, spider mites, whitefly and citrus mealybug, apply 2 1/2 tablespoons per gallon of water (1 quart per 25 gallons), preferably in June or July. Do not apply when trees are in bloom. Do not apply within 7 days of harvest.

AVACADO: For the control of wax scale, whitefly, and latania scale, apply 2 tablespoons (1 fl. oz.) per gallon of water, preferably in May, June or July. Do not apply within 7 days of harvest.

CAMELLIAS, GARDENIA, LIGUSTRUM AND IXORA: For the control of leaf scale, whitefly, citrus mealybug, soft scales, Florida red scales, and purple scale, use 2 tablespoons (1 fl. oz.) per gallon of water. Make 2 applications, 2 weeks apart, after blooming.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Do not reuse empty container. Wrap container and place in trash collection.

NOTICE: The seller makes no implied warranty of merchantability nor any other warranties which extend beyond the description on this label.

184121

NET CONTENTS: 1 FL. PINT

Security Products Company of Delaware, Inc.  
EPA Reg. No. 56644-56  
EPA Est. No. 769-GA-1

