

SECURITY
NATURE-GARD

Rotenone-Pyrethrin Insect Spray

ACTIVE INGREDIENTS:	
Rotenone.....	1.1%
Other Cube Extractives.....	2.2%
Pyrethrins.....	0.8%
INERT INGREDIENTS:	95.9%
TOTAL	<u>100.0%</u>


EPA REG. NO. 56644-71 EPA EST. 769-GA-1

Contains Petroleum Distillate

KEEP OUT OF REACH OF CHILDREN

WARNING: See additional precautionary statements on the side panel.

Not Reviewed. Registrant claims to be in accordance with Reregistration of Pesticide Product Guidance for Phase 2 Response, Page 2.12, Active to Inert Change in Status.



**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

WARNING! May be fatal if swallowed. Harmful if absorbed through the skin. Causes skin irritation. May cause substantial but temporary eye injury. Wear safety glasses. Do not get in eyes, on skin or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Do not induce vomiting. This product contains aromatic petroleum solvent. Aspiration may be a hazard. **IF IN EYES OR ON SKIN:** Flush with plenty of water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes, potholes). Runoff and drift from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

PHYSICAL AND CHEMICAL HAZARDS

DO NOT USE OR STORE NEAR HEAT OR FLAME. Flash Point minimum 120°F (48.9°C).

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FOR USE ON ORNAMENTAL AND FLOWERING PLANTS: such as poplar, linden, dogwood, willow, birch, boxelder, maple, elm, pine, spruce, cedar, yew, spirea, rose, lilac, zinnia, aster, petunia, geranium, dahlia, gladiolus, phlox, sweet pea, marigold, carnation, chrysanthemum and rose, mix this product with water as described below. Use hand or automatic sprayer as a coarse spray. Use sufficient spray mix to give complete coverage of the plant. Do not apply this product within one day of collection. Repeat every 5-10 days or as needed. Keep diluted spray away from light. Pests: aphids, leafhoppers, exposed thrips use 1 teaspoon per gallon of water; oblique-banded leaf-rollers, sawfly larvae, boxelder bugs, Japanese beetles, elm leaf beetle larvae, mites, lygus bugs, lace bugs, greenhouse whiteflies, stink bugs, garden flea hoppers, greenhouse leafhoppers use 2 teaspoons per gallon of water; cucumber beetles, rose chafers, ants, aster beetles, flea beetles use 4 teaspoons per gallon of water.

BEST AVAILABLE COPY

FOR USE ON VEGETABLE PLANTS: such as peas, bean, beet, cucumber, squash, muskmelon, potato, tomato, cabbage, spinach, lettuce, celery, asparagus, pepper, eggplant, onion, carrot and corn; mix with water as described below. Use hand or automatic sprayer as a coarse spray. Use sufficient spray mix to give complete coverage of the plants. Do not apply this product within one day of harvest. Repeat every 5-10 days or as needed. Keep diluted spray away from light. Pests: aphids, leafhoppers, exposed thrips use 1 teaspoon per gallon of water; cabbage worms, bean beetles, beet webworms, sod webworms, asparagus beetles, Japanese beetles, lygus bugs, whiteflies, garden fleahoppers, harlequin bugs, oblique-banded leaf rollers, stink bugs, celery leafleters, melon worms, tortoise beetles, garden webworms, alfalfa webworms and mites use 2 teaspoons per gallon of water; cucumber beetles, flea beetles, vegetable weevils, squash bugs (nymphs), squash vine borers, European corn borers 1 to 4 applications or more as eggs start to hatch and repeated at 5 to 7 day intervals use 4 teaspoons per gallon of water.

FOR USE ON FRUITS AND BERRIES: such as apple, pear, plum, grape, currant, gooseberry, blackberry, strawberry and blueberry. Use sufficient spray mix to give complete coverage of the plants. Do not apply this product within one day of harvest. Repeat every 5-10 days or as needed. Keep diluted spray away from light. Pests: aphids, leafhoppers, exposed thrips use 1 teaspoon per gallon of water. Imported currant worms, gooseberry fruit worms, raspberry fruit worms, blueberry maggots, cherry fruit flies, strawberry root worms (adult), rednecked borers, Japanese beetles and mites use 2 teaspoons per gallon of water; strawberry leafrollers and rose chafers use 4 teaspoons per gallon of water.

TO KILL FLEAS, TICKS AND LICE ON DOGS, CATS AND PREMISES: Do not use on puppies or kittens less than 4 weeks old. (Keep diluted spray away from light). ON DOGS: Do not use more than 1 fluid ounce (2 Tbsp.) per gallon on puppies. FOR FLEAS AND LICE: dilute this product at the rate of 1 fluid ounce (2 Tbsp.) per 1 gallon of water. For young animals, use 1/2 fluid ounce (1 Tbsp.) per 1 gallon of water. Apply directly to animal as a spray, sponge-on or a wash. Dip until animal's skin is wet. Do not rinse.

FOR TICKS: dilute this product at the rate of 2 fluid ounces (4 Tbsp.) per 1 gallon of water. For young animals, use 1 fluid ounce (2 Tbsp.) per 1 gallon of water. Apply directly to animal as a spray, sponge-on or a wash. Dip until animal's skin is wet. Do not rinse. Avoid contact with eyes, nose or mouth of animal. To prevent reinfestation, thoroughly spray the animal's living quarters with this product. ON CATS: Do not use on young kittens. On adult animals, do not use more than 1 fluid ounce (2 Tbsp.) per gallon. FOR FLEAS, TICKS AND LICE: dilute this product at the rate of 1 fluid ounce (2 Tbsp.) per gallon of water. Apply directly to animal as a spray, sponge-on or wash. Dip until animal's skin is wet. Do not rinse. Avoid contact with eyes, nose or mouth or animal. To prevent reinfestation, thoroughly spray the animal's living quarters with this product.

BEST AVAILABLE COPY