

PM 15 56644-86 Y
5/24/89

EPA-PI

Apply as a thorough foliar spray. Wet the plants to the drooping point. Try to hit the undersides of leaves and penetrate dense foliage.

FRUITS AND NUTS: Apples, Apricots, Blackberries, Cherries, Strawberries, Grapes, Grapes, Lemons, Oranges, Peaches, Walnuts. Controls Aphids, Apple Maggots, Spider Mites, Ticks (Clove or Brown, Two Spotted, European Red, Pacific, or Two Spotted Spider), Scale (Cottony-Cushion, San Jose), Cotton Moths, Fruit Leaf Rollers, Cherry Frost Fleas, Leafhoppers, Maggot (Apple), and Mealybugs. Use 1 Pint, or 3 gallons of water. Apply as a full cover spray. Start spraying after blooming, and again in the summer. Applications should be made according to infestation or when Aphids, Ticks, and crawler stages of Scale insects are first noticed. If warranted, repeat application at 7 to 10 day intervals. On apples, slight russetting may occur on some varieties such as Golden Delicious. Do not spray within the following number of days before harvest: Strawberries—3 days; Apricots, Cherries, Lemons, Grapes—10 days; Apples—14 days; Grapes, Lemons, Oranges, Peaches, Walnuts—21 days. Do not apply after fruits open or split.

VEGETABLES: Beans, Peas, Cabbage, Cauliflower, Broccoli, Radishes, Squash, Carrots, Lettuce, Cantaloupes, Cucumbers, Sweet Corn, Onions, Peppers, Potatoes, Tomatoes, Colorado Potato Beetle, Cucumber Beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Mites, Diamondback Moths, Flea Beetles, Imported Cabbage Worms, Dipterous Leaf Miners, Thrips, Colorado Potato Beetles, Fall Armyworms (Tomato), Onion Thrips, Southern Armyworms (Potato and Tomato). Use 1 Pint, or 3 gallons of water, covers 1000 sq. ft. of garden. Do not spray within the following number of days before harvest: 1 day—Peas, Tomatoes, Sweet Corn, 3 days—Cantaloupes; 5 days—Broccoli, Cauliflower; 7 days—Carrots, Cabbage, Cucumbers, Peppers, Squash; 10 days—Carrots, Lettuce, Onions, Radishes; 35 days—Potatoes.

STORAGE AND DISPOSAL

Storage: Store in original container in a cool place away from sports or open flame. Keep out of reach of children.
Disposal: Soak dry, wrap original container in several layers of paper, and discard in trash. Do not reuse container.
Notice: Buyer assumes all risk of use, storage and handling of this material not in strict accordance with directions given herewith.

Manufacturers
Security Lawn &
Garden Products Co.
Atlanta, Georgia

EPA REG. NO. 56644-86
EPA EST. NO. 769-GA.-1



ACTIVE INGREDIENTS:
80 methyl 0-(7-nitroindolin-6-methyl-4-pyromellityl)-
phenylcarbamate

INERT INGREDIENTS

Diazinon® Inertium of Ciba-Geigy

25 fl.
75.01
Lbs. 100 Lbs.

KEEP OUT OF REACH OF CHILDREN

WARNING: See back panel for additional precautionary statements.

NET 1 PINT
(0.473 L.)

CHACON DIAZINON® SPRAY
PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS:
Keep out of reach of children. May be fatal if swallowed. Do not breathe spray mist. May be absorbed through skin. Avoid contact with eyes, skin and clothing. Wash thoroughly after handling and before eating or smoking. Avoid contamination of food and feed. Do not use on household pets or humans. Do not eat or drink from treated surfaces. Do not use on household pets or humans. Do not eat or drink from treated surfaces. Statement of Practical Treatment: If swallowed, do not induce vomiting. Call a physician or Poison Control Center immediately. Never give anything by mouth to an unconscious or convulsing person. If inhaled, remove affected person from exposure. If in eyes, flush with plenty of water for at least 15 minutes. Get medical attention if irritation persists. If on skin, remove contaminated clothing and wash affected skin areas with soap and water. Note to Physician: Solvent presents aspiration hazard. Gastro-Intestinal irritation and/or material was taken internally. If symptoms of cholinesterase inhibition are present, atropine is antidotal. 2 PAM is also antidotal and may be used in conjunction with atropine.

ENVIRONMENTAL HAZARDS
This product is toxic to fish, birds, and other wildlife. Birds feeding in the treated areas may be killed. Do not apply directly to any body of water. Shrimp and Crab may be killed at application rate recommended on this label. Do not apply where these are important resources. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. This pesticide is highly toxic to bees exposed to direct treatment or indirect contamination in the treated area. Denatured when bees are actively visiting the crop. Cover crop, or weeds blooming in the treated area. Application should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

PHYSICAL OR CHEMICAL STANDARDS
Do not use, pour, spill or store near heat or open flame.
DIRECTIONS FOR USE
It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read entire label thoroughly. Apply this product only as specified on this label. DO NOT USE INSIDE THE HOME.
Determine size of spraying job. For lawns and vegetables measure the area to be sprayed. For all other applications, apply as thorough cover spray. Make applications when insects first appear. Be sure to spray underside of leaves. Repeat applications as needed or as stated in the directions. Use contents within 6 months after opening. (Note: Do not use on Ferns, Peacock's Tail, Azaleas, Peony, Pea, Gardenias and Orchids, since many may die off.)
LAWNS: Bent, Bermuda, Dechondra, Fescue, Merton, St. Augustine. Controls: Ants, Armyworms, Bermuda Grass Thrips, Crickets, Culicines, Earwigs, Fleas, Fruit Fleas, Lawn Boll-Bugs, Lawn Moths (Soil Webworms), Leafhoppers, Moles (Bermuda, Clover), Sawflies (Pillbugs), Springtails (Collembola), Rhododendrons, Scales. Use 1 Pint, or 3 gallons of water per 125 sq. ft. Thoroughly wet down grass a few hours before applying. Keep children and pets off of area until grass has completely dried. For Lawn Boll-Bugs, water lawn immediately after treatment.

TREES AND SHRUBS: Arborvitae, Azaleas, Birch, Boxwood, Camellia, Elm, Juniper, Maple, Oak, Pine, Poplar, Rhododendron, Spruce, Willow. Controls: Aphids, Bagworms, Leaf Miners, Miles (Clove, Cyclamen, European Red, Privet, Two Spotted Spider), Scale Crawlers (Cottony-Cushion, Lychee), Pine Needles Pt. Scale, San Jose, Soft. Use 1 Pint, or 3 gallons of water. Apply as thorough foliar spray. Wet both sides of leaves and branches to the drooping point. Try to penetrate dense foliage.
ROSES AND FLOWERS: Azaleas, Camellias, Crysanthemums, Carnations, Gladiolas, Lilies, Rhododendrons, Roses. Controls: Aphids, Euonymus Scale Crawlers, FleaBeetle, Dipterous Leaf Miners, Leafhoppers, Mealybugs, Thrips, Whiteflies. Use 1 Pint, or 3 gallons of water.

Not Reviewed. Registrant claims to be in accordance with
Registration of Pesticide Product Guidance for Phase 2
Response, Page 2.12, Active to Inert Change in Status.

68-42-03

84-843

BEST AVAILABLE COPY