

1/2/97

PM 23

57242-4

pg 1/5

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval expires 05-31-98



United States
Environmental Protection Agency
Washington, DC 20460

Registration
 Amendment
 Other

OPP Identifier Number
253373

Application for Pesticide - Section I

1. Company/Product Number 57242-4	2. EPA Product Manager Joanne I. Miller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) GLADEAMINE 3.8 2-4,D HERBICIDE	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) GLADES FORMULATING CORP. c/o RegWest Company P.O. Box 2220 Greeley, CO 80632-2220 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application NOTIFICATION
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Submit additional brand name notification pursuant to PR Notice 95-2. The ABN is *Weed-Amine 2,4-D Herbicide*. This notification is consistent with the provisions of PR Notice 95-2 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to the EPA. I further understand that if this notification is not consistent with the terms of PR Notice 95-2 and 40 CFR 152.46 this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	If "Yes" Unit Packaging wgt. No. per container		<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
* Certification must be submitted		If "Yes" Package wgt. No. per container			
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled				<input type="checkbox"/> Other _____	

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Kim Davis	Title Consultant/Agent	Telephone No. (Include Area Code) (970) 353-0611
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Consultant/Agent	
4. Typed Name Kim Davis	5. Date November 26, 1996	

245

WEED-AMINE

2,4-D HERBICIDE

For Selective Control of Many Broadleaf Weeds in Non-Crop Areas,
Grass Pastures, Rangelands and in Certain Crops

NOTIFICATION

Also for Control of Trees by Injection

Contains Dimethylamine Salt of 2,4-D*

ACTIVE INGREDIENT:		
2,4-Dichlorophenoxyacetic acid, dimethylamine salt*	46.3%	
INERT INGREDIENTS	53.7%	
	Total	100.0%

2,4-dichlorophenoxyacetic acid* - 33.4% - 3.3 lbs./gal.

*Isomer Specific by AOAC Method No. 978.05 (15th Ed.)

*Salts are the least volatile forms of 2,4-D and do not release enough vapors from treated areas to reduce yield of adjacent susceptible crops.

KEEP OUT OF REACH OF CHILDREN

WARNING

AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See back panel for additional precautionary statements

EPA REG. NO. 57242-4

EPA EST. 57242-FL-1

NET CONTENTS _____ GALS. (_____ L)

45035

07/94

BEST AVAILABLE COPY

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
& DOMESTIC ANIMALS

WARNING AVISO

Injurious to eyes. Harmful if swallowed. May cause skin irritation. Do not get in eyes, on skin, or on clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers, including persons repairing or cleaning equipment, must wear:

- Coveralls over short-sleeved shirt and short pants
- Waterproof gloves
- Chemical-resistant footwear plus socks
- Protective eye wear such as face shield or safety glasses (brow and temple protection recommended). Eye protection is not required for ground or aerial applicators in completely enclosed cabs or cockpits
- Chemical-resistant headgear for overhead exposure

NOTE: For containers of over 1 gallon but less than 5 gallons: Persons engaged in pouring this product must wear coveralls or a chemical-resistant apron.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. After each day of use, clothing or PPE must not be reused until it has been cleaned.

Engineering Controls Statements:

For Containers of 5 gallons or more: A mechanical system (such as probe and pump) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal. If the mechanical system is used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4)], the handler PPE requirements may be reduced or modified as specified in the WPS.

When handlers use enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

Wash hands, before eating, drinking, chewing gum, using tobacco or using the toilet.

Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

STATEMENT OF
PRACTICAL TREATMENT

If In Eyes: Immediately flush with plenty of water for 15 minutes. Call a physician.

If Swallowed: Call a physician or poison control center. Induce vomiting immediately by giving two glasses of water and sticking finger down throat. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person. **If On Skin:** Wash with soap and plenty of water. Remove and wash contaminated clothing before reuse. Discard contaminated shoes.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Mixing and Loading: Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

AGRICULTURAL USE REQUIREMENTS

Refer to label booklet for *Agricultural Use Requirements* in the *Directions for Use* section for information regarding the Worker Protection Standard, including Personal Protective Equipment PPE and Agricultural Use Requirements.

NOTICE: Read the entire label. Use only according to label directions. Before buying or using this product, read "Warranty Disclaimer" and "Limitation of Remedies" inside label booklet.

AGRICULTURAL CHEMICAL: Do not ship or store with food, feeds, drugs or clothing.

Distributed by **TERRA** for
GLADES FORMULATING CORPORATION
P.O. Box 1690
Belle Glade, FL 33430-6286

WEED-AMINE

2,4-D HERBICIDE

For Selective Control of Many Broadleaf Weeds in Non-Crop Areas, Grass Pastures, Rangefields and in Certain Crops

Contains Dimethylamine Salt of 2,4-D

ACTIVE INGREDIENTS:

2,4-Dichlorophenoxyacetic acid, dimethylamine salt	46.3%
INERT INGREDIENTS	53.7%
Total	100.0%

2,4-dichlorophenoxyacetic acid, 38.4%, 3.8 lbs. by 100 lbs. of Weed-Amine. The active ingredient is contained in the liquid volume form of 2,4-D and does not release through vapors from treated areas to reduce yield of adjacent susceptible crops.

KEEP OUT OF REACH OF CHILDREN

WARNING

Stirred or shaken for use. Always a suitable form for use as a liquid. Do not use if you do not understand the label, and cannot explain it to you (in Dutch).

Read the label of each bottle for additional precautionary information. For use on lawns, golf courses, cemeteries, parks, airfields, roadsides, vacant lots and drainage ditches. Do not use on lawns, golf courses, cemeteries, parks, airfields, roadsides, vacant lots and drainage ditches. Do not use on lawns, golf courses, cemeteries, parks, airfields, roadsides, vacant lots and drainage ditches. Do not use on lawns, golf courses, cemeteries, parks, airfields, roadsides, vacant lots and drainage ditches.

AGRICULTURAL CHEMICAL - Do not mix or store with food, feed, drug or clothing.

EPA REG. NO. 57242-4 EPA EST. 57242-FL-1

15

application, or other factors, all of which are beyond the control of Glades Formulating Corp. or the seller. All such risks shall be assumed by Buyer.

LIMITATION OF REMEDIES

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to: a) Glades Formulating Corp. shall be liable for the cost of the product bought or purchased; b) Replacement of amount of product used.

Glades Formulating Corp. shall not be liable for losses or damages resulting from handling or use of this product unless Glades Formulating Corp. is promptly notified of such loss or damage in writing. In no case shall Glades Formulating Corp. be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer above and the Limitation of Remedies shall be null and void if any written or verbal statements or representations are made by any agent of Glades Formulating Corp. or the seller, which are not authorized to vary or exceed the terms of the Warranty Disclaimer or the Limitation of Remedies in any respect.

Distributed by TERRA for
GLADES FORMULATING CORPORATION
P.O. Box 1690; Baita Glades, FL 33430-6288

45035 0784

12

WEED CONTROL IN SUGARCANE: See Table for recommended use rates. Apply as a pre-emergence or post-emergence spray in accordance with state recommendations.

WEED CONTROL ON FALLOW LAND: Use 1 to 2 quarts of this product per acre on annual broadleaf weeds and up to 3 quarts per acre on established perennial species, such as Canada thistle and field bindweed. Apply to weeds actively growing. Do not plant any crop for 3 months after treatment or until chemical has dissipated from soil.

RANGELAND, PASTURE, NON-CROP AND FOREST USES

Use Requirements for Rangeland, Pasture, Forest and Non-Crop Areas: No Worker Protection Standard worker entry restrictions or worker notification requirements apply when this product is applied to rangeland, pasture or non-crop land areas, and forest sites when applied by the tree injection method only.

WEED CONTROL IN ESTABLISHED GRASS PASTURES AND RANGELANDS: Use at 2 to 4 pints per acre. Apply preferably when weeds are small and growing actively before the bud stage. Do not use on bentgrass, alfalfa, clover or other legumes. Do not use on newly seeded areas until grass is well established. Do not use from early bud to milk stage where grass seed production is desired. **Grazing Restrictions:** Do not graze lactating dairy animals on treated areas within 7 days after application. Do not harvest grass cut for hay from treated areas for 30 days. Withdraw meat animals from treated areas at least 3 days before slaughter.

CONTROL OF SOUTHERN WILD ROSE: On pastures and rangeland, apply a maximum of 2 quarts of this product per acre per application. On roadsides and lawns, use 1 gallon of this product plus 4 to 8 fluid ounces of an agricultural surfactant per 100 gallons of water and spray thoroughly as soon as foliage is well

15

delayed following treatment. With spring application, reseed in the fall, with fall application, reseed in the spring. Legumes are usually damaged or killed. Deep-rooted perennial weeds such as bindweed and Canada thistle may require repeated applications.

WEED CONTROL IN GRASS SEED CROPS: Use 1 to 4 pints per acre in spring or fall to control broadleaf weeds in grass being grown for seed. Do not apply from early bud to the milk stage. Spray seedling grass only after the first stage, using 3/4 to 1 pint per acre to control small seedling weeds. After the grasses are established, higher rates of up to 4 pints can be used to control hard-to-treat annual or perennial weeds. For best results, apply when soil moisture is adequate for good growth. **NOTE:** Do not harvest any grain or hay from treated areas until 7 days after application. Do not harvest grass cut for hay from treated areas for 30 days. Withdraw meat animals from treated areas at least 3 days before slaughter.

WARRANTY DISCLAIMER

Glades Formulating Corp. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Glades Formulating Corp. makes no other express or implied warranty of merchantability or fitness for a particular purpose, or any other express or implied warranty.

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of

11

under high temperature and high soil moisture conditions. In such situations, use the low rate of 1/2 pint per acre. After application, delay cultivation for 8 to 10 days to allow the corn to overcome any temporary bulliness. **NOTE:** Hybrids vary in tolerance to 2,4-D. Some are easily injured. Spray only varieties known to be tolerant to 2,4-D. Consult the seed company or your Agricultural Experiment Station or Extension Service weed specialist for this information.

WEED CONTROL IN SORGHUM (MILO): See Table for recommended use rates. Treat only after the sorghum is 15 inches high and preferably before it is 15 inches high. Do not treat during the boot, tasseling and early dough stages. Reduce spray drift by keeping the boom and spray nozzles as low as possible. If crop is taller than 8 inches, use deep nozzles to keep the spray off the leaves. Temporary crop injury can be expected under conditions of high soil moisture and high air temperatures. If it is necessary to apply this product under these conditions, use no more than 2/3 pint per acre. **NOTE:** Hybrids vary in tolerance to 2,4-D. Some are easily injured. Spray only varieties known to be tolerant to 2,4-D. Consult the seed company or your Agricultural Experiment Station or Extension Service weed specialist for this information.

WEED CONTROL IN RICE: See Table for recommended use rates. Apply in the late tillering stage of rice development, at the time of first joint development (first to second green ring), usually 6 to 9 weeks after emergence. Do not apply after panicle initiation, after rice internodes exceed 1/2 inch, at early seedling, early panicle, boot flowering or early heading growth stages. **NOTE:** Some rice varieties under certain conditions can be injured by 2,4-D. Therefore, before spraying, consult local Extension Service or University specialists for appropriate rates and timing of 2,4-D sprays.

14

TURF USES

Use Requirements for Turf Including Sod Farms and Grass Seed Crops

Restricted Entry Interval: When used on sod farms or grass seed crops, follow PPE and entry instructions in the "Agricultural Use Requirements" section of this label. For use on other turf areas, do not allow people (other than applicators) or pets on treatment area during application. Do not enter into treated areas until sprays have dried.

Restrictions on Retreatment: Do not apply more than 2 broadleaf applications per year per treatment site.

BROADLEAF WEED CONTROL IN NON-CROPLAND GRASS AREAS (Including Sod Farms or Grass Seed Crops) SUCH AS LAWNS, GOLF COURSES, CEMETERIES, PARKS, AIRFIELDS, ROADSIDES, VACANT LOTS AND DRAINAGE DITCH BANKS: Use 1 to 2 quarts of this product per acre in the amount of water needed for uniform application. Treat when weeds are young and growing well. Do not use on ditches or other herbaceous ground covers. Do not use on creeping grasses such as bent grass except for spot treating nor on freshly seeded turf until grass is well established. Reseeding of lawns should be delayed following treatment. With spring application, reseed in the fall, with fall application, reseed in the spring. Legumes are usually damaged or killed. Deep-rooted perennial weeds such as bindweed and Canada thistle may require repeated applications.

WEED CONTROL IN SOD FARMS: Use 1 to 3 quarts of this product per acre in the amount of water needed for uniform application. Treat when weeds are young and growing well. Usually 2 quarts per acre will provide adequate weed control. Do not use on ditches or other herbaceous ground covers. Do not use on creeping grasses such as bent grass except for spot treating or on freshly seeded turf until grass is well established. Reseeding should be

10

of specific varieties and spray only those known to be resistant to 2,4-D injury. If plants are more than 6 inches tall, use directed spray and keep spray off corn and sorghum foliage. These higher rates may be needed to handle difficult weed problems in certain areas, such as under dry conditions, especially in western areas. However, do not use unless possible crop injury will be acceptable. Consult State Agricultural Experiment Station or Extension Service weed specialist for recommendations or suggestions to fit local conditions.

WEED CONTROL IN SMALL GRAINS NOT UNDERSOED WITH A LEAGUE (Barley, Oats, Rye, Wheat): See Table for recommended use rates. Spray after grain begins to emerge and before the boot stage (usually 4 to 8 inches tall) and weeds are small. Do not apply before the tiller stage nor from early boot through the milk stage. To control large weeds that will interfere with harvest or to suppress perennial weeds, preharvest treatment can be applied when the grain is in the dough stage. Best results will be obtained when soil moisture is adequate for plant growth and weeds are growing well. **NOTE:** Do not graze lactating dairy animals on treated areas within 7 days after application. Do not harvest grain fields cut for hay from treated areas for 30 days. Withdraw meat animals from treated areas at least 3 days before slaughter. Do not feed treated straw to livestock.

WEED CONTROL IN CORN: See Table for recommended use rates. **Preplant (preseed) tillage:** Apply any tillage prior to planting. **Pre-emergence:** Apply to soil anytime after planting but before corn emerges. Do not use on very light, sandy soil. **Emergence:** Apply just as corn plants are breaking ground. **Post-emergence:** Apply to emerging corn when corn is over 8 inches tall. Do not apply to corn until it is 8 inches tall. Do not apply to corn until it is 8 inches tall. Do not apply to corn until it is 8 inches tall. Do not apply to corn until it is 8 inches tall.

13

developed. Two or more treatments may be required. **Grazing Restrictions:** Do not graze lactating dairy animals on treated areas within 7 days after application. Do not harvest grass cut for hay from treated areas for 30 days. Withdraw meat animals from treated areas at least 3 days before slaughter.

SPOT TREATMENT IN NON-CROP AREAS: To control broadleaf weeds in small areas with a hand sprayer, use 1/2 pint of this product in 3 gallons of water and spray to thoroughly wet foliage. **TREE INJECTION TREATMENT:** To control unwanted broadleaf trees such as elm, hickory, oak and sweetgum in forest and other non-crop areas, apply this product by injecting 1 inch of the undiluted product through the bark around the trunk at intervals of 1 to 3 inches between edges of the injection wounds. For harder to control species such as ash, maple and dogwood use 2 ml of undiluted product per injection site. Continuous cuts around the trunk often provide improved control. Also, cuts near the ground level may be more effective than at higher levels. Treatments can be made at any season; however, effectiveness may be less during winter months. Maples should not be treated during the spring sap flow. **Note:** No Worker Protection Standard worker entry restrictions or worker notification requirements apply when this product is directly injected into agricultural plants.

FOR RIGHTS-OF-WAY: For perennial broadleaf weeds and susceptible woody species, use up to 2 gallons of this product per acre. For difficult to control perennial broadleaf weeds and woody species, use up to 2 gallons of the product, and 1 to 4 quarts GARLON 3A herbicide per acre. For ground application: High volume: Apply a total of 100 to 400 gallons per acre. Low volume: Apply a total of 10 to 100 gallons per acre. For helicopter: Apply 1 to 1.5 of 0.10 to 0.30 gallons per acre spray volume.

BEST AVAILABLE COPY

73% Reduction

445

2

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

WARNING **AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)
Injurious to eyes. Harmful if swallowed. May cause skin irritation. Do not get in eyes, on skin, or on clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers, including persons repairing or cleaning equipment, must wear:

- Coveralls or short-sleeved shirt and short pants
- Waterproof gloves
- Chemical-resistant footwear plus socks
- Protective eye wear such as face shield or safety glasses (if on label or label protection recommended). Eye protection is not required for ground or aerial applicators in completely enclosed cabs or cockpits
- Chemical-resistant headgear for overhead exposure

NOTE: For containers of over 1 gallon but less than 5 gallons: Persons engaged in pouring this product must wear coveralls or a chemical-resistant apron.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. After each day of use, clothing or PPE must not be reused until it has been cleaned.

Engineering Controls Statements:

For Containers of 5 gallons or more: A mechanical system (such as probe and pump) must be used for transferring the contents of

or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.
GENERAL: Consult Federal, state or local disposal authorities for approval of alternate procedures.

GENERAL INFORMATION

Weed-Amine is a selective herbicide for control of many broadleaf weeds in non-crop areas, grass pastures, rangelands and in certain crops; also for control of trees by injection.

Generally, the lower dosages given will be satisfactory for young succulent growth of sensitive weed species. For less sensitive species and under conditions where control is more difficult, the higher dosages will be needed. Apply this product during warm weather when weeds are young and growing actively. Use enough spray volume for uniform coverage by ground or air application. If only bands or rows are treated, leaving middles unsprayed, the dosage per crop acre is reduced proportionately. Do not apply where drift may be a problem due to proximity of susceptible crops or other desirable plants. Read and follow all Use Precautions given on this label.

General Use Precautions

Be sure that use of this product conforms to all application regulations. Chemigation: Do not apply this product through any type of irrigation system.

Crop Injury: Do not apply this product directly to, or otherwise permit it to come into contact with cotton, flowers, fruit trees, grapes, ornamentals, vegetables, or other desirable plants which are sensitive to 2,4-D herbicides and do not use in a greenhouse. Do not permit spray mist containing it to drift onto them, since even very small quantities of the spray which may not be visible, can cause severe injury during both growing and dormant periods. Use coarse sprays to minimize drift.

Avoid Spray Drift: Applications should be made only when there is

3 this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal. If the mechanical system is used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

STATEMENT OF PRACTICAL TREATMENT

If In Eyes: Immediately flush with plenty of water for 15 minutes. Call a physician.

If Swallowed: Call a physician or poison control center. Induce vomiting immediately by giving two glasses of water and sticking finger down throat. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person. If On Skin: Wash with soap and plenty of water. Remove and wash contaminated clothing before reuse. Discard contaminated shoes.

no hazard from spray drift since very small quantities of the spray, which may not be visible, may severely injure susceptible crops during both growing and dormant periods. Use coarse sprays to minimize drift since, under adverse weather conditions, fine spray droplets may drift a mile or more. A spray thickening agent, such as Nalco-Trot or Sta-Pul, may be used with this product to aid in reducing spray drift. If used, follow all use recommendations and precautions on the product label.

Drift from aerial or ground application may be reduced by:

1. Applying as near to the target as possible in order to obtain coverage.
2. By increasing the volume of spray mix per acre.
3. By decreasing the pounds of pressure at the nozzle tips.
4. By using nozzles which produce a coarse spray pattern.
5. By not applying when wind is blowing toward susceptible crops or valuable plants.
6. (Aerial) By using straight stream nozzles directed straight back and using a spray boom no longer than 3/4 the wing or rotor span of aircraft.

Determine Air Movement and Directions Before Foliage Application: Use a smoke generator or other means at or near the application site for the detection of air movement, air stability or temperature inversions. Such a condition exists when there is little or no wind and air temperature is lower near the ground than at higher levels. Use appropriate drift control measures or avoid application when smoke is moving toward nearby desirable susceptible plants or sensitive areas.

To avoid injury to desirable plants, do not store, handle, or apply other agricultural chemicals with the same containers or equipment used for this product except as specified on this label. Excessive amounts of 2,4-D in the soil may temporarily inhibit seed germination or plant growth.

Consult your state Agricultural Experiment station or Extension Service weed specialists in many states for recommendations from

4 **ENVIRONMENTAL HAZARDS**

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Mixing and Loading: Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the

8 this label that best fit local conditions. Be sure that use of this product conforms to all applicable regulations. Apply this product only as specified on this label.

WEEDS CONTROLLED

blitterweed	loco, bigbend	shepherdspurse
broomweed	lupines	sicklepod
burdock	mallow, Venice	smartweed
carpetweed	marshelder	sneezeweed, bitter
carrot, wild	morningglory, annual	spittlebug, annual
chicory	mustards	spunkneedles
cocklebur	penycress	sunflower
coffeebean	penworth	sweatclover
croton	pepperweed	tansymustard
dandelion	pigweed	thistle, bull
dock	plantains	thistle, musk
flixweed	poorjoe	thistle, Russian
gallinoga	pusley, Florida	tumbleweed
hemp, wild	radish, wild	velvetleaf
jewelfeed	ragweed, common	vetch
limonweed	rape, wild	waterplantain
kochia	rocket, yellow	witchweed
lambsquarters		wormwood

To Prepare the Spray: Mix this product only with water, unless otherwise directed on this label. Add about half the water to the mixing tank, then add this product with agitation, and finally the rest of the water with continuing agitation. Note: Adding oil, wetting agent or other surfactant to the spray may increase effectiveness on weeds, but also may reduce selectivity to crops resulting in crop damage.

**APPROVED USES
CROP USES**

Agricultural Use Requirements for Crops: For the following crop uses, follow PPE and Roentry instructions in the "Agricultural Use Requirements" section of this label.

5 statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to use of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls over short-sleeved shirt and short pants
- Waterproof gloves
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. STORAGE: Keep container tightly closed when not in use. If exposed to subfreezing temperatures, the product should be warmed to at least 40°F and mixed thoroughly before using.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional office for guidance.

CONTAINER DISPOSAL: Metal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Plastic Containers: 5 Gallons or Less: Triple rinse (or equivalent). Then dispose of in a sanitary landfill.

AMOUNT OF THIS PRODUCT TO USE IN CROPS

By Air or Ground Application

NOTE: Do not apply when weather conditions favor drift from treated areas. Read complete directions and use precautions before using.

CROP	DOSAGE PER ACRE	
	Normal Rates (Usually Safe To Crop)	Higher Rates for Special Situations ¹ (More Likely to Injure Crop)
SMALL GRAINS		
Spring postemergence		
wheat, barley, rye	2 1/2 to 1 1/2 pts.	2 to 3 pts.
oats	1/2 to 1 pt.	1 1/2 to 2 pts.
Preharvest (dough stage)		
wheat, barley, oats	1 to 2 pts.	2 to 3 pts.
CORN¹		
Preemergence	2 to 4 pts.	
Emergence ²	1 pt.	1 1/2 pts.
Postemergence ³		
up to 8 inches tall	1/2 to 1 pt.	
8 inches to tasseling	1 pt.	1 1/2 to 2 1/2 pts.
(use only directed spray)		
SORGHUM (Milo)¹		
Postemergence		
6 to 8 inches tall	2/3 to 1 pt.	
8 to 15 inches tall	1 pt.	1 1/2 to 2 pts.
(use only directed spray)		
RICE	1 to 2 1/2 pts.	2 to 3 pts.
SUGARCANE	2 to 4 pts.	

¹Corn and sorghum varieties vary in tolerance to 2,4-D; some are easily injured. Before spraying, get information on 2,4-D tolerance

BEST AVAILABLE COPY