

PM 23 57542-5 10/6

US ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (TS-767) WASHINGTON, DC 20460	EPA REGISTRATION NO. 57542-5	DATE OF ISSUANCE Jan 31, 1995
	TERM OF ISSUANCE Indefinite	
	NAME OF PESTICIDE PRODUCT [Faint text]	

NOTICE OF PESTICIDE:  REGISTRATION  
 REREGISTRATION  
 (Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

[Faint text]  
 Clades Formulating Corp.  
 c/o Regent Company  
 P.O. Box 2220  
 Stanley, CA 95322-2220

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

Under section 3(c)(7)(A) provided that you:

1. [Faint text]
2. [Faint text]
3. [Faint text]
4. [Faint text]

If the conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 3(d). Your failure to comply with the conditions will constitute a violation of the Act and may result in the suspension or revocation of the registration of your product.

**BEST AVAILABLE COPY**

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

DATE

JAN 31 1995

2-76

# ASULAM HERBICIDE

For Agricultural or Commercial Use Only.  
Not for Use by Homeowners.

For Postemergent Weed Control in Sugarcane, Turf, Ornamentals,  
Christmas Tree Plantings and Non-Cropland

**ACTIVE INGREDIENT:**

Sodium salt of asulam (methyl sulfanylylcarbamate)*	36.2%
<b>INERT INGREDIENTS</b>	<u>63.8%</u>
Total	100.0%

\*Equivalent to 33.1% asulam or not less than 3.34 lbs. per gallon.

**Keep Out of Reach of Children**

## CAUTION

See side panel for additional precautionary statements.

ACCEPTED  
with COMMENTS  
In EPA Letter Dated

JAN 21 1995  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
registered for the pesticide  
EPA Reg. No.  
57242-5

NET CONTENTS: \_\_\_\_\_ GALLONS

EPA Reg. No. 57242-L

EPA Est. 57242-FL-1

GLADES FORMULATING CORPORATION  
P.O. Box 1690; Belle Glade, FL 33430-6286

276

**PRECAUTIONARY STATEMENTS**  
**Hazards to Humans & Domestic Animals**

**CAUTION:** Avoid contact with skin, eyes or clothing.

**Statement of Practical Treatment**

**If On Skin:** Wash with plenty of soap and water. Get medical attention if irritation persists. **If In Eyes:** Flush with water for 15 minutes. Get medical attention if irritation persists.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Applicators and other handlers must wear long-sleeved shirt and long pants, waterproof gloves, and shoes plus socks. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**ENGINEERING CONTROL STATEMENTS**

When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 172.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**USER SAFETY RECOMMENDATIONS**

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

**Environmental Hazards**

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not clean equipment or dispose of equipment washwater in a manner that will contaminate resources. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read entire label before using this product. Do not apply this product through any type of irrigation system. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box apply only to use of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil or water is coveralls, waterproof gloves and shoes plus socks.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

**Storage:** Store at temperatures above 20°F.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal:** Triple rinse or equivalent. Then puncture and dispose of in a sanitary landfill, by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**BEST AVAILABLE COPY**

**Returnable/Refillable Containers:** After use, return the container to the point of purchase or designated locations. This container must be refilled only with Asulam Herbicide. **DO NOT REUSE THE CONTAINER FOR ANY OTHER PURPOSE.** Prior to refilling, inspect thoroughly for damage such as cracks, punctures, abrasions and damaged or worn out threads on closure devices. Do not refill or transport damaged or leaking containers. Check for leaks after refilling and before transportation. For information on the disposal of unused, unwanted product, contact Glades Formulating Corporation at (407) 996-4200. For information on cleanup of spills, contact Rhône-Poulenc at 1-800-334-7577. If the container is not being refilled, return it to the point of purchase.

### SUGARCANE

This product can be applied to either plant cane or cane grown from stubble. Apply as a water mix spray for ground applications. Use 15 to 100 gallons of water per acre, depending on local practice. For aerial application, this product should be mixed in 3 to 5 gallons of water per acre, except in Hawaii, where 5 to 10 gallons of water per acre should be used.

Addition of an adjuvant, cleared for use on growing crops, to the Asulam Herbicide water mix spray will improve weed control when environmental conditions are not optimal. Use either a non-ionic surfactant containing a minimum of 80% active ingredient at the rate of 1 to 2 quarts per 100 gallons (0.25 to 0.5% V/V) of water mix spray or a crop oil concentrate containing 80 to 85% paraffin based petroleum oil and 15 to 20% non-ionic surfactant at the rate of 4 quarts per 100 gallons (1% V/V) of water mix spray.

The rates of Asulam Herbicide given below are for broadcast applications. For banded application, reduce the rate proportionally to the width of the band according to the following formula:

$$\frac{\text{BAND WIDTH (inches)}}{\text{ROW WIDTH (inches)}} \times \text{Broadcast Rate} = \text{Band Rate/Acre}$$

For spot treatments, use a 5% V/V Asulam Herbicide spray (1 gallon per 20 gallons of water). Do not exceed 8 pints per acre per treatment.

### SUGARCANE

#### Single Application Per Growing Season

WEED SPECIES	SPECIAL INSTRUCTIONS	RATE
Itchgrass or Raoulgrass ( <i>Rottboellia exaltata</i> )	Apply when the grass is 8 inches tall or less (addition of surfactant is necessary).	8 pints/Acre
Johnsongrass ( <i>Sorghum halepense</i> )	Apply when the grass is between 12 to 18 inches tall. Johnsongrass should be actively growing and the average air temperature should be at least 60°F or higher.	
Paragrass or Califormagrass ( <i>Brachiaria mutica</i> or <i>Panicum purpurascens</i> )	Apply when the grass is 6 to 8 inches tall or less.	6-8 pints/Acre
Crabgrass ( <i>Digitaria</i> spp.)	If treatment is made before the grass reaches seed head formation then lower rate should be used. If the grass is in early seed head formation then the higher rate should be used.	
Alexandergrass ( <i>Brachiaria plantaginea</i> ) Foxtail ( <i>Setaria</i> spp.) Goosegrass ( <i>Eleusine indica</i> ) Broadleaf Panicum ( <i>Panicum adspersum</i> ) Barnyardgrass ( <i>Echinochloa crusgalli</i> )	If treatment is made when the grass is 6 to 8 inches tall or less, then the lower rate should be used. If the grass is greater than 8 inches tall, then the higher rate should be used.	

### SUGARCANE

#### Two Applications Per Growing Season

This may be required when initial weed infestations are heavy and/or when rhizome Johnsongrass is present. Two applications may also be used when treating weed species which germinate at different times during the growing season.

WEED SPECIES	SPECIAL INSTRUCTIONS	1ST APPLICATION	2ND APPLICATION
Crabgrass ( <i>Digitaria</i> spp.)	At each application the grass should be treated before seed head formation.	6 to 8 pints/Acre	6 to 8 pints/Acre
Itchgrass or Raoulgrass ( <i>Rottboellia exaltata</i> )	At each application to the grass should be 8 inches tall or less (addition of surfactant is necessary).	8 pints/Acre	8 pints/Acre
Johnsongrass ( <i>Sorghum halepense</i> )	At each application the grass should be between 12 and 18 inches tall.	8 pints/Acre	8 pints/Acre

#### RESTRICTIONS AND PRECAUTIONS: Sugarcane

- Asulam Herbicide should be used when the weeds are actively growing.
- Do not plant crops other than sugarcane in treated area. Cover crops may be planted if plowed under and not grazed.
- Do not treat sugarcane within 90 days of harvest.
- Do not graze or feed sugarcane fodder and forage to livestock.

**BEST AVAILABLE COPY**

2/6

- Do not rotate with any crop which is not registered for use with this product for one year following the last application of this chemical.
- Cultivation and/or fertilizer applications or any other cultural practice that disturbs the root system of targeted weed species may result in less than optimum control when applying this product. These practices are not recommended within 7 days prior to or within 7 days after applications of this product.
- Differences in crop tolerance to this product among Sugarcane varieties has been reported in Louisiana. Contact your local County Agent or University Extension Specialist for further information.

**NON-CROPLAND**

This product may be used as postemergent treatment to control weeds on non-cropland areas such as:

- |                                    |  |
|------------------------------------|--|
| Boundary fences                    | Railroad rights-of-way and yards         |
| Fence rows                         | Storage areas and industrial plant sites |
| Highway and roadside rights-of-way | Utility rights-of-way and yards          |
| Lumberyards                        | Warehouse lots                           |
| Pipeline rights-of-way             |  |

A surfactant may be added to the spray solution at 0.25% by volume. Use an approved non-ionic surfactant. Apply this product as a water-mix spray for ground applications using 20 to 100 gallons of solution per acre, depending upon local practice, to control the following weed species:

WEED SPECIES	SPECIAL INSTRUCTIONS	RATE
Crabgrass ( <i>Digitaria</i> spp.)	Apply before the grass reaches seed head formation.	
Johnsongrass ( <i>Sorghum halepense</i> )	Apply when the grass is 18 inches or taller. Use the higher rate in well established heavy infestations. For spot treatment in Hawaii, use the higher rate in 100 gallons of solution and apply an amount not to exceed 50 gallons of total solution per acre.	1 to 2 gallons per Acre
Paragrass or Californiagrass ( <i>Brachiaria mutica</i> or <i>Panicum purpurascens</i> )	Apply before the grass reaches seed head formation. For spot treatment in Hawaii, use the same rate in 100 gallons of solution and apply an amount not to exceed 50 gallons of total solution per acre.	
Western Bracken ( <i>Pteridium aquilinum</i> var. <i>pubescens</i> )	Apply when the fern is in full frond.	7 to 8 pints per Acre

**CHRISTMAS TREE PLANTINGS**

This product may be used as a postemergent treatment in Christmas Tree Plantings where Douglas Fir, Grand Fir, Nobel Fir or Scotch Pine are grown. Do not graze or feed foliage from treated areas to livestock. This product should be applied as a water mix spray. For ground application, use a minimum of 20 gallon of solution per acre; and for aerial application, use a minimum of 10 gallons of solution per acre. Do not use a wetting agent with this product.

WEED SPECIES	SPECIAL INSTRUCTIONS	RATE
Western Bracken ( <i>Pteridium aquilinum</i> var. <i>pubescens</i> )	Apply after bud break and hardening or firming of new tree growth. Bracken should be in full frond prior to treatment.	1 gal./Acre

**TURF**

This product can be applied on St. Augustinegrass and Tifway 419 Bermudagrass turf. Apply one application per season postemergence to the weeds listed below. Use 20 to 50 gallons of water per acre in the spray solution.

TURF SPECIES	WEED SPECIES	RATE
St. Augustinegrass	Bullgrass ( <i>Paspalum supinum</i> ), Crabgrass ( <i>Digitaria</i> spp.), Goosegrass ( <i>Eleusine indica</i> )	5 pints/Acre
Tifway 419 Bermudagrass	Sandbur ( <i>Cenchrus</i> spp.)	

Do not use a surfactant. Do not apply to turf which is under stress or freshly mowed.

**ORNAMENTALS**

This product can be applied as a single, postemergent, broadcast application on the following ornamentals:

JUNIPERS	YEWS
<i>Juniperus andorra</i> <i>Juniperus chinensis</i> <i>Juniperus conferta</i>	<i>Taxus cuspidata</i> <i>Taxus media</i> <i>Podocarpus macrophyllus</i>
<i>Juniperus horizontalis</i> <i>Juniperus litoralis</i> <i>Juniperus sabina</i>	

Treatment should be made with a minimum of 20 gallons of water per acre. Do not use a surfactant.

6976

WEED SPECIES	SPECIAL INSTRUCTIONS	RATE
Barnyardgrass ( <i>Echinochloa crusgali</i> ) Crabgrass ( <i>Digitaria</i> spp.) Fall Panicum ( <i>Panicum dichotomiflorum</i> ) Foxtails ( <i>Setaria</i> spp.) Goosegrass ( <i>Eleusine indica</i> ) Horseweed (marestail) ( <i>Conyza canadensis</i> )	Apply when the weeds are between the stages of early seedling and early seed head formation.	1 gal./Acre

Local conditions may affect the use of this chemical. Consult State Agricultural Extension or Experiment Station weed specialists for specific recommendations for local weed problems and for information on possible lower dosages.

**LIMITED WARRANTY AND DISCLAIMER**

The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions, and (c) that the directions, cautions and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops, and upon reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE, TO MAKE ANY OTHER WARRANTIES, EXPRESS OF IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED AT THE MANUFACTURER'S OPTION TO REPLACEMENT OF OR THE REPAYMENT OF THE PURCHASE PRICE FOR THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL THE MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

**NOTICE TO BUYER**

Purchase of this material does not confer any rights under patents governing this product or the use thereof in countries outside of the United States.

**BEST AVAILABLE COPY**