

57538-18

01/11/2002

1/14



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Biopesticides and Pollution Prevention Division (7511C)
1200 Pennsylvania Avenue NW
Washington, DC 20460

EPA Reg. Number:

57538-18

Date of Issuance:

1/11/02

Term of Issuance:

Unconditional

Name of Pesticide Product:

PRO GA₃

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Stoller Enterprises
8580 Katy Freeway, Suite 200
Houston, TX 77024

~~Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Biopesticides and Pollution Prevention Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.~~

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA Sec. 3(c)(5) provided you:

1. Submit and /or cite all data required for registration/reregistration of your product under FIFRA Sec.3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA Sec.4.
2. Revise the EPA Registration Number to read, "EPA Reg. No. 57538-18".
3. Submit three(3) copies of the revised final printed label for the record.

Signature of Approving Official:

JZA


Janet L. Andersen, Director,
Biopesticides and Pollution Prevention Division

Date:

1/11/02

Concur
7511C
Rull 1/10/02

2/14



STOLLER

ProGA₃

ACTIVE INGREDIENTS:

Gibberellic acid (GA₃)..... 4.0%

OTHER INGREDIENTS:..... 96.0%

Total 100.0%

This product contains approximately 1.0 gram active ingredient per fluid ounce.

**KEEP OUT OF REACH OF CHILDREN
 WARNING-AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

FIRST AID	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.</p>	

See additional Precautionary Statements on side panel.

EPA Reg. No. 57538-

**Manufactured By
 STOLLER ENTERPRISES, INC.
 8580 Katy Freeway, Suite 200
 Houston, Texas 77024
 Phone (713) 461-1493**

NET CONTENTS ___ GAL. (___ L)

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

WARNING

Causes substantial but temporary eye injury. Harmful if inhaled or absorbed through skin. Do not get in eyes or on clothing. Avoid breathing vapor or spray mist, and avoid contact with skin. Wear goggles or safety glasses. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category C on an EPA chemical-resistant category selection chart.

Applicators and other handlers must wear: long-sleeved shirt and long pants, chemical-resistant gloves, such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride and viton, shoes plus socks, and protective eyewear.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwater. Exposed treated seed may be hazardous to birds and other wildlife. Dispose of all excess treated seed and seed packaging by burial away from bodies of water.

PHYSICAL OR CHEMICAL HAZARDS

Flammable. Keep away from heat and open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exception: If the product is soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls over short-sleeved shirt and short pants, chemical-resistant gloves, such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride and viton, shoes plus socks, and protective eyewear.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Keep containers tightly closed when not in use. Keep away from heat and open flame.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Do not reuse empty containers. Triple rinse or equivalent. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by the state authorities, by burning. If burned, stay out of smoke.

APPLICATION RECOMMENDATIONS

Use only as directed. The label should be read thoroughly and understood before making applications.

ProGA₃ contains gibberellic acid which is an extremely potent plant growth regulator. When applying plant growth regulators, deviations from the label directions in the rates, timings, water volumes, or the adoption of untested spray mixes, may result in undesirable effects. Always consult the State Extension Service Specialist in your area for the spray regimen best suited to your conditions.

- Do not apply to plants under pest, nutritional or water stress.
- When a range of rates is indicated, use the concentration and spray volume recommended locally by the State Extension Service Specialist.
- For optimum effectiveness, thorough spray coverage must be achieved. All parts of the plant or crop must receive the spray or desired results will not occur. Prepare solution concentrations by mixing the required amount of product with water in a clean, empty spray tank. Discard any unused spray material at the end of each day following local, state or Federal Law.

- For best results, the water pH should be around neutral and always below 8.5.
- ProGA₃ applications made under slow drying conditions (cool to warm temperatures, medium to high relative humidity and no wind) will increase absorption by the plant, thus optimizing effectiveness. Night-time applications are encouraged when day-time conditions are not conducive to slow drying conditions.
- Product persistence: ProGA₃ should be reapplied if significant rain occurs within 2 hours of application.
- Compatibility: The ProGA₃ spray guidelines refer to the use of the product alone. The use of surfactants and other additives has been reported to be beneficial. Stoller does not assume responsibility for unexpected results due to the tank mixing of ProGA₃ with other products.
- DO NOT apply using ULV application methods. For aerial applications, spray volumes must be greater than 2 gallons per acre (10 gallons per acre for tree crops).
- ProGA₃ can be applied up to 7 days before harvest.

SPRAY GUIDELINES FOR GRAPES

ProGA₃ contains approximately 1.0 gram ai/fl. oz.

For all grapes, application is recommended by ground sprayer. Apply in sufficient water volume to insure thorough wetting. It is important to wet all flower clusters or berries thoroughly. For specific spray rates and timings, by variety, see accompanying table. Do not exceed maximum rates.

SEEDLESS GRAPE

Do not apply more than 208 grams ai/acre per growing season for all uses.

For cluster elongation and looser cluster forms (“Stretch”). To reduce costs of thinning, allow better air circulation to aid in the control of bunch rot, and increase light penetration to aid in sugar development.

Guide: Apply one to two applications before bloom when flower clusters are 2 to 5 inches long.

For decreased berry set (“Thinning”), reduced hand-thinning costs, and hastened maturity. **Guide:** Apply one to three applications during bloom. When the bloom period is extended, subsequent sprays should be made 1 to 7 days after the first application.

Note: Higher amounts or multiple applications may cause an excess of shot berries or over-thinning, especially in young vines or vines with high vigor.

For larger berries (“Sizing”) and larger clusters when used in conjunction with established girdling and thinning practices. **Guide:** Apply one to three applications beginning when the average berry size reaches “target” diameter (see Table 1). Timing of the subsequent sprays will be dictated by experience in the vineyard and temperatures occurring between sprays. Potential effect will be reduced if the final spray occurs more than 2 weeks after the first application.

Table I: Application Rates (Grams A.I./Acre) for Seedless Grape, Including Target Berry Diameters

Seedless Grapes	Stretch Grams ai/Acre	Thinning Grams ai/Acre	Sizing	
			Target	Grams ai/Acre
Perlette	8-16	*	4-5 mm	32-80
Flame	8-16	3-16	6-9 mm	20-80
Thompson	8-16	8-16	3-5 mm	32-80
Raisin	8-16	3-12	4-5 mm	4-12
All Others	*	*	12-14 mm	8-48

*No recommendations available for this variety/timing at this time.

Note: Do not apply more than 208 grams ai/acre per growing season for all uses.

SEEDED GRAPE

EMPEROR GRAPE

For reducing berry shrivel. This use can also increase berry size. **Guide:** Apply 20 grams ai/acre as one application approximately 2 weeks after completion of berry shatter. This timing should correspond to a period when the predominant berry diameter ranges from 10 to 15 mm.

BLACK CORINTH (ZANTE CURRANT) GRAPE

For improving berry size: **Guide:** Apply 1 to 8 grams ai/acre as one application 3 to 5 days after full bloom but before shatter begins.

SPRAY GUIDELINES FOR CITRUS

ProGA₃ contains approximately 1.0 gram ai/fl. oz.

For all citrus, apply in sprays of sufficient water volumes to insure thorough fruit wetting. Application to trees of low vigor or under stress (pest, nutritional or water, etc.) may cause severe leaf and/or fruit drop. In most cases some drop of older mature leaves will occur after application. Do not apply in white wash sprays in which lime or other caustic material has produced a high pH in the spray tank.

NAVEL ORANGE

To delay rind aging and reduce rind disorders (e.g., rind staining, water spotting, sticky or tacky surface, puffy rind and rupture under pressure) and to produce a more orderly harvesting pattern: The delay in rind aging is greatest when an early spray is applied. This spray timing produces the firmest rind possible. **Guide:** Apply 16 to 48 grams ai/acre in sufficient water volume to insure thorough wetting.

Early Spray: Apply one spray approximately 2 weeks prior to color break, which normally occurs August through November.

OR

Late Spray: Apply one spray after marketable color has developed, normally from October through December. This late application may cause fruit re-greening.

Note: Do not apply the early spray to groves that may be harvested early, as fruit coloring will be delayed. Do not apply from January through July, as production may be reduced the following year.

VALENCIA ORANGE

To reduce rind creasing and to delay rind aging and softening: **Guide:** Apply a single spray in August to October to trees with a target crop of young fruit. Apply 40 to 80 grams ai/acre in sufficient water volumes to insure thorough wetting.

Note: Slower color development should be expected in the target crop. Increased re-greening of mature fruit may occur. After marketable color is achieved, treatment effects may be reduced the longer treated fruit remains on the tree.

OTHER ORANGES (All States Except California)

To reduce rind creasing and to delay aging and softening of the rind: **Guide:** Apply a single spray in August to October to trees with a target crop of young fruit. Apply 40 to 80 grams ai/acre in sufficient water volume to insure thorough wetting.

Note: Slower color development should be expected in the target crop. Increased re-greening of mature fruit may occur. After marketable color is achieved, treatment may be reduced the longer treated fruit remains on the tree.

LEMON/LIME

To decrease the amount of small ripe fruit and to produce a more desirable production pattern relative to market demand. **Guide:** Apply one spray when target crop is 1/2 to 3/4 full size, but still green. Use 10 to 32 grams ai/acre in sufficient water volume to insure thorough wetting. When applied 2 years in a row, an even larger difference in harvest pattern and maturity occurs.

TANGERINE HYBRIDS

To delay disorders associated with rind aging, puffiness, and softening, and to increase peel strength of tangerine hybrids such as Orlando, Robinson, Minneola and Sunburs.: **Guide:** Apply 20 to 40 grams ai/acre approximately 2 weeks prior to color break. Apply as a dilute spray in sufficient water volume to insure thorough wetting.

Note: Do not apply if early harvest is planned. Do not apply after coloring as preharvest rind staining may occur. Application during coloring may cause variation in rind color development.

(All States Except California)

To increase fruit set and yields as tangerine hybrids with pollination problems such as the Orlando, Robinson, Minneola and Sunburst: **Guide:** Apply 8 to 30 grams ai/acre during full bloom. Apply as a dilute spray in sufficient water volume to insure through wetting.

Note: Fruit sizes may be reduced and color development slightly retarded. A slight increase in mature leaf drop may occur in trees under stress.

GRAPEFRUIT (All States Except California)

To delay disorders associated with rind aging (e.g., puffiness, softening and orange coloration), to prevent preharvest drop of mature fruit, to increase peel strength and reduce water loss during storage, and to produce a more orderly harvesting pattern. The delay in rind aging is greatest when an early spray is applied before color change. This spray timing produces the firmest rind possible. **Guide:** Apply 16 to 48 grams ai/acre in a minimum of 250 gallons per acre.

Early Spray: Apply one spray approximately 2 weeks prior to color break, which normally occurs August through September.

OR

Late Spray: Apply one spray after marketable color has developed which is normally from October through December. This late application may cause fruit re-greening.

Note: Do not apply the early spray to groves that may be harvested early as fruit coloring will be delayed. It is advisable to spot pick heavy crops to aid early marketing and to avoid reduction of yields, which generally follow late held crops. Applications made to fully colored fruit will begin to re-green if allowed to remain on the tree for extended periods.

Applications made after December, or when trees begin to break dormancy may adversely affect new crop. Do not use concentrate sprays. Results may vary from season to season depending on environmental conditions.

STAR RUBY GRAPEFRUIT (All States Except California)

To reduce early-season small fruit drop of Star Ruby Variety thereby increasing yields. **Guide:** Apply a single spray during the bloom period. Use 25 grams ai/acre in a minimum of 250 gallons of water per acre.

Note: Results may vary from season to season depending on environmental conditions. Maintain a well-balanced fertilization and watering program.

SPRAY GUIDELINES FOR FRUIT CROPS

ProGA₃ contains approximately 1.0 gram ai/fl. oz.

BLUEBERRY (All States Except California)

To improve fruit set. For natural fruit set problems due to insufficient natural honeybee pollination, adverse weather conditions, or physiological factors.

Highbush Blueberry (for varieties such as Coville, Jersey, Stanley, Earliblue, Weymouth, Walcott, Berkeley, Blue-ray, Bluecrop, 1316A, Concord and others)

Guide: Make one or two applications at 40 grams ai/acre in 100 gallons of water. The single application should be made at full bloom (when 75% of the flowers are fully open). When two applications are made, spray the first one at full bloom, and the second one within 10 to 14 days of the first one. For Weymouth, application can be delayed up to 2 weeks after bloom to increase size of "shot" berries.

Rabbiteye Blueberry (for varieties such as Aliceblue, Beckyblue, Bonita, Brightwell, Climax, Delite, Tiftblue, Woodward and others)

Guide: Make a single application of 40 grams ai/acre in 100 to 300 gallons per acre when most of the flowers are elongated but not yet open (bloom Stage 5).

OR

For multiple applications make two to four applications 10 to 14 days apart starting at bloom Stage 5. Spray 40 grams ai/acre in 50 to 300 gallon per application.

SWEET CHERRY

To produce larger, brighter colored, firmer fruit. **Guide:** Apply a single spray when the fruit is light green to straw colored. Use 16 to 48 grams ai/acre in sufficient water volume to insure thorough wetting.

Note: Color development and harvest may be slightly delayed.

RED TART CHERRY (All States Except California)

To maintain and extend high fruiting capacity of bearing tart cherry trees and reduce the occurrence of "blind" nodes. Treatment will cause bud differentiation, which is apparent the year after application. Therefore, changes in shoot, spur and flower production will not be evident until 2 or 3 years after program initiation. Applications must be applied annually to insure vegetative development and subsequent yield improvement year after year. **Guide:** Apply one spray 14 to 28 days after bloom. Optimum timing is defined as that stage when three to five terminal leaves have fully expanded, or, at least 1 to 3 inches of terminal shoot extension has occurred. Use 4 to 18 grams ai/acre, depending on tree age and vigor (see Table 2). Apply in sufficient water volume to insure thorough wetting.

Table 2: Recommended Application Rates (grams ai/acre) for Tart Cherry Trees by Age

Tree Age (years)	Rate (grams ai/acre)
6-10	4-6
11-15	8-10
16-20	10-14
20+	14-18

Note: Rates are based on expected normal tree vigor at various ages. Adjust rate according to tree vigor. If trees are vigorous, use lowest recommended rates. Lowest rates should also be used on trees that have been heavily pruned or hedged. Use higher rates for trees low in vigor and weak in shoot and spur production. Excessive application rates will increase vegetative growth at the expense of fruit production the following year.

Applications will not improve growth of trees under stress conditions, such as nutritional, moisture, or pest. Best results will be obtained when combined with good cultural practices.

SPRAY GUIDELINES FOR NON-BEARING YOUNG TART AND SWEET CHERRY TREES (All States Except California)

To reduce flowering and fruiting in young tart and sweet cherry trees in order to minimize the competitive effect of early fruiting on tree development.

Note: DO NOT SPRAY TREES IN THE FIRST YEAR. Treat in the second season for reduction of flowering in the third season, and again in the third season if flower reduction and fruiting is desired in the fourth season. Treat only trees that are in good physiological condition. Discontinue treatment the year before desired harvest.

Guide: Apply a single spray of 20 to 40 grams ai/acre, 2 to 4 weeks after bloom. Apply a foliar spray of 25 to 50 gallons per acre, assuming a tree density of 100 trees per acre equivalent. Under conditions of

low vigor, two applications are recommended. If two spray applications are made, allow at least a 7 day interval between sprays.

SPRAY GUIDELINES FOR OTHER FRUIT

ProGA₃ contains approximately 1.0 gram ai/fl. oz.

OLYMPUS STRAWBERRY (All States Except California)

To increase runner production of mother plants of the Olympus cultivar: **Guide:** Apply a single spray of 20 grams ai/acre to mother plants 10 to 30 days after planting. At the time of spraying, plants should have one to six leaves. Apply 100 gallons per acre to thoroughly wet new foliage to the point of run-off.

Note: Not for use on fruiting plants. Treatments may not be effective on plantings set out after mid-May.

PINEAPPLE

Apply a single spray of 4 grams ai/acre (10 grams ai/hectare) at the start of flowering.

SPRAY GUIDELINES FOR VEGETABLE CROPS

ProGA₃ contains approximately 1.0 gram ai/fl. oz.

RHUBARB

To break dormancy on plants receiving insufficient chilling and to increase marketable yield of forced rhubarb. **Guide:** Make a single application of 2 fluid ounces (60 ml) of solution (containing 20 grams ai in 10 gallons of water) to each cleaned crown, when the rest period is not completely broken. When the rest period is broken by cold weather, apply 2 fluid ounces (60 ml) of a solution containing 10 grams ai in 10 gallons of water to each cleaned crown.

Note: Keep forcing house temperatures at 40°F to 50°F for 24 hours after application. If house is warmer than 50°F, the crowns should be covered with plastic. Temperatures in the forcing house above 50°F may lower yields and cause poor stalk color.

ARTICHOKE

To accelerate maturity and shift harvest to an earlier date: **Guide:** For perennials, apply one to three applications at bud initiation stage. For annuals, apply one to four applications at 2 week intervals, beginning at the fourth true leaf. Use 10 to 20 grams ai/acre per application in sufficient water volume to insure thorough wetting of the entire plant (leaves, stems and buds).

CELERY

To increase plant height and yield and overcome stress due to cold weather conditions or saline soils, and to obtain earlier maturity: **Guide:** Apply a single spray 1 to 4 weeks prior to harvest at a rate of 2.5 to 10 grams ai/acre. Use 25 to 50 gallons per acre by ground application or 5 to 10 gallons per acre for aerial application*. Use lower concentrations applying 3 to 4 weeks before harvest and higher concentrations within 1 to 2 weeks before harvest.

*Do not apply by air in California.

Note: Do not apply earlier than 4 weeks before harvest as bolting (seed stalk formation) may occur.

LETTUCE FOR SEED

To obtain uniform bolting and increase seed production. **Guide:** Apply one to four applications at 2-week intervals, beginning at the fourth true leaf. Use 1 to 4 grams ai/acre per application in sufficient water volume to insure thorough wetting.

MELON AND CUCUMBER (All States Except California)

To stimulate fruit set during periods of cool temperatures. **Guide:** Use 2 grams ai/acre in sufficient water volume for thorough coverage of exposed foliage. Make one application prior to bloom followed by two additional applications at intervals of 10 to 14 days on cantaloupes and watermelons. On cucumbers, up to four applications may be required.

For maximum benefits, vines must be in good condition, except for reduced rate of growth due to cool temperatures.

PEPPER (All States Except California)

To promote plant growth: **Guide:** Apply one to two sprays of 1 to 3 grams ai/acre in 25 to 50 gallons per acre at 2 week intervals. Begin sprays 2 weeks after transplanting.

Note: This use is recommended for areas with short growing seasons, or when low temperatures slow plant growth.

To increase fruit set and promote fruit growth: **Guide:** Apply one to two sprays of 1 to 3 grams ai/acre at the beginning of the picking period. The high rate is recommended for plants with heavy fruit loads.

SEED POTATO

To stimulate uniform sprouting to aid in maximum production, more uniform development, fewer late maturing plants, and to break dormancy of newly harvested potatoes that have not had a full rest period. **Guide:** Dip whole or cut seed pieces in a solution containing 0.2 to 0.4 grams ai/acre in 100 gallons of water prior to planting.

Note: Under high soil temperatures use the minimum concentration for dormant seed. Do not treat rested seed.

SPINACH (All States Except California)

To facilitate harvest, increase yield and improve quality of fall and over-winter spinach. **Guide:** Apply a single spray 10 to 18 days before each anticipated harvest on fall or over-winter spinach, ideally when daytime temperatures are 40°F to 70°F and during early morning hours when dew is present on crop. Apply 6 to 8 grams ai/acre in 10 to 50 gallons per acre by ground sprayer or in a minimum of 5 to 10 gallons per acre by air. When applied to promote growth of second cutting, wait until some regrowth has started before spraying. Maximum benefit is obtained when below normal temperatures predominate following application and growth would be otherwise slowed in untreated spinach.

Note: Since the promotion of bolting may occur, do not apply after the mid-winter period or if temperatures may be expected to exceed 75°F within several days of application. Do not apply on spring plantings.

SPRAY GUIDELINES FOR OTHER CROPS

ProGA₃ contains approximately 1.0 gram ai/fl. oz.

COTTON (All States Except California)

To promote early plant growth, increase early seedling vigor, and to overcome stress caused by cool weather. **Guide:** Apply 1 to 3 grams ai/acre as a foliar application for the 2-leaf stage through the 5-leaf stage. Make one to two applications as needed. Use 5 to 40 gallons of water by ground application or 3 to 10 gallons by air.

Note: Use higher rates when temperatures will likely average 75°F or less during the 14 days following the application. Do not tank mix with herbicides. Do not apply more often than necessary to achieve the desired height, as overdosage may result in excessive growth.

HOPS (Northwestern U.S. Only)

For seeded and seedless Fuggle hop and similar varieties adapted to the Northwestern states. To increase yield and fruit set. **Guide:** Apply a single spray when vine growth is 5 to 8 feet in length. Use 4 to 6 grams ai/acre in 100 to 150 gallons/acre.

RICE SEED TREATMENT (Except California)

For use as a seed treatment on both semi-dwarf and tall rice varieties to promote germination, emergence and final stand densities when planted at greater depths where soil moisture levels are more adequate for germination. ProGA₃ is particularly effective on semi-dwarf varieties such as Lemont, Gulfmont, and Texmont. This will also result in more uniform emergence thus allowing more accurate and efficient herbicide, fertilizer, fungicide and insecticide applications and may maximize yield and improve grain quality.

Apply only to rice seed intended for drill seeded or dry broadcast systems. Do not apply to rice used in a 24-hour presoak prior to broadcast. Do not use more than 2 grams ai per 100 pounds of seed. **DO NOT USE TREATED SEED FOR FOOD, FEED, OR OIL PURPOSES.**

Guide: Use 1 to 2 grams ai in 8 to 20 oz. water per 100 pounds of rice seed. ProGA₃ can be applied to dry seed with standard mist-treating equipment. Best results are obtained using a higher treatment volume (12 to 20 oz./cwt. of seed) to insure the seed is completely and uniformly covered with ProGA₃. Fill the seed treatment tank with water to one-half the final tank mix volume. Add the required amount of ProGA₃, mixing thoroughly while adding water and other seed treatment products to the desired final volume.

An approved dye must be added to distinguish treated seed and prevent inadvertent use for food, feed or oil purposes. Seed commercially treated with this product must be labeled in accordance with all applicable requirements of the federal and state seed laws. ProGA₃ is compatible with most commonly used fungicide seed treatments such as Vitavax® and Dithane®, standard dyes and sticker-binding agents. When preparing tank mixes, the user should ensure adequate physical compatibility and mixing characteristics.

RICE POSTEMERGENCE SEEDLING TREATMENT (All States Except California)

For use as a postemergence seedling application on rice grown in the U.S. to promote more uniform and vigorous growth prior to permanent flooding. Early season foliar applications of ProGA₃ will promote vigorous and more uniform seedling growth of rice prior to permanent flood establishment. This will allow earlier (five to ten days) flooding of drill or dry broadcast seeded varieties and is particularly effective on semi-dwarf varieties. Early flooding may reduce additional flushing costs associated with a delay in permanent flooding, weed infestations and the number of herbicide applications as well as promote earlier and more uniform grain maturity. ProGA₃ application may result in a temporary lighter green foliage color due to accelerated growth rates.

Avoid drift or accidental application to other crops. Do not apply when rice is subject to drought stress conditions. ProGA₃ can be tank mixed with most commonly used rice herbicides and fungicides. When ProGA₃ is applied in tank mixes with Arrosolo®, Stam® M4 combined with labeled herbicides, Stam® 80EDF or Wham® EZ, plus a recommended adjuvant, the use of a surfactant is not necessary. Do not apply with Whip®.

ProGA₃ applied between split-boot and 100% heading can increase height of semi-dwarf rice. This may facilitate harvest efficiency in the field by allowing the rice grain to be cut above the leaf canopy at faster combine speeds and at reduced vegetative load. Grain quality and maturity can be advanced with the promotion of tiller panicle development. Heading applications to the first crop can also accelerate regrowth of second crop rice. This can result in earlier second crop maturity and maximize grain yield.

SEEDLING APPLICATIONS

ProGA₃ may be applied at a rate of 1 to 3 grams ai/acre to rice between the 1 to 2 leaf stage and the 4 to 5 leaf stage of growth. Timing and dosage is based on environmental conditions, tank mix combinations with herbicides, and preferred permanent flood practice in relation to rice leaf stage.

For best results, apply ProGA₃ at a rate of 1 to 2 grams ai/acre using a non-ionic surfactant such as Golden Pest Spray Oil or in tank mix combination with rice herbicides. Use higher rates (1.5 to 3 grams ai/acre) with some dry and water based herbicide formulations, or when temperatures will likely average 75°F or less during fourteen days after application.

PANICLE EXTENSION APPLICATIONS

Apply 3 to 8 grams ai/acre between split-boot and 100% panicle heading to promote main culm and tiller panicle extension. Applications should be made with fixed wing aircraft at 10 gallons spray volume per acre. Tank mixing with Natur'l Oil spray adjuvant is recommended.

APPLICATION EQUIPMENT

Apply ProGA₃ by fixed wing aircraft equipped with spray systems capable of producing a uniform medium to fine spray droplet pattern. Do not apply less than 10 gallons total spray volume per acre. Low pressure ground sprayers equipped with boom and flat fan nozzles and applying 10 to 15 gallons total spray volume per acre may be used.

CONVERSION TABLE

ProGA₃ is a liquid and each fluid ounce contains approximately one (1) gram of active ingredient.

Amount Active per Acre	To	Amount of ProGA ₃ per Acre
0.5 g	=	0.5 oz.
1 g	=	1 oz.
2 g	=	2 oz.
4 g	=	4 oz.
5 g	=	5 oz.
8 g	=	8 oz.
10 g	=	10 oz.

NOTICE: The information contained herein is true and accurate to the best of our knowledge and belief, but it is presented without guarantee since field conditions and use are beyond our control. Neither the manufacturers nor the seller make any warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use of this material when such use is contrary to label instructions. Read and follow the label directions carefully.