

N-LARGETM

Plant Growth Regulator Solution

ACTIVE INGREDIENT:

Gibberellic acid (GA₃)..... 4.0%

OTHER INGREDIENTS:..... **96.0%**

Total 100.0%

This product contains approximately 1.0 gram active ingredient per fluid ounce (30mL).

EPA Reg. No. 57538-18 EPA Est. Nos. 57538-TX-1, 57538-TX-2

INDEX:

- 1.0 First Aid
- 2.0 Precautionary Statements
 - 2.1 Hazards to Humans and Domestic Animals
 - 2.2 Personal Protective Equipment (PPE)
 - 2.3 User Safety Recommendations
 - 2.4 Environmental Hazards
 - 2.5 Physical or Chemical Hazards
- 3.0 Directions for Use
- 4.0 Agricultural Use Requirements
- 5.0 Non-Agricultural Use Requirements
- 6.0 General Instructions for Use
 - 6.1 Application Instructions
- 7.0 Spray Guidelines for Grapes
 - 7.1 Seedless Table Grape
 - 7.2 Seeded Grape
- 8.0 Spray Guidelines for Citrus
 - 8.1 Citrus: Field Applications
 - 8.2 Citrus: Postharvest Applications
- 9.0 Spray Guidelines for Fruit Crops
- 10.0 Spray Guidelines for Non-Bearing Fruit Trees
- 11.0 Spray Guidelines for Vegetable Crops
- 12.0 Spray Guidelines for Other Crops
- 13.0 Spray Guidelines for Ornamentals, Cut Flowers, Turfgrass, Bedding Plants, etc.
- 14.0 Conversion Table (g/fl. oz.)
- 15.0 Conversion Table (ppm)
- 16.0 Storage and Disposal
- 17.0 Warranty

ACCEPTED
FEB 17 2005
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 as amended, for the pesticide
 registered under
 EPA Reg. No. 57538-18

KEEP OUT OF REACH OF CHILDREN

CAUTION

1.0 FIRST AID	
If swallowed	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If inhaled	Move person to fresh air. If person is not breathing, call 911 or an ambulance; then give artificial respiration, by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
If on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-539-5283 for emergency medical treatment information.

2.0 PRECAUTIONARY STATEMENTS

2.1 Hazards To Humans And Domestic Animals

Caution. Harmful if inhaled, swallowed, or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove and wash contaminated clothing before reuse. Wear the appropriate Personal Protective Equipment (PPE).

2.2 Personal Protective Equipment

Mixers, loaders, applicators and other handlers must wear:

- long-sleeved shirt and long pants,
- waterproof gloves, and
- shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

2.3 User Safety Recommendations

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

2.4 Environmental Hazards

For terrestrial uses: Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwater or rinsate. Exposed treated seed may be hazardous to birds and other wildlife. Dispose of all excess treated seed and seed packaging by burial away from bodies of water.

2.5 Physical or Chemical Hazards

Flammable! Keep away from heat and open flame.

3.0 DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide application.

4.0 AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours unless wearing appropriate PPE.

Exception: If the product is soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves anything that has been treated such as plants, soil, or water is: Coveralls, waterproof gloves, and shoes plus socks.

5.0 NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses. Do not enter without appropriate protective clothing until sprays have dried.

Manufactured by:
Stoller Enterprises, Inc.
 4001 W Sam Houston Pkwy N, Suite 100
 Houston, Texas 77043
 Questions? Phone (713) 461-1493
 NET CONTENTS (____ Gals.) (____ L)

6.0 GENERAL INSTRUCTIONS FOR USE

Use only as directed. The label should be read thoroughly and understood before making applications. Do not apply this product through any type of irrigation system.

6.1 Application Instructions

N-LARGE™ contains gibberellic acid, which is an extremely potent plant growth regulator. When applying plant growth regulators, deviations from the label directions in the rates, timings, water volumes, or the adoption of untested spray mixes, will result in undesirable effects. Always consult the State Extension Service Specialist in your area for the spray regimen best suited to your conditions.

- Do not apply to plants under pest, nutritional or water stress.
- When a range of rates is indicated, use the concentration and spray volume recommended locally by the State Extension Service Specialist.
- For optimum effectiveness, thorough spray coverage must be achieved. All parts of the plant or crop must receive the spray or desired results will not occur. Prepare solution concentrations by mixing the required amount of product with water in a clean, empty spray tank. Discard any unused spray material at the end of each day following local, state or Federal Law.
- For best results, the water pH must be around neutral and always below 8.5.
- N-LARGE™ applications made under slow drying conditions (cool to warm temperatures, medium to high relative humidity and no wind) will increase absorption by the plant, thus optimizing effectiveness. Night-time applications are encouraged when day-time conditions are not conducive to slow drying conditions.
- Product persistence: N-LARGE™ must be reapplied if significant rain occurs within 2 hours of application.
- Compatibility: Except when noted elsewhere, the N-LARGE™ spray guidelines refer to the use of the product alone. The use of surfactants and other additives has been reported to be beneficial. Stoller Enterprises does not assume responsibility for unexpected results due to the tank mixing of N-LARGE™ with other products. Conduct a jar test before tank mixing to ensure compatibility of N-LARGE with other pesticides or products if the tank mix combination has not been used previously. To test for compatibility, use a small jar and mix a small amount of spray, combining all ingredients in the same ratio as the anticipated use. If any indications of physical incompatibility develop, do not use this mixture for spraying. Indications of incompatibility usually will appear within 5 to 15 minutes after mixing. To ensure maximum crop safety and product performance, follow all precautions and limitations on this label and labels of products used in the tank mixture with N-LARGE.
- DO NOT apply using ULV application methods. For aerial applications, spray volumes must be greater than 2 gallons per acre (20 l/ha), 10 gallons per acre for tree crops (100 l/ha).
- No harvest interval is required for this product. Observe the 4-hr. REI.

7.0 SPRAY GUIDELINES FOR GRAPES

For all grapes, application shall be by ground sprayer. Apply as a concentrate or dilute spray in sufficient water volume to ensure thorough wetting. It is important to wet all flower clusters or berries thoroughly. For cultivar specific spray rates and timings, see accompanying tables.

7.1 SEEDLESS TABLE GRAPE CLUSTER STRETCH SPRAYS

Objective/benefit	Application timing/instructions
For cluster elongation and looser cluster forms. To reduce costs of thinning, allow better air circulation to aid in the control of bunch rot, and increase light penetration to aid in sugar development.	Make one to three applications before bloom when flower clusters are 2 to 7 inches long.
Crop/Cultivar	Rate (grams a.i./acre)
Perlette Seedless	8-24
Flame Seedless	
Thompson Seedless	
Raisin	
Other Seedless Grapes	

BERRY THINNING SPRAYS

Objective/benefit	Application timing/instructions
For decreased berry set, reduced hand-thinning costs, and hastened maturity.	Make one to four applications during bloom. Only 1-2 applications for "Other Seedless Grapes." When the bloom period is extended, subsequent sprays are to be made 1 to 7 days

Crop/Cultivar	after the first application. Rate (grams a.i./acre)
Perlette Seedless	Not applicable
Flame Seedless	3-16
Thompson Seedless	8-20
Raisin	3-12
Other Seedless Grapes	0.5-12

NOTE: Higher amounts or multiple applications will cause an excess of shot berries or overthinning, especially in young vines or vines with high vigor. For "Other Seedless Grapes" use caution as some of the new cultivars are very responsive and will over-thin easily. A grower shall consult the local specialist before thinning cultivars with which he has no familiarity.

BUMP SPRAY - For Thompson Seedless

Objective/benefit	Application timing/instructions
To help initiate the beginning of the berry growth period.	Make one application of 16-24 grams a.i./acre during the period between the last thinning spray and the first sizing spray.

BERRY SIZING SPRAYS

Objective/benefit	Application timing/instructions
For larger berries and larger clusters when used in conjunction with established girdling and thinning practices.	Make one to four applications beginning when the average berry size reaches "target" diameter (see below). Timing of the subsequent sprays will be dictated by experience in the vineyard and temperatures occurring between sprays. Sprays made after 15-20 days from the first sizing spray are less effective.

Crop/Cultivar	Target Berry Diameter*	Rate (grams a.i./acre)
Perlette Seedless	4-5 mm	32-128
Flame Seedless	6-9 mm	20-128
Thompson Seedless	3-5 mm	32-128
Raisin	3-5 mm	4-20
Other Seedless Grapes	3-14 mm	8-60

* Target average berry diameter for the first application.
NOTE: In some growing regions and for some cultivars, high amounts of gibberellic acid will reduce fruitfulness (cluster counts) the following year. High amounts of gibberellic acid will also delay berry skin color development, sugars accumulation and overall maturation. A grower shall consult the local specialist before sizing cultivars with which he has no familiarity.

7.2 SEEDED GRAPE

BERRY SIZING SPRAYS

Objective/benefit	Application timing/instructions		
To increase berry size in listed cultivars; and also to reduce berry shrivel in Emperor.	Make one application during the indicated berry diameter range. Application is made as a whole vine spray, or as a spray or dip directly to the cluster.		
Crop/Cultivar	Berry Diameter (mm)*	Whole vine spray, Rate in grams a.i./acre	Direct spray to the cluster only or dip the clusters. Rate in ppm's of a.i.
Emperor	12-16	20	40-50
Red Globe	12-18		
Calmeria	12-16		
Christmas Rose	12-16		
Rogue	12-16		
Queens	12-15		

* Predominant average berry diameter for this application.
NOTE: The whole vine application will reduce fruitfulness (cluster counts) the following year. High amounts of gibberellic acid will also delay berry skin color development, sugars accumulation and overall maturation. A grower shall consult the Stoller representative or local specialist before sizing cultivars with which he has no familiarity.

Objective/benefit	Application timing/instructions
To increase berry size.	Make one application 3-5 days after full bloom, but before shatter begins.
Crop/Cultivar	Rate (grams a.i./acre)
Black Corinth (Zante Currant)	1-12

8.0 SPRAY GUIDELINES FOR CITRUS

For citrus, apply in sprays of sufficient water volumes to ensure thorough fruit wetting. In most cases, this application will cause some drop of older mature leaves; this drop of older leaves is inconsequential. However, application to trees of low vigor or under stress (pest, nutritional, or water, etc.) will cause severe leaf and/or fruit drop. Do not apply in white wash sprays in which lime or other caustic material has produced a high pH in the spray tank. Applications of copper fungicides and/or oils within three weeks (before or after) the N-LARGE application will result in significant leaf drop and fruit drop.

8.1 CITRUS: FIELD APPLICATIONS

Crop/ Variety	Objective/ Benefit	Rate (grams a.i./ acre)	Application Timing/ Instructions
Navel Orange	To delay rind aging, reduce physiological disorders (e.g., rind staining, water spotting, sticky or tacky surface, puffy rind and rupture under pressure), and produce a more orderly harvesting pattern.	16-48	Make one or two applications as a concentrate or dilute spray. 1) Early application: spray approximately 2 weeks prior to color break (typically August-November). This timing causes the greatest delay in rind aging and produces the firmest rind possible. AND/OR 2) Late spray: one application after marketable color (typically October-December). Late sprays cause re-greening.
Valencia Orange (For California and Arizona use only)	To reduce rind creasing and to delay rind aging and softening.	40-80	Make a single application as a concentrate or dilute spray in August to October to target crop of young fruit.
NOTE: Do not apply the early spray to groves that may be harvested early, as fruit coloring will be delayed. Do not apply from January through July, as production will be reduced the following year. Slower color development and increased re-greening of mature fruit is to be expected in the target crop. After marketable color is achieved, treatment effects will be reduced the longer treated fruit remain on the tree.			
All round Oranges (For Florida use only)	To delay aging and softening of the rind, and to reduce creasing and puffiness.	20-60	Make a single application in August to October to trees with a target crop of young fruit. The addition of pure organo-silicone type surfactant at 0.05% (6 fl. oz. in 100 gallons) is beneficial.
Lemon Lime	To increase the amount of small ripe fruit and produce a more desirable production pattern relative to market demand.	10-32	Make a single application when target crop is 1/2 to 3/4 full size, but still green.
NOTE: When applied two years in a row, an even larger difference in harvest pattern and maturity will occur.			
Tangerine Hybrids: Orlando, Robinson, Minneola, Sunburst, and others (All States Except CA)	To delay disorders associated with rind aging, puffiness, and softening, and to increase peel strength of tangerine hybrids.	20-40	Make one spray application two weeks prior to color break. Apply as a dilute spray.
NOTE: Do not apply if early harvest is planned. Do not apply after coloring as pre-harvest rind staining may occur. Application during coloring causes variation in rind color development.			
Grapefruit (All States Except CA)	To delay disorders associated with rind aging (e.g. puffiness, softening, and orange coloration) prevent preharvest drop of	16-48	Make one or two dilute spray applications in sufficient volume to ensure coverage. Do not exceed 20 ppm a.i. in spray solution.

	mature fruit, increase peel strength, reduce water loss during storage, and produce a more orderly harvesting pattern.		EARLY: Make application two weeks prior to color break. Apply as a dilute spray (Aug-Sept). AND/OR LATE: Make application after marketable color has developed (Oct-Dec).
NOTE: Do not spray groves that may be harvested early since fruit coloring will be delayed. Treated fruit will re-green if allowed to remain on the tree for extended periods. Application made after December, or when trees begin to break dormancy, will adversely affect new crop. Do not use concentrate sprays. Results will vary from season to season depending on environmental conditions. The delay in rind aging is greatest when spray is applied before color change. This spray timing produces the firmest rind possible.			

Star Ruby Grapefruit (All States Except CA)	To reduce early-season small fruit drop of Star Ruby Variety thereby increasing yields.	25-35	Make a single dilute application during the bloom period.
NOTE: Results will vary from season to season depending on environmental conditions. Maintain a well-balanced fertilization and watering program.			

Clementine Mandarin	To increase fruit set and yield.	1-8	Make one or two applications from 50% petal fall up to 3 weeks after petal fall. Use a dilute spray with sufficient spray volume for adequate coverage of tree canopy.
NOTE: The number of applications depends upon amount of desired fruit set. Generally, more fruit will be set by 2 applications, earlier applications, higher rates, and climatic conditions more favorable to set. Differences in the crop strain will also interact with the above factors to affect the degree of fruit set achieved. Reductions in final fruit size will occur as a result of excessive fruit set.			

Tangerine Hybrids (Orlando, Robinson, Minneola, Sunburst, and others)	To increase fruit set and yield. The number of applications depends on desired fruit set.	8-30	Make one to two applications during the bloom period. Apply as a dilute spray.
NOTE: Fruit sizes will be reduced and color development slightly retarded. A slight increase in mature leaf drop will occur in trees under stress.			

Navel and Valencia Orange (for Florida use only)	To enhance fruit set and yield.	15-25	Make a single application in Dec-Jan. Apply in 125-175 gallons of water per acre with a pure organo-silicone type surfactant at 0.05% (6 fl. oz/100 gallons).
Amber- sweet Or- ange (For Florida use only)	To enhance fruit set and yield.	15-25	Make a single application in January. Apply in 125-175 gallons of water per acre with a pure organo-silicone type surfactant at 0.05% (6 fl. oz/100 gallons).
Grapefruit	To enhance fruit set and yield.	15-25	Make a single application in Dec-Jan. Apply in 125-175 gallons of water per acre with a pure organo-silicone type surfactant at 0.05% (6 fl. oz/100 gallons).

8.2 CITRUS: POSTHARVEST APPLICATIONS

Lemon	To delay fruit senescence and prolong storage life. The delay in senescence will reduce the incidence of infection by sour rot (<i>Geotrichum candidum</i>).	50-100	Add 2 to 4 fluid ounces of product (2 to 4 grams of a.i.) in 10 gallons of storage wax, which has been diluted as per wax label instructions.
Yellow lemons and other ma-	To delay aspects of rind senescence and color changes	50-100	Add 2 to 4 fluid ounces of product (2 to 4 grams of a.i.) in 10 gallons of stor-

ture citrus fruit			age wax, which has been diluted as per wax label instructions.
-------------------	--	--	--

9.0 SPRAY GUIDELINES FOR FRUIT CROPS

FRUIT CROPS

Crop/Cultivar	Objective/Benefit	Rate (grams a.i./acre)	Application Timing/Instructions
Banana	To stimulate plant growth, and to overcome the effects of stress caused by insect, disease or adverse weather. These applications will also improve fruit size and quality and overall yield.	1-6	Apply by air or ground equipment once every 30 to 90 days throughout the year. Use sufficient water volume to achieve good coverage of the foliage. Make more frequent applications (monthly) during the 6 months prior to anticipated weather stress periods.
Banana	To extend storage life.	1-2	Mix 1 to 2 grams/liter of water and spray directly on the banana fingers from 30 days before harvest until harvest. One to two applications are to be used.
Blueberry <u>Highbush:</u> Coville, Jersey, Stanley, Earlibluc, Weymouth, Walcott, Berkeley, Bluecray, Bluecrop, 1316A, Concord, and others.	To improve fruit set.	40-80	Make a single application of 80 grams a.i. in 40 to 100 gallons of water/acre. The application shall be made at full bloom (when 75% of the flowers are fully open). OR Make two applications at 40 grams a.i./acre in 40 to 100 gallons of water. Make the first application at full bloom, and the second one within 10-14 days of the first one. For Weymouth, application shall be delayed up to two weeks after bloom to increase size of "shot" berries.
Blueberry <u>Rabbiteye:</u> Aliceblue, Beckyblue, Bonita, Brightwell, Climax, Delite, Tiftblue, Woodward, and others.	To improve fruit set.	40-80	Make a single application of 40 to 80 grams a.i. acre in 40 to 100 gallons of water per acre when most of the flowers are elongated but not yet open (bloom stage 5). OR Make two to four application 10 to 14 days apart starting at bloom Stage 5. Spray 20 to 40 grams a.i. acre in 40 to 100 gallons of water per application.
Sweet Cherry	To produce larger, brighter colored, firmer fruit.	16-48	Apply a single spray when the fruit is translucent green to straw colored. Use sufficient water volume to ensure thorough wetting.
NOTE: Color development and harvest date will be slightly delayed.			
Red Tart Cherry	To maintain and extend high fruiting capacity of tart cherry trees and reduce the occurrence of "blind" nodes. Treatment will	4-18	Apply one spray 14 to 28 days after bloom. Optimum timing is defined as that stage when 3 to 5 terminal leaves have fully expanded, or, at least 1 to 3 inches of terminal shoot extension has occurred. Use 4 to 18 grams

	cause bud differentiation, which is apparent the year after application. Therefore, changes in shoot, spur, and flower production will not be evident until two or three years after program initiation. Applications must be applied annually to ensure vegetative development and subsequent yield improvement year after year.		a.i./acre, depending on tree age and vigor (See Table below). Apply as a concentrate or dilute spray in sufficient water volume to ensure thorough wetting.
--	---	--	---

NOTE: Rates are based on expected normal tree vigor at various ages. Adjust rate according to tree vigor. If trees are vigorous, use lowest recommended rates. Lowest rates must also be used on trees that have been heavily pruned or hedged. Use higher rates for trees low in vigor and weak in shoot and spur production. Excessive application rates will increase vegetative growth at the expense of fruit production the following year. Applications will not improve growth of trees under stress conditions, such as nutritional, moisture, or pest. Best results will be obtained when combined with good cultural practices.

Tree Age (years)	Rate (grams a.i./acre)
6-10	4-6
11-15	8-10
16-20	10-14
20+ years	14-18

STONE FRUITS			
Stone Fruit Group	To increase fruit firmness and improve fruit quality in the season of application	16-32	Apply as a single spray one to four weeks prior to the beginning of the harvest period. Use sufficient water to achieve complete coverage of fruits and foliage.

NOTE: This application will cause reduction in flower counts the year following the application, particularly if it is made during the months of May through July

Italian Prune (All States Except CA)	To reduce internal browning, improve quality, and increase size.	16-48	Make a single application four to five weeks before expected harvest. Apply in sufficient water volume to ensure thorough wetting.
NOTE: Color development and harvest will be slightly delayed. Will reduce bloom the following season			

10.0 SPRAY GUIDELINES FOR NON-BEARING FRUIT TREES AND OTHER CROPS

Crop/Variety	Objective/Benefit	Rate (grams a.i./acre)	Application Timing/Instructions
Non Bearing Stone Fruit	To reduce flowering and fruiting in young stone fruit trees in order to minimize the competitive effect of early fruiting on tree development.	20-80	Make a single application during the period of flower bud initiation for the following year. Consult with the local horticulturist for timings and rates for specific cultivars in your area. Use sufficient water to achieve good coverage of the canopy.

NOTE: Do not spray trees in the first year. Treat in the second season for reduction of flowering in the third season, and again in the third season if flower

reduction and fruiting is desired in the fourth season. Treat only trees that are in good physiological condition.
Discontinue treatment the year before desired harvest.

Strawberry	To increase runner production of mother plants.	15-25	Make a single application to mother plants 10-30 days after planting. Plants must have 1-6 leaves at spraying. Apply 100 gallons spray/acre to point of run-off.
------------	---	-------	--

NOTE: Not for use on fruiting plants. Treatments will not be effective on plantings set out after mid-May. Response varies with cultivar and location. Consult local horticulturist for specific instructions.

Cranberry (All States Except CA)	To reduce or completely eliminate the crop in the year of application	10-50	Make a single application at early bloom (2-5% scatter bloom). Use sufficient water to ensure thorough coverage.
----------------------------------	---	-------	--

NOTE: Applications made later than indicated will result in no effect or actually result in increased fruit set (opposite effect). Responses will vary with cultivar, age of the bog and location. Consult the local specialist for specific information.

Pineapple	To shape fruit	120 grams a.i./acre	Make 1 to 2 applications per crop cycle of 14 to 18 months.
-----------	----------------	---------------------	---

11.0 SPRAY GUIDELINES FOR VEGETABLE CROPS

VEGETABLE CROPS

Crop/Variety	Objective/Benefit	Rate (grams a.i./acre)	Application Timing/Instructions
Artichoke	To accelerate maturity and shift harvest to an earlier date.	10-20	For perennials: Apply 1 to 3 applications at bud initiation stage. For annuals: Apply 1 to 4 applications at 2-week intervals, beginning at the fourth true leaf. Use sufficient water volume to ensure thorough wetting of the entire plant (leaves, stems and buds).
Carrots, Fresh and Processing	To delay leaf senescence. Maintaining vigorous foliage will reduce the incidence of infection by <i>Ather-naria dauci</i> .	1-6	Make the first application 4-6 weeks after emergence using commercial ground or aerial equipment with spray concentrations of 20-30 ppm. In severe disease situations or cool weather a second spray 14 days later will be required to achieve the desired amount of foliar recovery. Do not apply more than twice per crop.

NOTE: Dilutions of greater concentration will increase the risk of excessive top growth, particularly with a second application.

Celery	To increase plant height and yield and to overcome stress due to cold weather conditions or saline soils, and obtain earlier maturity.	2.5-10	Make a single application one to four weeks prior to harvest. Use 25 to 50 gallons of water per acre by ground application or 5 to 10 gallons of water per acre for aerial application (except in California). Use lower concentrations if applying 3 to 4 weeks before harvest and higher concentrations within 1 to 2 weeks before harvest.
--------	--	--------	---

NOTE: Do not apply by air in California. Do not apply earlier than 4 weeks before harvest as bolting will occur.

Cucumber	To stimulate fruit set during	1-4	Make one application prior to bloom followed by two addi-
----------	-------------------------------	-----	---

	periods of cool temperatures.		tional applications at intervals of 10 to 14 days. Up to four applications are required. Use sufficient water volume for thorough coverage of exposed foliage.
--	-------------------------------	--	--

NOTE: For maximum benefits, vines must be in good condition, except for reduced rate of growth due to cool temperatures.

Lettuce for Seed	To obtain uniform bolting and increase seed production.	1-4	Apply one to four applications at two-week intervals, beginning at the fourth true leaf. Use sufficient water volume to ensure thorough wetting.
------------------	---	-----	--

Melon	To stimulate fruit set during periods of cool temperatures.	1-4	Make one application prior to bloom followed by two additional applications at intervals of 10 to 14 days on cantaloupes and watermelons.
-------	---	-----	---

NOTE: For maximum benefits, vines must be in good condition, except for reduced rate of growth due to cool temperatures.

Pepper (All States Except CA)	To promote plant growth.	1-3	Apply one to two sprays in 25 to 50 gallons of water per acre at two-week intervals. Begin sprays 2 weeks after transplanting.
-------------------------------	--------------------------	-----	--

NOTE: This use is recommended for acres with short growing season, or when low temperatures slow plant growth.

Pepper (All States Except CA)	To increase fruit set and promote fruit growth.	1-3	Apply one to two sprays in 25 to 50 gallons of water per acre at weekly intervals during the flowering period.
-------------------------------	---	-----	--

NOTE: The high rate is recommended for areas and/or varieties with pollination and/or fruit set problems.

Pepper (All States Except CA)	To increase fruit size.	1-3	Apply in 25 to 50 gallons of water per acre at the beginning of the picking period.
-------------------------------	-------------------------	-----	---

NOTE: Use the highest rate for plants with heavy fruit loads.

Potato seed	To stimulate uniform sprouting to aid in maximum production, more uniform development, fewer late maturing plants, and to break dormancy of newly harvested potatoes that have not had a full rest period.	0.2-0.4 (grams in 100 gallons)	Dip whole or cut seed pieces in a solution containing 0.2 to 0.4 grams a.i. in 100 gallons of water prior to planting.
-------------	--	--------------------------------	--

NOTE: Under high soil temperatures use the minimum concentration for dormant seed. Do not treat rested seed pieces.

Rhubarb	To break dormancy on plants receiving insufficient chilling and to increase marketable yield of forced rhubarb.	10-20 (grams in 10 gallons)	1) When the rest period is not completely broken, make a single application of 2 fluid ounces (60 ml) of a solution containing 20 grams a.i. in 10 gallons of water to each cleaned crown. 2) When the rest period is broken by cold weather, apply 2 fluid ounces (60 ml) of a solution containing 10 grams a.i. in 10 gallons of water to each cleaned crown.
---------	---	-----------------------------	--

NOTE: Keep forcing house temperatures at 40°F-50°F for 24 hours after application. If house is warmer than 50°F, crowns must be covered with plastic. Temperatures above 50°F will lower yields and cause poor stalk color.

Spinach (All States Except CA)	To facilitate harvest, increase yield and improve quality of fall and over-winter spinach.	6-10	Apply in a single spray 10 to 18 days before each anticipated harvest on fall or over-winter spinach, ideally when daytime temperatures are
--------------------------------	--	------	---

			40°F to 70°F and during early morning hours when dew is present on crop. Make applications in 10 to 50 gallons of water per acre by ground sprayer or in a minimum of 5 to 10 gallons of water per acre by air. When applied to promote growth of second cutting, wait until some re-growth has started before spraying. Maximum benefit is obtained when below normal temperatures predominate following application and growth would be otherwise slowed in untreated spinach.
--	--	--	--

NOTE: Since the promotion of bolting will occur, do not apply after the mid-winter period or if temperatures are expected to exceed 75°F within several days of application. Do not apply on spring planting.

**12.0 SPRAY GUIDELINES FOR OTHER CROPS
COTTON, HOPS, AND RICE**

Crop/Variety	Objective/ Benefit	Rate (grams a.i./acre)	Application Timing/ Instructions
Cotton (All States Except CA)	To promote early plant growth and increase seedling vigor.	1-6	Apply as an in-furrow application to seed or as a foliar application from the cotyledon leaf stage through the 7 leaf/node stage. Up to three applications are to be made as needed. To mix, fill the treatment tank with half the final tank mix volume. Add the required amount of N-LARGE and mix thoroughly while adding water to the desired final volume. Compatibility information regarding tank mixtures of N-LARGE with herbicides used in cotton is not available. Aerial application: Use a spray system capable of producing a uniform spray pattern of medium to fine spray droplets at 10 gallons per acre (GPA). Apply no less than 3 GPA of total spray volume. Ground application: For low pressure ground sprayers equipped with boom and flat fan nozzles, apply 10 to 15 GPA spray volume. Dispose of unused spray mixture according to the label directions at the end of the day.

NOTE: Use higher rates when temperatures will likely average 75°F or less during the 14 days following the application. Do not apply more often than necessary to achieve the desired height, as over-dosage will result in excessive growth. Do not apply to cotton plants under drought stress.

Hops: Seeded and seedless Fuggle hops and similar varieties adapted to the	To increase fruit set and yield	4-6	Make a single application in 100-150 gallons of water per acre when vine growth is 5-8 feet in length.
--	---------------------------------	-----	--

Northwestern U.S.			
Rice Seed Treatment	For use as a seed treatment of both semi-dwarf and tall rice varieties to promote germination, emergence and final stand densities when planted at greater depths where soil moisture levels are more adequate for germination.	0.5-2.1	Use in 8 to 20 oz. water per 100 pounds of rice seed. N-LARGE is to be applied to dry seed with standard mist-treating equipment. Best results are obtained using a higher treatment volume (12 to 20 fl. oz. per 100 pounds of seed) to ensure the seed is completely and uniformly covered with N-LARGE. Fill the seed treatment tank with water to one-half the final tank mix volume. Add the required amount of N-LARGE mixing thoroughly while adding water and other seed treatment products to the desired final volume.

NOTE: Apply only to rice seed intended for drill seeded or dry broadcast systems. Do not apply to rice used in a 24-hour presoak prior to broadcast or to water used for the presoak. Do not use more than 2.1 grams a.i. per 100 pounds of seed. DO NOT USE TREATED SEED FOR FOOD, FEED, OR OIL PURPOSES.

An approved dye must be added to distinguish N-LARGE treated seed and prevent inadvertent use of food, feed or oil purposes. Seed commercially treated with this product must be labeled in accordance with all applicable requirements of the federal and state seed laws. N-LARGE is compatible with most commonly used fungicide seed treatments such as VITAVAX® and DITHANE®, standard dyes and sticker-binding agents. When preparing tank mixes, the user must ensure adequate physical compatibility and mixing characteristics.

Rice Post-Emergent Seedling Treatment	For use as a post-emergence seedling application on rice grown in the United States to promote more uniform and vigorous growth of rice prior to permanent flood establishment. This will allow earlier (five to ten days) flooding of drill or dry broadcast seeded varieties and is particularly effective on semi-dwarf varieties. Early flooding will reduce additional flushing costs associated with delay in permanent flooding, weed infestations and the number of herbicide applications as well as promote earlier and more uniform grain maturity.	1-3	Apply to rice between the 1 to 2 leaf stage and the 4 to 5 leaf stage of growth. Timing and dosage is based on environmental conditions, tank mix combinations with herbicides and method of permanent flood practice in relation to rice leaf stage.
---------------------------------------	--	-----	---

NOTE: N-LARGE application will result in a temporary lighter green foliage color due to accelerated growth rates. Do not apply when rice is subject to drought stress conditions. N-LARGE may be tank mixed with most commonly used rice herbicides and fungicides. When N-LARGE is applied in tank mixes with Arrosolone®, Riverside Propanil® 60 DF, Stam® 80 EDF or WHAM® EZ, plus a recommended adjuvant, the use of a surfactant is not necessary. Do not apply N-LARGE with products containing fenoxaprop-p-ethyl as the active ingredient. When preparing tank mixes, ensure

adequate physical compatibility and mixing characteristics. Refer to the paragraph on Compatibility under the APPLICATION INSTRUCTIONS section of this label for additional information.

N-LARGE applied between split-boot and 100% heading will increase panicle height of semi-dwarf rice. This will facilitate harvest efficiency in the field by allowing the rice grain to be cut above the leaf canopy at faster combine speeds and at reduced vegetative load. Grain quality and maturity will be advanced with the promotion of tiller panicle development. Heading applications to the first crop will also accelerate regrowth of second crop rice. This will result in earlier second crop maturity and maximize grain yield.

Hybrid Rice: Seed Production	Apply N-Large to facilitate main culm and tiller panicle extension to increase pollination and harvest efficacy	20-100	Make 1 to 5 applications at regular intervals during the heading period.
------------------------------	---	--------	--

13.0 SPRAY GUIDELINES FOR ORNAMENTALS, CUT FLOWERS & TURFGRASS

The following instructions are based on results with common cultivars. Differences in responsiveness will vary from one cultivar to another, or from one set of growing conditions to another, or from one cultural management system to another. Therefore, prior to widespread usage, test a small number of plants from each cultivar under a specific set of growing and cultural management conditions to verify desired efficacy.

13.1 ORNAMENTALS

Crop/Variety	Objective/Benefit	Rate (grams a.i./acre)	Application Timing/Instructions
Azalea (All states except California)	As a partial replacement of cold treatment to break flower dormancy.	250-500 ppm	Apply three sprays at weekly intervals after three to four weeks of chilling.
NOTE: Initiate treatment when plants are at Stage 5 of floral development (i.e. style elongated and open). A representative spray schedule consists of applications made at 3, 10 and 17 days after four weeks of chilling. Flowers will not develop properly if applied prior to Stage 5. Do not apply after flower buds show color. To ensure uniform flowering, apply thoroughly.			
Azalea (All states except California)	To break dormancy on some cultivars (e.g. 'Gloria', 'Prize', and 'Redwing').	1000 ppm a.i.	Apply after three to four weeks of chilling
Azalea (All states except California)	As a complete substitution of cold treatment to break flower dormancy.	1000 ppm a.i.	Apply four to six sprays at weekly intervals. Plants must be at Stage 5 of floral development (style elongated and open) before first spray is applied.
NOTE: Flowers will not develop properly if applied prior to Stage 5 of floral development. Do not apply after flower buds show color. To ensure uniform flowering, apply thoroughly.			
Azalea (All states except California) Flower Bud Initiation	To inhibit flower bud initiation during vegetative growth.	100-750 ppm a.i.	Approximately 2 to 3 weeks after each pinch, apply a single foliar application. After the first application, continue applying on a weekly basis for 1 to 2 weeks.
NOTE: Make a maximum of three applications.			
Calla Lily (All states except California)	For increased flowering.	500 ppm a.i.	Prepare a solution and soak rhizome or tuber for 10 minutes prior to planting.
NOTE: Leaf or flower stretching will be observed in some cultivars. If this occurs, reduce rates.			
Camellia (All)	To substitute for	2% a.i. solution	Mix equal vol-

States Except CA)	chilling requirements and increase bloom size.		umes of product and water. After removing the vegetative bud, found immediately adjacent to or below the floral bud, place a single drop of the prepared solution on the vegetative bud scar.
NOTE: Adding a deposition aid (e.g., carboxymethylcellulose) to thicken the solution will reduce run-off.			
Cyclamen (all states except California) - Bud Application	To promote uniform flowering.	0.25 fl. oz. 10 to 15 ppm a.i.	Apply a single application of 8 ml (0.25 fl. oz.) of a 10 to 15 ppm a.i. solution directly to the crown when buds are pinhead size in the leaf axils.
Cyclamen (all states except California) - Foliar Application	To promote uniform flowering.	25 ppm a.i.	Thoroughly wet the crown by applying a single foliar application directly toward the crown and adjacent leaves when buds are pinhead size in the leaf axils.
NOTE: Both bud and foliar applications have been shown to promote uniform flowering. Late or excessive applications will result in poorly formed flowers or weakened stems.			
Fuchsia (all states except California)	To produce tree forms of common fuchsia cultivars by stem elongation.	250 ppm a.i.	Apply a foliar application beginning after the fuchsia plant has reached the desired size and continuing for four consecutive weeks. Spray plant to point of run-off.
NOTE: Staking will be required after application. Higher concentrated solutions will cause long, spindly and weak stems.			
Geranium (all states except California)- Cuttings	To increase number and size of flowers.	1-5 ppm a.i. solution	Apply when inflorescence first begins to show color. Apply spray to the developing inflorescence.
NOTE: Peduncle stretching will be observed if application is made prior to inflorescence showing color or if concentrations in excess of 5 ppm are used.			
Geranium (all states except California) - Seedlings	To advance flowering.	5-15 ppm a.i.	Apply a single application when the first flower bud set is noted. Spray plant to point of run-off. Depending on type of geranium, flowering will be advanced 10 to 21 days.
NOTE: Overuse or incorrect timing will cause long, spindly and weak stems.			
Geranium (all states except California) Tree Forms	To produce tree forms of common geranium cultivars by stem elongation.	250 ppm a.i.	Apply a foliar application for four consecutive weeks spraying plant to point of run-off.

NOTE: Staking will be required after application.

Hydrangea (all states except California)	To substitute for chilling requirements and break flower bud dormancy.	2-5 ppm a.i.	Apply a single foliar application for one to four consecutive weeks beginning at the start of forcing. Thoroughly apply solution to all growing points containing flower buds.
--	--	--------------	--

NOTE: Overuse or incorrect timing will cause long, spindly and weak stems.

Pompom Chrysanthemums (all states except California)	For elongating peduncles on pompom chrysanthemums.	25-60 ppm a.i.	Apply a single spray four to five weeks after initiation of short day conditions. Apply spray towards the flower buds.
--	--	----------------	--

NOTE: Overuse or incorrect timing will cause long, spindly and weak stems.

Spathiphyllum (all states except California)	To induce flowering of spathiphyllum.	150-250 ppm a.i.	Apply single full coverage spray approximately nine to twelve weeks prior to sale. Spray plant to point of runoff, thoroughly wetting all growing points.
--	---------------------------------------	------------------	---

NOTE: Distorted bloom, increased petiole length and narrower leaves will appear on some cultivars such as 'Petite', 'Starlight', 'Tasson', and 'Mauna Loa'. For other cultivars, prior to application on a commercial basis, evaluate the effects of N-LARGE on a small number of plants.

Aglaonema, Anthurium, Dieffenbachia (Dumb Cane) (all states except California)	To accelerate bloom and increase flowering.	250-500 ppm a.i.	Apply a single foliar application for one to four consecutive weeks beginning at the start of forcing.
Syngonium (all states except California)		500-2000 ppm a.i.	Apply a single foliar application for one to four consecutive weeks beginning at the start of forcing. Thoroughly apply solution to all growing points containing flower buds.

NOTE: Applying N-LARGE will increase flower yield and decrease time to flowering. To induce bloom, make 1 to 2 application while plant is in the vegetative phase. For other Araceae cultivars, prior to application on a commercial basis, evaluate the effects of N-LARGE on a small number of plants.

13.2 CUT FLOWERS

NOTE: Applying N-LARGE to ornamental plants grown for cut flowers will aid in promoting longer stems and increased flower yield. Gibberellic Acid is a potent plant growth regulator and overuse will result in undesirable effects. Assess the effects of N-LARGE on a small number of plants prior to making large-scale applications.

Crop/Variety	Objective/Benefit	Rate (grams a.i./acre)	Application Timing/Instructions
Aster (all states except California) - Monte Carlo type, Novitype and Belgitype	To aid in promoting longer stems and increased flower yield.	50-100 ppm a.i.	Apply 1 to 3 applications when plants are 2" to 6" tall. Make applications at 2 to 3 week intervals.
Baby's Breath	To promote	150-500	Make 3 to 4 application

(Gypsophila) (all states except California)	plant growth, increase flower yield and uniformity.	ppm a.i.	of a solution at 4 weeks of growth (after pinching). Make applications at 2 week intervals.
Bells of Ireland (Moluccella) (all states except California)	To promote plant growth and longer stems	50-100 ppm a.i.	Apply when plants are 4" to 8" tall. Make applications at 2 to 3 week intervals.
Bupleurum (all states except California)	To promote plant growth and longer stems.	50-100 ppm a.i.	Apply solution as a foliar spray when plants are 4" to 8" tall. Make applications at 2 to 3 week intervals.
Campanula (all states except California)	To promote plant growth and longer stems.	50-100 ppm a.i.	Apply solution as a foliar spray when plants are 4" to 8" tall. Make applications at 2 to 3 week intervals.
Candy Tuft (Iberis) (all states except California)	To promote plant growth and longer stems.	50-100 ppm a.i.	Apply solution as a foliar spray when plants are 4" to 8" tall. Make applications at 2 to 3 week intervals.
Column Stock (Matthiola) (all states except California)	To promote plant growth and longer stems.	50-100 ppm a.i.	Apply solution as a foliar spray when plants are 4" to 8" tall. Make applications at 2 to 3 week intervals.
Delphinium including <i>D. belladonna</i> , <i>D. bellamosum</i> , <i>D. cardinale</i> , <i>D. elatum</i> , <i>D. grandiflorum</i> , <i>D. nudicale</i> , and Delphinium hybrids (all states except California)	To promote plant growth and longer stems.	50-100 ppm a.i.	Apply solution as a foliar spray when plants are 4" to 8" tall. Make applications at 2 to 3 week intervals.
Didiscus (Torch) (all states except California)	To promote plant growth and longer stems.	50-100 ppm a.i.	Apply solution as a foliar spray when plants are 4" to 8" tall. Make application at 2 to 3 week intervals.
Hydrangea (all states except California)	To promote plant growth and longer stems.	50-100 ppm a.i.	Apply solution as a foliar spray when plants are 4" to 8" tall. Make applications at 2 to 3 week intervals.
Larkspur (<i>Consolida ambigua</i> , <i>C. orientalis</i> , <i>Delphinium ajacis</i>) (all states except California)	To promote plant growth and longer stems.	50-100 ppm a.i.	Apply solution as a foliar spray when plants are 4" to 8" tall. Make applications at 2 to 3 week intervals.
Lisianthus (Eustoma) <i>Eustoma grandiflora</i> (all states except California)	To promote plant growth and longer stems.	50-100 ppm a.i.	Apply solution as a foliar spray when plants are 4" to 8" tall. Make applications at 2 to 3 week intervals.
Phlox (<i>Phlox paniculata</i> and <i>Drummondii hybrida</i>) (all states except California)	To promote plant growth and longer stems.	50-100 ppm a.i.	Apply solution as a foliar spray when plants are 4" to 8" tall. Make application at 2 to 3 week intervals.
Queen Anne's Lace (Ammi) (all states except California)	To promote plant growth and longer stems.	50-100 ppm a.i.	Apply solution as a foliar spray when plants are 4" to 8" tall. Make applications at 2 to 3 week intervals.
Safflower (Carthamus) (all states except California)	To promote plant growth and longer stems.	50-100 ppm a.i.	Apply solution as a foliar spray when plants are 4" to 8" tall. Make

California)			applications at 2 to 3 week intervals.
Solidaster (Solidago) (all states except California)	To promote plant growth and longer stems.	50-100 ppm a.i.	Apply solution as a foliar spray when plants are 4" to 8" tall. Make applications at 2 to 3 week intervals.
Statice (Limonium) (all states except California)	To promote earlier flowering and to increase flower yield.	10 ml of a 400-500 ppm a.i.	Apply as a foliar spray when plants are more than 10 inches in diameter (approximately 90 to 110 days after normal seeding time).
NOTE: Do not exceed specified rates. Do not apply repeated sprays. Accelerated flowering is influenced by extended photoperiod, adequate nutrition and reduced night temperature. Treatment with Gibberellins lessens the requirement for the cold requirement and/or the long photoperiod.			
Statice (Limonium) (all states except California)	To promote plant growth and longer stems.	50-100 ppm a.i.	Apply solution as a foliar spray when plants are 4" to 8" tall. Make applications at 2 to 3 week intervals.
Sunflower (Helianthus) (all states except California)	To promote plant growth and longer stems.	50-100 ppm a.i.	Apply solution as a foliar spray when plants are 4" to 8" tall. Make applications at 2 to 3 week intervals.
Sweet William (Dianthus) (all states except California)	To promote plant growth and longer stems.	50-100 ppm a.i.	Apply solution as a foliar spray when plants are 4" to 8" tall. Make applications at 2 to 3 week intervals.

13.3 BEDDING PLANTS, ANNUAL AND PERENNIAL POTTED CROPS, FIELD GROWN ORNAMENTALS AND BULB CROPS

Crop/Variety	Objective/Benefit	Rate (grams a.i./acre)	Application Timing/Instructions
Bedding Plants, Annual and Perennial Potted Crops, Field Grown Ornamentals and Bulb Crops (all states except California)	To promote plant growth and/or overcome the effects of excessive use of a gibberellin inhibiting plant growth regulator.	1-25 ppm a.i.	Begin by applying a single foliar application of a 1 ppm a.i. solution unless experience dictates a higher rate is appropriate. If desired results are not achieved, a reapplication or increased rate will be necessary. Do not use more than 25 ppm a.i.

NOTE: Gibberellic Acid is a potent plant growth regulator and overuse will result in undesirable effects including stem elongation. Assess the effect of N-LARGE on a small number of plants prior to making large scale applications.

13.4 TURFGRASS

Crop/Variety	Objective/Benefit	Rate (grams a.i./acre)	Application Timing/Instructions
Bermudagrass Tidwarf, Tif-green, and other cultivars (all states except California)	To initiate or maintain growth and prevent color change during periods of cold stress and light frosts.	10-25 grams a.i.	Apply 10 grams a.i. per acre weekly or 25 grams a.i. per acre biweekly in 25 to 100 gallons of water per acre.

NOTE: Application of N-LARGE to Bermudagrass grown in golf courses, parks and turf farms has been shown to initiate or maintain growth and prevent color change during periods of cold stress. Do not exceed specific rates. Maintain adequate moisture and proper fertilization programs as indicated for the local area. Discontinue treatments if thinning is observed. Do not apply the high rate more frequently than every two weeks. More frequent mowing will be necessary. Do not use on dormant turf.

Bermudagrass Tidwarf, Tif-green (all states except California)	To maintain or enhance regrowth of golf course Bermudagrass during summer months.	1-3 grams per acre	Apply weekly in 25 to 100 gallons of water per acre.
--	---	--------------------	--

NOTE: Application of N-LARGE to Bermudagrass grown in golf courses, parks and turf farms has been shown to initiate or maintain growth and prevent color change during periods of cold stress. Do not exceed specific rates. Maintain adequate moisture and proper fertilization programs as indicated for the local area. Discontinue treatments if thinning is observed. Do not apply the high rate more frequently than every two weeks. More frequent mowing will be necessary. Do not use on dormant turf.

14.0 CONVERSION TABLE (G/FL. OZ.)

N-LARGE contains approximately 1 gram of active ingredient per fluid ounce of product.

Grams of active ingredient	Fluid ounces of N-LARGE
0.5	0.5 oz.
1.0	1 oz.
2.0	2 oz.
4.0	4 oz.
5.0	5 oz.
8.0	8 oz.
10.0	10 oz.
12.0	12 oz.
16.0	16 oz.
20.0	20 oz.
25.0	25 oz.
32.0	32 oz.
40.0	40 oz.
48.0	48 oz.
50.0	50 oz.

15.0 CONVERSION TABLE (PPM)

Volume of N-LARGE to use in water spray to provide the desired parts per million (ppm) spray.

Gibberellic Acid (GA ₃) ppm (parts per million)	N-LARGE milliliters (mL) per liter of spray	N-LARGE milliliters (mL) per gallon of spray	N-LARGE fl. oz. per gallon of spray
1	0.03	0.1	0.003
5	0.15	0.6	0.02
10	0.3	1.1	0.04
25	0.74	2.8	0.09
50	1.5	5.6	0.18
100	3.0	11.2	0.4
250	7.4	28	0.95
500	14.8	56	1.9
750	22.2	84	2.8
1000	29.6	112	3.8

Arrosolo® is a registered trade name for Syngenta Crop Protection, Inc.
 Dithane® is a registered trade name for Dow AgroSciences L.L.C.
 Stam® is a registered trade name for Dow AgroSciences L.L.C.
 Vitavax® is a registered trade name for Unroyal Chemical Co., Inc.
 Wham® is a registered trade name for RiceCo.

16.0 STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
Pesticide Storage: Keep containers tightly closed when not in use. Store away from any heat source.
Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **Container Disposal:** Do not reuse empty containers. Triple rinse or equivalent; then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

17.0 WARRANTY

To the fullest extent permitted by law, neither the manufacturers nor the seller make any warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use of this material when such use is contrary to label instructions. Read and follow the label directions carefully.