



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

November 14, 2022

Jeremy D. Malone
Consultant
Stoller Enterprises, Inc.,
c/o Spring Regulatory Sciences
6620 Cypresswood Dr,
Suite 250,
Spring, TX 77379

Subject: Non-PRIA (Pesticide Registration Improvement Act) Labeling Amendment – Revision of product label to add optional marketing claims, the addition of chemigation to the commercial sublabel, additional crops added to the label (with efficacy data on-file), metric units added in addition to US units in several portions of the label, updates to EPA standard wording to the latest guidance included in the label review manual, and significant formatting updates for easier readability of the label
Product Name: Fortified Stimulate
EPA Registration Number: 57538-49
EPA Receipt Date: 04/11/2022
Action Case Number: 00354389

Dear Mr. Malone:

The amended labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable.

This approval does not affect any terms or conditions that were previously imposed on this registration. You continue to be subject to existing terms or conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one (1) copy of the final printed labeling before you release this product for shipment with the new labeling. In accordance with 40 CFR § 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. “To distribute or sell” is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR § 152.3.

Should you wish to add/retain a reference to your company’s website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the U.S. Environmental Protection Agency (EPA). If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration

Page 2 of 2

EPA Reg. No. 57538-49


Action Case No. 00354389

process. Therefore, should the EPA find or if it is brought to our attention that a website contains statements or claims substantially differing from statements or claims made in connection with obtaining a FIFRA section 3 registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6.

If you have any questions, please contact Sydnie Hetzel by phone at (202) 566-1606 or via email at hetzel.sydney@epa.gov.

Sincerely,

A handwritten signature in blue ink, appearing to read "James Parker", with a long, sweeping underline that extends to the left.

James Parker, Team Leader
Biochemical Pesticides Branch
Biopesticides and Pollution
Prevention Division (7511M)
Office of Pesticide Programs

Enclosure

[Denotes Optional Text]
 {Denotes Notes to EPA Reviewer}
 <Notes to Stoller Reviewer>
 <EPA MASTER LABEL>

FORTIFIED STIMULATE
YIELD ENHANCER® PLUS
[PLANT GROWTH PROMOTER]
Sublabel A FOR AGRICULTURAL/COMMERCIAL USE
Sublabel B FOR RESIDENTIAL USE

ACTIVE INGREDIENTS:

| | |
|------------------------------|--------|
| Cytokinin (as kinetin) | 0.009% |
| Gibberellic Acid..... | 0.005% |
| Indole-3-butyric Acid | 0.005% |
| Indole-3-acetic Acid | 0.005% |

| | |
|---------------------------------|----------------|
| OTHER INGREDIENTS: | <u>99.976%</u> |
| Total | 100.000% |

This product contains: 0.000095 oz (2.7 mg) of cytokinin, 0.000053 oz (1.5 mg) of gibberellic acid, 0.000053 oz (1.5 mg) of indole-3-butyric acid, and 0.000053 oz (1.5 mg) of indole-3-acetic acid per fluid ounce.

KEEP OUT OF REACH OF CHILDREN
CAUTION

See additional Precautionary Statements, First Aid and Directions for Use [inside booklet] [on [back panel] [side panel] [other panel]].

Z-FORTSTIMYEP

| FIRST AID | |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| If inhaled | <ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice. |
| If in eyes | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eye. • Call a poison control center or doctor for treatment advice. |
| If on skin or clothing | <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice. |
| Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information concerning this product, call the National Pesticides Information | |

Center (NPIC) at 1-800-858-7378 seven days a week, 6:30 am to 4:30 pm Pacific Time (NPIC web site: www.npic.orst.edu).

EPA Reg. No.: 57538-49

EPA Est. Nos.: [57538-FL-1],
[57538-IA-1], [57538-TX-2]

[DENSITY:]
[8.6 lb/gal or 1.04 kg/L]

NET CONTENTS:
 1 Gal (3.8 L) 2.5 Gal (9.5 L) 5 Gal (19 L) 55 Gal (209 L)

[NET WEIGHT:]
 8.6 lb (3.9 kg) 21.5 lb (9.7kg) 43.0 lb (19.5 kg) 473.0 lb (214.5 kg)]

[*Not for use in California]

[Lot Number:]

[Manufacture Date:]

[Best if used by:]

[Rev: 22J21] [revision code for each label change will go here]

[1.0 Quart (0.95 L) 2.2 lb (1.0 kg)]

[30 Gal (114 L) 258.0 lb (117.0 kg)]

[275 Gal (1045 L) 2365.0 lb (1072.5 kg)]

[Bulk]

[Will replace [™] with ® throughout the label when it is a Registered trademark)

[Patents: <https://www.stollerusa.com/about/patents/>]

[ABN: Fortified Stimulate Yield Enhancer]

{Optional Marketing Claims – can appear anywhere on the labels}

[Fortified Stimulate Yield Enhancer® Plus improves genetic potential, root growth, plant vigor and early plant development.]

[Fortified Stimulate® Yield Enhancer® Plus is the only EPA-registered product with four essential plant growth hormones: Cytokinin, Gibberellic Acid, Indole Butyric Acid, Indole Acetic Acid]

[Fortified Stimulate Yield Enhancer® Plus helps plants improve hormone balance and sets the foundation for higher yield.]

[Improves emergence]

[Boosts root growth and plant vigor]

[Improves plant health and physiology]

[Increases yield potential and grain quality]

[Fortified Stimulate Yield Enhancer® Plus improves emergence and germination, while boosting early-stage root growth and plant vigor when applied at planting.]

[Fortified Stimulate Yield Enhancer® Plus is an EPA registered plant growth regulator that has four growth hormone ingredients in a stable formulation that promotes plant growth and improves essential plant functions.]

[Fortified Stimulate Yield Enhancer® Plus contains the plant hormone IAA (indole acetic acid) which is critical for optimizing fast, vigorous root growth and early plant development.]

[Improve] [Protect] [Enable] [Enhance] your yield potential]

[Enhances emergence. Fortified Stimulate Yield Enhancer® Plus boosts plant vigor, promoting strong, uniform emergence, and balanced growth above and below ground.]

[Boosts root growth + plant vigor. Fortified Stimulate Yield Enhancer® Plus contains the signaling molecules that enable optimal plant functioning, to produce uniform growth and high yielding characteristics.]

[Improves plant health + physiology. Fortified Stimulate Yield Enhancer® Plus improves cell division and nutrient uptake, so your plants can better defend themselves and recover from abiotic stressors such as cold weather, drought and herbicide applications]

[Increases yield potential + grain quality. An EPA-registered yield enhancer, Fortified Stimulate Yield Enhancer® Plus supports hormones within the plant to improve genetic expression.]

[Fortified Stimulate Yield Enhancer® Plus is formulated for the following highlighted applications [seed] [soil] [foliar] [irrigation]]

[Fortified Stimulate Yield Enhancer® Plus improves emergence and germination, while boosting early-stage root growth and plant vigor when applied at planting.]

[Fortified Stimulate Yield Enhancer® Plus is formulated for in-tank compatibility with most industry fertilizers, herbicides, fungicides and insecticides.]

[Irrigation provides an ideal scenario for the application of Fortified Stimulate Yield Enhancer® Plus at key growth stages.]

{Front Panel Start}

FORTIFIED STIMULATE
YIELD ENHANCER® PLUS
[PLANT GROWTH PROMOTER]
 {Sublabel A FOR AGRICULTURAL/COMMERCIAL USE}

ACTIVE INGREDIENTS:

| | |
|------------------------------|--------|
| Cytokinin (as kinetin) | 0.009% |
| Gibberellic Acid | 0.005% |
| Indole-3-butyric Acid | 0.005% |
| Indole-3-acetic Acid | 0.005% |

| | |
|---------------------------------|----------------|
| OTHER INGREDIENTS: | <u>99.976%</u> |
| Total | 100.000% |

This product contains: 0.000095 oz (2.7 mg) of cytokinin, 0.000053 oz (1.5 mg) of gibberellic acid, 0.000053 oz (1.5 mg) of indole-3-butyric acid, and 0.000053 oz (1.5 mg) of indole-3-acetic acid per fluid ounce.

KEEP OUT OF REACH OF CHILDREN
CAUTION

See additional Precautionary Statements, First Aid and Directions for Use [inside booklet] [on [back panel] [side panel] [other panel]].

Z-FORTSTIMYEP

EPA Reg. No.: 57538-49

EPA Est. Nos.: [57538-FL-1,]
[57538-IA-1,] [57538-TX-2]

[DENSITY:]
[8.6 lb/gal or 1.04 kg/L]

NET CONTENTS:
 1 Gal (3.8 L) 2.5 Gal (9.5 L) 5 Gal (19 L) 55 Gal (209 L)

[NET WEIGHT:]
 8.6 lb (3.9 kg) 21.5 lb (9.7kg) 43.0 lb (19.5 kg) 473.0 lb (214.5 kg)]

[*Not for use in California]
[Lot Number:]
[Manufacture Date:]
[Best if used by:]

[Rev: 22J21] [revision code for each label change will go here]

[1.0 Quart (0.95 L) 2.2 lb (1.0 kg)]

[30 Gal (114 L) 258.0 lb (117.0 kg)]

[275 Gal (1045 L) 2365.0 lb (1072.5 kg)]

[Bulk]

[Will replace [™] with © throughout the label when it is a Registered trademark)

[Patents: <https://www.stollerusa.com/about/patents/>]

[ABN: Fortified Stimulate Yield Enhancer]

{End Front Panel}

| FIRST AID | |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| If on skin or clothing | <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice. |
| If swallowed | <ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or a doctor. • Do not give anything by mouth to an unconscious person. |
| If inhaled | <ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice. |
| If in eyes | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. |
| HOT LINE NUMBER | |
| -Have the product container or label with you when calling a poison control center or doctor or are going for treatment. -For general information on product use, call the National Pesticide Information Center at 1-800-858-7378. -For emergencies, call the Poison Control Network at 1-800-222-1222. | |
| FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure or accident, call CHEMTREC at 1-800-424-9300. | |

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if inhaled. Avoid breathing vapor or spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Remove and wash contaminated clothing before reuse. Wear the appropriate Personal Protective Equipment (PPE).

PERSONAL PROTECTIVE EQUIPMENT

Waterproof gloves are sufficiently chemical-resistant for this product. If you want more options, follow the instructions for category A on an Environmental Protection Agency (EPA) chemical-resistance category selection chart.

Applicators and other handlers must wear:

- long-sleeved shirt and long pants,
- chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride, and
- shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If there are no instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4.6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. Remove clothing immediately if pesticide gets inside, then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not apply when weather conditions favor drift from treated areas. Do not apply where runoff is likely to occur. Do not contaminate water when cleaning equipment or disposing of equipment wash water or rinsate. Exposed treated seed may be hazardous to birds and other wildlife. Treat only those seeds needed for immediate use and planting. Do not store excess treated seed beyond planting time. Dispose of all excess treated seed and seed packaging by burial away from streams and bodies of water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard (WPS), 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval (REI). The requirements in this box only apply to uses of this product that are covered by the WPS.

Do not enter or allow worker entry into treated areas during the REI of 4 hours unless wearing appropriate PPE.

Exception: If the product is soil-incorporated, the WPS, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the WPS and that involves contact with anything that has been treated, such as plants, soil, or water is:

- long-sleeved shirt and long pants,
 - chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride,
- and
- shoes plus socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others to enter treated areas until sprays have dried.

CHEMIGATION [*]

Application and Calibration Techniques for Sprinkler Irrigation

Apply this product only through the following types of irrigation systems: center pivot, traveler, big gun, motorized lateral move, end tow, side (wheel) roll, solid set, and hand move irrigation equipment. Do not apply through any other types of irrigation systems. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Experiment Station specialists, equipment manufacturers or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person shall shut the system down and make necessary adjustments should the need arise.

[*Not for Use in California]

A. Center Pivot, Traveler, Big Gun, Motorized Lateral Move, End Tow, and Side (Wheel) Roll

Irrigation Equipment: Operate system and injection equipment at normal pressures recommended by the manufacturer of injection equipment used. Fill tank of injection equipment with water. Operate system for one complete circle for center pivot or one complete run for the other recommended equipment, measuring time required, amount of water injected, and acreage contained in circle or run. Mix recommended amount of Fortified Stimulate Yield Enhancer® Plus for acreage to be covered into same amount of water used during calibration and inject into system continuously for one revolution or run, but continue to operate irrigation system until Fortified Stimulate Yield Enhancer® Plus has been cleared from last sprinkler head. Spray mixture in the chemical supply tank must be continuously agitated, otherwise settling and uneven application may occur.

B. Solid Set and Hand Move Irrigation Equipment: Determine acreage covered by sprinkler. Fill tank of injection equipment with water and adjust flow to use contents over a thirty-to-forty-five-minute period. Mix desired amount of Fortified Stimulate Yield Enhancer® Plus for acreage to be covered into quantity of water used during calibration and operate entire system at normal pressures recommended by the manufacturer of injection equipment used for amount of time established during calibration. Provide constant mechanical agitation in the mix tank to ensure that Fortified Stimulate Yield Enhancer® Plus will remain in suspension during the injection cycle. Fortified Stimulate Yield Enhancer® Plus can be injected at the beginning or end of the irrigation cycle or as a separate application. Stop injection equipment after treatment is completed and continue to operate irrigation system until Fortified Stimulate Yield Enhancer® Plus is cleared from last sprinkler head.

Safety Devices for Sprinkler Chemigation

- (1) The systems designated above must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- (2) All pesticide injection pipelines must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- (3) The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- (4) The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- (5) The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- (6) Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- (7) Do not apply when wind speed favors drift beyond the area intended for treatment.

Systems Connected to Public Water Sources

- (1) Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of a year.
- (2) Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
- (3) The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- (4) The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- (5) The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops or, in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
- (6) Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- (7) Do not apply when wind speed favors drift beyond the area intended for treatment.

Furrow Chemigation[*]

- (1) Systems using a gravity flow pesticide dispensing system must meter the pesticide into the water at the head of the field and downstream of a hydraulic discontinuity such as a drop structure or weir box to decrease potential for water source contamination from backflow if water flow stops.
- (2) Systems utilizing a pressurized water and pesticide injection system must meet the following requirements:
 - a. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
 - b. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
 - c. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
 - d. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
 - e. The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
 - f. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Apply Fortified Stimulate Yield Enhancer® Plus with sufficient water to penetrate into the root zone without excessive leaching into deeper soil.

[*Not for use in California]

Drip (Trickle) Chemigation[*]

- (1) The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- (2) The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- (3) The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- (4) The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- (5) The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- (6) Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Apply Fortified Stimulate Yield Enhancer[®] Plus with sufficient water to penetrate into the root zone without excessive leaching into deeper soil.

[*Not for use in California]

MODE OF ACTION

Fortified Stimulate Yield Enhancer[®] Plus is a plant growth regulator. Fortified Stimulate Yield Enhancer[®] Plus enhances plant growth and development by stimulating cell division, cell differentiation and enlargement, nutrient uptake and nutrient utilization. It is especially effective when applied with foliar fertilizer, but it is also compatible with pesticides.

MIXING INSTRUCTIONS: Fortified Stimulate Yield Enhancer[®] Plus is water soluble and suitable for use in conventional liquid application systems including sprinkler irrigation systems. Shake Fortified Stimulate Yield Enhancer[®] Plus thoroughly and dilute in sufficient water to assure adequate, even coverage without producing excessive runoff. Agitate the spray mixture during application and apply within 12 hours of dilution. Fortified Stimulate Yield Enhancer[®] Plus can be applied tank mixed with most insecticides, fungicides, herbicides and foliar fertilizers but must be the last addition to the spray mixture. Follow the most restrictive of the labeling limitations and precautions of all products used in mixtures.

APPLICATION INSTRUCTIONS

Apply Fortified Stimulate Yield Enhancer[®] Plus (by ground or air) to foliage diluted in 2 to 10 gallons of water per acre (19-94 liters per Hectare). Larger volumes of water may be used if not associated with excessive runoff. For best results, apply in the early morning or late evening. When applying Fortified Stimulate Yield Enhancer[®] Plus in a band or as a foliar-directed spray, reduce the application rate from the labeled broadcast rate in proportion to the percent of the field surface area covered by the foliar spray, but not below the minimum rate listed in the table.

VEGETABLES

| Crop | No. of Applications | Broadcast Spray | | Band Spray | | In Seed Furrow | | Transplant Water | | Spray Timing |
|------------------------------|---------------------|-----------------|-------|------------|-------|----------------|---------|------------------|---------|--------------------------------------------------------------------------------------------------|
| | | Fl oz/A | MI/Ha | Fl oz/A | MI/Ha | Fl oz/A | MI/Ha | Fl oz/A | MI/Ha | |
| Beans[*] | 1 | 8 | 560 | 4 | 280 | | | | | Between the 3rd trifoliate leaf stage and flower bud formation. |
| | 2-3 | 4 | 280 | 3 | 210 | 2-8 | 140-560 | | | Begin at the 3rd trifoliate leaf stage and then at 7-10 day intervals. |
| | 4-6 | 3 | 210 | 2 | 140 | | | | | Begin at the 2nd trifoliate leaf stage and then at 7-14 day intervals. |
| Broccoli[*] | 3 | 6 | 425 | 4 | 280 | 2-8 | 140-560 | 2-8 | 140-560 | Begin at the 4-5 leaf stage and then at 10-14 day intervals. |
| Brussels Sprouts[*] | 3 | 6 | 425 | 4 | 280 | 2-8 | 140-560 | 2-8 | 140-560 | Begin at the 4-5 leaf stage and then at 10-14 day intervals. |
| Cabbage[*] | 3 | 6 | 425 | 4 | 280 | 2-8 | 140-560 | 2-8 | 140-560 | Begin at the 4-5 leaf stage and then at 10-14 day intervals. |
| Cauliflower[*] | 3 | 6 | 425 | 4 | 280 | 2-8 | 140-560 | 2-8 | 140-560 | Begin at the 4-5 leaf stage and then at 10-14 day intervals. |
| Corn[*], Sweet[*] | 1 | 8 | 560 | 5 | 350 | 2-8 | 140-560 | | | 2-6 leaf stage. |
| | 2 or more | 4 | 280 | 3 | 210 | | | | | Begin at 2-6 leaf stage and then at 7-21 day intervals through the end of tasseling. |
| Cucumbers[*] | 1 | 8 | 560 | 4 | 280 | | | | | Between flower bud initiation and first bloom. |
| | 2-3 | 4 | 280 | 3 | 210 | 2-8 | 140-560 | 2-8 | 140-560 | Begin at flower bud initiation and then at 7-10 day intervals. |
| | 3-4 | 4 | 280 | 3 | 210 | | | | | Begin at transplant, or at the 3-4 leaf stage for direct seeded, and then at 7-10 day intervals. |
| Lettuce[*] | 3 | 6 | 425 | 4 | 280 | 2-8 | 140-560 | 2-8 | 140-560 | Begin at the 4-5 leaf stage and then at 10-14 day intervals. |
| Melons[*] | 1 | 8 | 560 | 4 | 280 | | | | | Between flower bud initiation and first bloom. |
| | 2-3 | 4 | 280 | 3 | 210 | 2-8 | 140-560 | 2-8 | 140-560 | Begin at flower bud initiation and then at 7-10 day intervals. |
| | 4-6 | 3 | 210 | 2 | 140 | | | | | Begin 2 weeks after emergence and then at 7-14 day intervals. |
| Onions[*] | 3 | 6 | 425 | 4 | 280 | 2-8 | 140-560 | | | Begin 2 weeks after emergence and then at 10-14 day intervals. |
| Peppers[*] | 4-6 | 4 | 280 | 3 | 210 | 2-8 | 140-560 | 2-8 | 140-560 | Begin at transplant, or at the 3-4 leaf stage for direct seeded, and then at 7-14 day intervals. |
| Potatoes[*] | 1 | 16 | 1,135 | 8 | 560 | 2-8 | 140-560 | | | Tuber initiation. |
| | 3 | 8 | 560 | 4 | 280 | | | | | Begin at stolon formation (8-10 leaf stage) and then at 10-14 day intervals. |
| Squash[*] | 1 | 8 | 560 | 4 | 280 | 2-8 | 140-560 | | | Between flower bud initiation and first bloom. |
| | 2-3 | 4 | 280 | 3 | 210 | | | | | Begin at flower bud initiation and then at 7-10 day intervals. |
| | 4-6 | 3 | 210 | 2 | 140 | | | | | Begin 2 weeks after emergence and then at 7-14 day intervals. |
| Tomatoes[*] (Eggplant[*]) | 1 | 8 | 560 | 4 | 280 | | | | | Between flower bud initiation and first bloom. |
| | 2-3 | 4 | 280 | 3 | 210 | 2-8 | 140-560 | 2-8 | 140-560 | Begin at flower bud initiation and then at 7-10 day intervals. |
| | 4-6 | 3 | 210 | 2 | 140 | | | | | Begin 2 weeks after emergence, or at transplant, and then at 7-14 day intervals. |

[*]Not for Use in California]

FIELD CROPS

| Crop | No. of Applications | Broadcast Spray | | Band Spray | | In Seed Furrow | | Transplant Water | | Spray Timing |
|---------------------------------------------------------------------------------------------------------------------------|---------------------|-----------------|-------|------------|-------|----------------|---------|------------------|---------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| | | Fl oz/A | MI/Ha | Fl oz/A | MI/Ha | Fl oz/A | MI/Ha | Fl oz/A | MI/Ha | |
| Alfalfa[*] | 3-4 | 4 | 280 | 4 | 289 | | | | | Apply after each cutting |
| | 1 | 16 | 1,135 | 8 | 560 | 2-8 | 140-560 | | | 6-8 leaf stage |
| Beets[*], sugar[*] | 2-3 | 8 | 560 | 4 | 280 | | | | | Begin at the 2 leaf stage and then at 7-14 day intervals. |
| Canola[*] | 3 | 6 | 425 | 4 | 280 | 2-8 | 140-560 | | | Begin at the 3-5 leaf stage and then at 10-14 day intervals. |
| Corn[*] | 1 | 8 | 560 | 5 | 350 | 2-8 | 140-560 | | | 2-6 leaf stage. |
| | 1 | 8 | 560 | 4 | 280 | 2-8 | 140-560 | | | Between flower initiation and final bloom. |
| Cotton[*] | 2-3 | 4 | 280 | 3 | 210 | | | | | Early bloom and 7-14 days later. |
| | 3-4 | 4 | 280 | 3 | 210 | | | | | Begin at pinhead square and then at 7-10 day intervals. |
| Peanuts[*] | 4-6 | 4 | 280 | 3 | 210 | 2-8 | 140-560 | | | Begin 30 days after planting and then at 7-14 day intervals. |
| Rice[*] | 1 | 8 | 560 | | | | | | | 2-5 leaf stage or panicle initiation. |
| | 2 | 4 | 280 | | | | | | | 2-5 leaf stage or panicle initiation. |
| Sorghum[*] | 1 | 8 | 560 | 5 | 350 | 2-8 | 140-560 | | | 2-6 leaf stage. |
| Soybeans[*] | 1 | 8 | 560 | 5 | 350 | 2-8 | 140-560 | | | 3-7 trifoliolate leaf stage (V4-V8). |
| | 2 | 4 | 280 | 3 | 210 | | | | | 3-7 trifoliolate leaf stage (V4-V8) and 10-17 days later. |
| Tobacco[*] | 2 | 8 | 560 | 5 | 350 | | | 2-8 | 140-560 | 1 st 40 days after planting 2 nd after topping |
| Tree Nuts[*] (Almonds[*], Cashew[*], Chestnuts[*], Macadamia[*], ,Pecan[*], Pistachio[*], Walnut[*]) | 5 | 4-16 | | | | | | | | 1st application: Petal fall. 2nd application: 4 weeks post petal fall. 3rd application: 8 weeks post petal fall. 4th application: Bud initiation. 5th application: Bud development. |
| Wheat[*], Barley[*], Oats[*], Rye[*] | 1-2 | 8 | 560 | 6 | 180 | 2-8 | 140-560 | | | Start at tillering in the fall and/or spring and when 2 to 3 leaves form on main stem. |

[*Not for Use in California]

SMALL FRUITS, VINES AND TREE FRUITS[*]

Bananas[*]: Apply 4 to 8 fl. oz./A (300 to 600 ml/Ha) in a band around the root mat and repeat every 10 to 14 days for a total of 10 applications.

Grapes[*] (Wine Grape[*], Table Grape[*]: Apply 4 to 16 fl. Oz/A, (300 to 1200 ml/Ha) every two weeks starting at bud break to the beginning of veraison and repeat 2 weeks post-harvest.

Oranges[*]: Apply Fortified Stimulate Yield Enhancer® Plus at 1 to 2 pints/100 gal(126 ml to 252 ml/100 L) at first bloom and repeat at each flush of new growth.

Pineapple[*]: Spray after transplanting with 4 to 8 fl. oz./100 gal (120 to 240 ml/300 L) and repeat every 10 to 14 days or apply in an irrigation system to provide 4 to 8 fl. oz./A (324 to 648 ml/Ha).

Strawberries[*]: Begin sprays at first bloom at 4 fl oz/A (286 ml/Ha) broadcast or 2 fl oz/A (143 ml/Ha) as band sprays directed at the row. Repeat sprays at 2 to 4 week intervals for a total of 3 to 6 sprays.

[*Not for Use in California]

YOUNG TREES AND ORNAMENTALS[*]

Shrubs, Established[*]: For increased vigor, rapid growth and healthy plant appearance, spray foliage with transplant solution of 1 fl. oz. of Fortified Stimulate Yield Enhancer® Plus /2 gal of water (3.9 ml/L) to point of run-off two to three times per year.

Flowering Plants (Roses, Azaleas, etc.)[*]: 3-4 year old--using stock solution of 1 fl. oz. of Fortified Stimulate Yield Enhancer® Plus /2 gal of water (3.9 ml/L) —Take 4 fl. oz. stock solution in 1 gallon (31.25 ml/L) of water and water in around root zone one time per year, preferably early spring.

Sod[*]: To improve growth, heavy rooting, broadcast 8 fl. oz./A (585 ml/Ha).

1. To speed up regrowth after harvest, broadcast 4 fl. oz./A (292 ml/Ha).
2. To boost with another 4 fl. oz./A (292 ml/Ha)., broadcast six weeks later.

Turf[*]: For quick “tie down” after laying and to get turf off to a quick start, use Fortified Stimulate Yield Enhancer® Plus as follows:

1. Broadcast 2 fl. oz./5,000 sq. ft. (0.13 ml/sq m) and water in following application.
2. Second application--30 days later--2 fl. oz./5,000 sq. ft. (0.13 ml/sq m) and water in following application.

[*Not for Use in California]

NOTICE: FORTIFIED STIMULATE YIELD ENHANCER® PLUS IS NOT A FERTILIZER. USE IN COMBINATION WITH A GOOD FERTILIZER PROGRAM WHERE INDICATED.

GOLF COURSES[*]

Greens[*]: Make an initial treatment with Fortified Stimulate Yield Enhancer® Plus to promote root development and protect against “winter kill” using 2 fl. oz./5,000-7,000 sq. ft. (0.13-0.10 ml/sq m). Thereafter, use 1 fl. oz./green (30 ml) every 30 days.

Tees[*]: Use Fortified Stimulate Yield Enhancer® Plus at ½ fl. oz./1,200-1,500 sq. ft. (0.13-0.10 ml/sq m) of tee area every 30 days to maintain a healthy mass root growth.

Fairways[*]: To establish the necessary root growth to fully utilize applied fertilizer, use Fortified Stimulate Yield Enhancer® Plus at 8 fl. oz./A (585 ml/Ha) two times the first year. Thereafter, use 8 fl. oz./A (585 ml/Ha) one to two times a year .

[*Not for Use in California]

TRANSPLANTS[*]

FORTIFIED STIMULATE YIELD ENHANCER® PLUS MAY BE USED WHEN TRANSPLANTING ORNAMENTALS AND YOUNG TREES FOR A FAST AND HEALTHY START.

Use 1 fl. oz./2 gal (3.9 ml/L) water stock solution as follows: At time of transplanting—

1. Bare (naked) roots - dip or spray with stock solution of 1 fl. oz. of Fortified Stimulate Yield Enhancer® Plus/2 gal of water (3.9 ml/L).
2. Balled plants - spray ball at time of transplant.
3. Mist (not run-off) foliage lightly at time of transplant.
4. Using stock solution, apply 1 gallon Fortified Stimulate Yield Enhancer® Plus in furrow/acre (9.4 L/Ha).

[*Not for Use in California]

SEED TREATMENT[*]

Use only on seeds for crops listed elsewhere on the label. Do not use treated seed for food, feed or oil purposes. If this product is intended for commercial seed treatment, treated seed must be labeled in accordance with the requirements of the Federal Seed Act and applicable State seed laws. An approved dye must be added to distinguish treated seed and prevent inadvertent use for food, feed, or oil purposes. If this product is intended for “at planting” use, treat only those seeds needed for immediate use and planting. Do not store excess treated seed beyond planting time. Dispose of excess treated seed by burial away from streams and bodies of water. A dye is not required.

For crops listed: Apply 2 to 4 fluid ounces per 100 lbs. of seed (1.3 to 2.6 ml per kg). Dilute with water and mist spray on seed while mixing. Do not store seed wet as germination can be reduced if not planted soon after treatment.

[*Not for Use in California]

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in a cool place and out of direct sunlight.

Pesticide Disposal: To avoid waste, use all of the material in this container by application according to label directions. If waste cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

Container Handling:

Nonrefillable container. Do not reuse or refill this container

For Nonrefillable containers equal to or less than 5 gallons(19 L).

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

For Nonrefillable containers greater than 5 gallons(19 L)

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY STATEMENT: To the fullest extent permitted by law, Stoller Enterprises, Inc. warrants that the product conforms to the description on the label and is reasonably fit for the purposes set forth on the label, when used according to directions under normal use conditions. Neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to the label instructions; the buyer assumes the risk of any such uses.

**Manufactured By
STOLLER ENTERPRISES, INC.**

9090 Katy Freeway, Suite 400
Houston, Texas 77024, U.S.A.

Toll Free 1-800-539-5283 • Tel.: 1(713) 461-1493

{Front Panel Start}

**FORTIFIED STIMULATE
YIELD ENHANCER® PLUS
[PLANT GROWTH PROMOTER]
{Sublabel B FOR HOMEOWNERS'/GARDEN USE}**

ACTIVE INGREDIENTS:

| | |
|------------------------------|--------|
| Cytokinin (as kinetin) | 0.009% |
| Gibberellic Acid | 0.005% |
| Indole-3-butyric Acid | 0.005% |
| Indole-3-acetic Acid | 0.005% |

| | |
|---------------------------------|----------------|
| OTHER INGREDIENTS: | <u>99.976%</u> |
| Total | 100.000% |

This product contains: 0.000095 oz (2.7 mg) of cytokinin, 0.000053 oz (1.5 mg) of gibberellic acid, 0.000053 oz (1.5 mg) of indole-3-butyric acid, and 0.000053 oz (1.5 mg) of indole-3-acetic acid per fluid ounce.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

**See additional Precautionary Statements, First Aid and Directions for Use [inside booklet] [on
[back panel] [side panel] [other panel].
Z-FORTSTIMYEP**

EPA Reg. No.: 57538-49

EPA Est. Nos.: [57538-FL-1],
[57538-IA-1], [57538-TX-2]

[DENSITY:]
[8.6 lb/gal or 1.04 kg/L]

NET CONTENTS:
 1 Gal (3.8 L) 2.5 Gal (9.5 L) 5 Gal (19 L) 55 Gal (209 L)

[NET WEIGHT:]
 8.6 lb (3.9 kg) 21.5 lb (9.7kg) 43.0 lb (19.5 kg) 473.0 lb (214.5 kg)]

[*Not for use in California]

[Lot Number:]

[Manufacture Date:]

[Best if used by:]

[Rev: 22J21] [revision code for each label change will go here]

[1.0 Quart (0.95 L) 2.2 lb (1.0 kg)]

[30 Gal (114 L) 258.0 lb (117.0 kg)]

[275 Gal (1045 L) 2365.0 lb (1072.5 kg)]

[Bulk]

[Will replace TM with © throughout the label when it is a Registered trademark)

[Patents: <https://www.stollerusa.com/about/patents/>]

[ABN: Fortified Stimulate Yield Enhancer]

| FIRST AID | |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| If on skin or clothing | <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice. |
| If swallowed | <ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or a doctor. • Do not give anything by mouth to an unconscious person. |
| If inhaled | <ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice. |
| If in eyes | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. |
| HOT LINE NUMBER | |
| -Have the product container or label with you when calling a poison control center or doctor or are going for treatment. -For general information on product use, call the National Pesticide Information Center at 1-800-858-7378. -For emergencies, call the Poison Control Network at 1-800-222-1222. | |
| FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure or accident, call CHEMTREC at 1-800-424-9300. | |

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if inhaled. Avoid breathing vapor or spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Remove and wash contaminated clothing before reuse. Wear the appropriate Personal Protective Equipment (PPE).

ENVIRONMENTAL HAZARDS

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

MODE OF ACTION

Fortified Stimulate Yield Enhancer® Plus is a plant growth regulator. Fortified Stimulate Yield Enhancer® Plus enhances plant growth and development by stimulating cell division, cell differentiation and enlargement, nutrient uptake and nutrient utilization. It is especially effective when applied with foliar fertilizer, but it is also compatible with pesticides.

MIXING INSTRUCTIONS: Fortified Stimulate Yield Enhancer® Plus is water soluble and suitable for use in conventional liquid application systems. Shake Fortified Stimulate Yield Enhancer® Plus thoroughly and dilute in sufficient water to assure adequate, even coverage without producing excessive runoff. Agitate the spray mixture during application and apply within 12 hours of dilution. If Fortified Stimulate Yield Enhancer® Plus is tank-mixed with other insecticides, fungicides, herbicides or foliar fertilizers, Fortified Stimulate Yield Enhancer® Plus must be the last addition to the spray mixture. Follow the most restrictive of the labeling limitations and precautions of all products used in mixtures.

APPLICATION INSTRUCTIONS

Apply Fortified Stimulate Yield Enhancer® Plus to foliage diluted in 1 gallon of water per 1000 sq. ft. Larger volumes of water may be used if not associated with excessive runoff. For best results, make early morning or late evening applications.

[*]VEGETABLES

| Crop | No. of Sprays | Amount/Gallon Per 1000 sq. ft. | Timing |
|-------------------------------------------------|---------------|--------------------------------|------------------------------------------------------------------------------------------------------|
| Bean[*] | 3 | 2 teaspoons | Begin at the 3rd trifoliolate leaf and repeat every 7-10 days. |
| Broccoli[*], Brussels Sprout[*], Cauliflower[*] | 3 | 1 tablespoon | Begin at the 4 to 5 leaf-stage followed by 2 more applications at 10-14 day intervals. |
| Sweet Corn[*] | 2 or more | 2 teaspoons | Begin at 2-6 leaf stage and then at 7-21 day intervals through the end of tasseling. |
| Cucumbers, Melon, Squash[*] | 3 | 2 teaspoons | Begin at flower bud initiation and then follow with 2 more sprays at 7-10 day intervals. |
| Lettuce[*] | 3 | 1 tablespoon | Begin at the 4-5 leaf stage and then 2 more sprays at 10-14 day intervals. |
| Onion[*] | 3 | 1 tablespoon | Begin 2 weeks after emergence and 2 more sprays at 10-14 day intervals. |
| Peppers[*] | 4 | 2 teaspoons | Begin at transplant or at the 3 to 4 leaf stage for direct seeded and repeat at 10-14 day intervals. |
| Potatoes[*] | 3 | 1 1/2 tablespoons | Begin at 8-10 leaf stage and then 2 more sprays at 7-10 day intervals. |
| Tomatoes[*] | 3 | 2 teaspoons | Begin at flower bud initiation and then 2 more sprays at 10-14 day intervals. |

[*Not for Use in California]

SMALL FRUITS, VINES AND TREE FRUITS[*]

| Crop | No. of Sprays | Amount/Gallon Per 1000 sq. ft. | Timing |
|-----------------------------------------------------------------------------------------------------|---------------|--------------------------------|------------------------------------------------------------------------------------------|
| Strawberries[*] | 3-6 | 2 teaspoons | Begin sprays at first bloom. Repeat at 2-4 week intervals for for a total of 3-6 sprays. |
| Oranges[*] | 3-6 | 2 tablespoons | Spray to wet foliage at first bloom and repeat at each flush of new growth. |
| Shrubs, Established[*] | 2-3 | 4 tablespoons | Spray foliage to point of run-off 2-3 times a year. |
| Flower Plants (Roses, Azalea etc.) [*] | 1 | 1 teaspoon | In early spring, water in over root zone 1 time per year. |
| Tree Nuts[*] (Almonds[*], Cashew[*], Chestnuts[*], Macadamia[*], Pecan[*], Pistachio[*], Walnut[*]) | 3-6 | 2 tablespoons | Spray to wet foliage at first bloom and repeat at each flush of new growth. |

Turf[*]: To improve growth and heavy rooting, spray broadcast at 1 1/2 tablespoon per gallon per 1000 sq. ft. To speed up regrowth after cutting, spray broadcast 2 teaspoons per gallon of water per 1000 sq. ft.

Sod[*]: For quick “tie down” after laying and to get turf off to a quick start, use as follows:

1. Broadcast 2 fl. oz./5000 sq/ ft. and water in.
2. Make second application 30 days later and water in.

[*Not for Use in California]

[*]Transplants: Prepare stock solution with 4 tablespoons of Fortified Stimulate Yield Enhancer® Plus per 1 gallon of water. When transplanting with bare roots, dip in stock solution before planting. When transplanting with balled plants, spray ball at time of planting with stock solution. Mist foliage at time of transplant.

[*Not for Use in California]

NOTICE: FORTIFIED STIMULATE YIELD ENHANCER® PLUS IS NOT A FERTILIZER. USE IN COMBINATION WITH A GOOD FERTILIZER PROGRAM WHERE INDICATED.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store unused product in original container only in a cool, dry area out of reach of children and animals. If container is damaged, place the container in a plastic bag. Do not store in direct sunlight. Avoid freezing temperatures.

Pesticide Disposal: If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

Container Disposal: Nonrefillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available.

WARRANTY STATEMENT: To the fullest extent permitted by law, Stoller Enterprises, Inc. warrants that the product conforms to the description on the label and is reasonably fit for the purposes set forth on the label, when used according to directions under normal use conditions. Neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to the label instructions; the buyer assumes the risk of any such uses.

Manufactured By
STOLLER ENTERPRISES, INC.
9090 Katy Freeway, Suite 400
Houston, Texas 77024, U.S.A.
Toll Free 1-800-539-5283 • Tel.: 1(713) 461-1493