

Phytar 560

herbicide

Sodium Cacodylate
Liquid Plus
Surfactant

For General
Post-Emergent
Weed Control


ACTIVE INGREDIENTS:
Sodium Cacodylate . . . 22.73%
Dimethylarsinic Acid
(Cacodylic Acid) . . . 3.88%

INERT INGREDIENTS: 73.39%
Total Arsenic (as elemental)
in water soluble
form 12.75%

Product contains 2.48 lbs.
cacodylic acid equivalent
per gallon.

CAUTION:
Keep out of the
reach of children

Read entire label
before using
this product



ACCEPTED

JUL 6 1973

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND REPLICABLE ACT
FOR ECONOMIC POISON REGISTERED
AND UNDER NO. 39511-105
IN ATTACHED COMMENTS.

*comments in
letter
transf. to
39511-105*

SPECIMEN LABEL

CAUTION: KEEP OUT OF THE REACH OF CHILDREN

CAUTION: Harmful if swallowed. Avoid contact with skin. Avoid breathing spray mist. Wash hands after using. Avoid storage near feed or food products. Keep children and domestic animals off treated areas until this material has been washed into the soil.
Do not contaminate waters used for domestic consumption, or by animals, wildlife and aquatic life, or for irrigation purposes. Do not graze treated areas to livestock.

ANTIDOTE: If taken internally, induce vomiting and call physician at once.
READ ENTIRE LABEL BEFORE USING THIS PRODUCT.

WARRANTY - CONDITION OF SALE; DIRECTIONS FOR USE of this product are based on field use and tests believed reliable and should be followed carefully. It is however impossible to eliminate all risks associated with use of this product. Because such factors as weather conditions, foreign material and manner of use for application are all beyond the control of The Ansul Company or the Seller of this product, such things as crop injury, ineffectiveness or other unintended consequences may result. **ALL SUCH RISKS ARE ASSUMED BY THE BUYER.**

Ansul warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the directions for use as modified by the above. Ansul makes no other warranties, express or implied, including **FITNESS or MERCHANTABILITY.** In no case shall Ansul or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. The foregoing is a condition of sale by The Ansul Company and is accepted as such by the Buyer.

GENERAL INFORMATION: PHYTAR 560 Herbicide is useful for general post-emergent weed control. It is non-selective in its action. Its phytotoxic properties are directly related to its contact with soil. It contains a surfactant. It is unnecessary to add any other surfactant to the spray solution. Best results are obtained on young actively growing weeds. It prevents seed germination, so repeat applications are required for a long-term control of perennials.

MIXING INSTRUCTIONS: PHYTAR 560 Herbicide is compatible with water. Any spray equipment that gives good coverage may be used. Fill the spray tank with water and add the required amount of herbicide with agitation. Finish by adding water and spray. After use, clean equipment thoroughly by flushing with water. Do not store spray solution in tank for application. Although PHYTAR 560 Herbicide is only moderately corrosive, do not use in galvanized or aluminum equipment.

CONTAINER DISPOSAL: Do not reuse empty container. Wash thoroughly with water and detergent, crush if possible, and discard in a safe place.

DIRECTIONS FOR USE:

NON-CROP: PHYTAR 560 Herbicide is useful for general post-emergent control of weeds, grasses and annual crops, along roadsides, driveways and fences, around buildings and ornamentals. Apply similar non-crop areas. Mix 1 to 2 quarts of PHYTAR 560 Herbicide in 100 gallons of water. Spray unwanted vegetation to just short of runoff. If regrowth occurs repeat, as required.

Do not allow spray to come in contact with foliage, green bark, grafted unions, or scuffed, damaged or broken bark when spraying around trees and ornamentals.

AGRICULTURAL PLANTINGS - CITRUS: Except Florida, PHYTAR 560 Herbicide is not used as a directed spray on citrus trees or citrus groves. It is not recommended for use on citrus trees or citrus groves. It should be applied at the rate of 1 to 2 quarts per acre.

Mix 1 to 2 quarts of PHYTAR 560 Herbicide in 100 gallons of water. Apply as directed spray. Do not allow spray to come in contact with foliage, green bark, grafted unions, or scuffed, damaged or broken bark when spraying around trees and ornamentals.

Do not allow spray solution to contact leaves, stems or bark of established trees, shrubs, vines, or young trees or young trees. Do not apply around trees from which fruit will be harvested within one year of treatment.

LAWN RENOVATION: Mow lawn to about 1 inch height before treatment. Mix 1 to 2 quarts of PHYTAR 560 Herbicide in 4 gallons of water and apply to 1 to 2 sq. ft. Spray large lawn thoroughly, preferably in a warm sunny day, foliage will usually turn brown in about five days. Before mowing and other deep-rooted perennials, or partially sprayed foliage, may require a second treatment. Re-apply five days later if green areas remain.

When foliage is all brown, matted areas should be raked to remove dead vegetation. The lawn may then be reestablished according to local practice.

Keep children and pets off treated areas until after first rain or sprinkling following treatment. Do not track from treated to untreated areas.

PHYTAR 560 Herbicide is manufactured by The Ansul Company, Marinette, Wisconsin 54143.
U.S. Patents 3,173,937 and 3,056,668 and others pending.

X. II. PHYTAR and ANSUL are registered trademarks of The Ansul Company.

Net Contents 5 Gallons
Form No.

FROM LABEL FILE 4663-41

THE ANSUL COMPANY,
MARINETTE, WISCONSIN
BEST AVAILABLE COPY