300/ 5-50598

May 23, 1996

RJ Advantage, Inc. c/o ERM Program Management Company 7926 Jones Branch Drive McLean, VA 22102

Attention: Ms. Cathy Shea

Subject: ReJeX-iT AG-145

EPA Reg. No. 58035-13

Your letter of May 22, 1996

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable provided that you make the following changes and submit one (1) copy to us before your ship product:

1. In the STORAGE AND DISPOSAL instructions, change "Avian Aversive Agent", a term not defined on the label, to "Product".

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

Robert A. Forrest

Product Manager (14)

Insecticide-Rodenticide Branch Registration Division (H7504C)

Enclosures: 1. Stamped label

Peacock WP#9:A:\MA\58035-13.MAY:305-5407,-6600:5/23/96

PRECAUTIONARY STATEMENT

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark.

STORAGE & DISPOSAL

STORAGE: Store only in original container, in a dry place inaccessible to children, pets and domestic animals. Store apart from other pesticides. fertilizers, food or feed that may cause cross-contamination from odor.

Store above 32°F. Keep container closed when not in use.

REPELLENT DISPOSAL: Avian Aversion Agent or rinse water that cannot be used according to label instructions must be disposed of according to Federal or approved State procedures. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse. Then offer for recycling or reconditioning, or puncture and dispose in a sanitary landfill, or by other procedure approved by State and local authorities.

ReJeX-iT® AG-145

for agricultural use

ACTIVE INGREDIENT:

Methyl Anthranilate (MA), methyl 2-aminobenzoate	14.5%
INERT INGREDIENTS	85.5%
TOTAL	100.0%


This product contains 1.23 pounds of Methyl Anthranilate per gallon of product.

ReJeX-iT® AG-145 has been formulated from Food Grade ingredients with an active ingredient that exceeds US Food Chemical Codex (FCC) specifications.

KEEP OUT OF REACH OF CHILDREN

CAUTIO MCCEPTED COMMENTS in EPA Letter Dated

MAY 2 3 1996 See left panel for additional precautionary statements.



RJ Advantage, Inc.

A Subsidiary of PMC Specialties Group, Inc. 501 Murray Road, Cincinnati, Ohio 45217

NET WEIGHT: 21 lb - NET CONTENT: 2.5 gallons

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

COMPATIBILITY WITH OTHER INGREDIENTS: Compatibility with other adjuvants has not been tested. If adding other adjuvants or products, test for compatibility first or ask your field man for advice. Do not mix with dry formulations such as wettable powder formulations which may result in residue on the fruit.

CHERRIES

USE RESTRICTIONS: This product may be used to limit damage to ripening cherries caused by American robins (Turdus migratorius), European startings (Sturnus vulgaris), mockingbirds (Mimus polyglottos), sparrows (Spizella, Melospiza, and Zonotrichia spp.), gray cathirds (Dumetella carolinensis), cardinals (Cardinalis cardinalis), American goldfinches (Carduelis tristis), house finches (Carpodacos mexicanus), and gulla (Larus app.). This product must be applied using appropriate spray equipment, such as a pump sprayer or a power-blast sprayer. Do not apply when fruit is wet or rain is expected. Withhold treatment until fruit is susceptible to bird damage.

DILUTION DIRECTIONS: Shake or stir ReJeX-iT® AG-145 container well prior to diluting. For aerial application, mix AG-145 with water at a ratio of 1 part product to 1-3 parts water. For aerial applications (1:3 dilution with water), apply at rates of 10-20 gallons of spray maiture per acre. For ground application, mix AG-145 with water at a ratio of 1 part product to 10-20 parts water. For ground applications of 1:10 dilutions with water, apply at rates of 27.5-55 gallons of sprau mxiture. At 1:20 dilutions, apply at rates of 52.5-105 callons of spray maiture per acre. Mix product outside or in ventilated areas.

APPLICATION DIRECTIONS: Apply spray mixture so that fruit receives thorough coverage. Apply at a rate of 2.5-5 gallons AG-145 (3.1-6.2 lbs a.i.) per acre. Repeat treatment within 7 days if bird activity persists. The product may be applied up to the day of harvest.

WARRANTY STATEMENT

NOTICE: The manufacturer warrants that this product conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this product is fit for its intended purpose. Since timing, method of application, weather and ground conditions, mixture with other chemicals, and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions, or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

95/05/21