



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

305/17 5-515605

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

February 28, 1997

RJ Advantage, Inc.
c/o 6526 Fairlawn Drive
McLean, VA 22101

Attention: Ms Cathy Shea

Subject: ReJeX-iT AG-145
EPA Registration No. 58035-13
Your submission of November 26, 1996

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable provided that you make the following changes and submit one (1) copy to us before you ship your product:

The "DIRECTIONS FOR USE" on the 3-fruit label submitted on November 26, 1996, are acceptable provided that the changes indicated below are made to them.

1. Correct the pluralized common name used for *Toxostoma rufum* to "brown thrashers" in the "USE RESTRICTIONS:" paragraph for "BLUEBERRIES AND GRAPES".
2. Move the text indicated below from its present location in the "DILUTION DIRECTIONS" for "BLUEBERRIES AND GRAPES" so that it appears between the current second and third sentences of the "APPLICATION DIRECTIONS:" paragraph for "BLUEBERRIES AND GRAPES".

"For aerial applications (1:3 dilution with water), apply at rates of 10-20 gallons of spray mixture per acre. For ground applications of 1:10 dilutions with water, apply at rates of 27.5-55 gallons of spray mixture per acre. At 1:20 dilutions, apply at rates of 52.5-105 gallons of spray mixture per acre."

BEST AVAILABLE COPY

Be sure to retain the sentence "Mix product outside or in a ventilated area" in the "DILUTION DIRECTIONS".


- 3. In the use directions pertaining to "CHERRIES", make the same relocation of text that is described under item "2." above. Note that making this change will put the text in the location that we prescribed for it in our registration acceptance letter of May 21, 1996.

As we have noted previously, the "AG-145" portion of the proposed product name reflects the concentration of active ingredient in this product times 10. "AG-14" would be a more appropriate and less misleading product name.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,



William W. Jacobs, PhD
Product Manager (14)
Insecticide-Rodenticide Branch
Registration Division (H7504C)

- Enclosures:
- 1. Stamped Label
 - 2. A-79 Enclosure

Peacock:11:A:\MA\58035-13.FEB:305-5407/-6600:2/28/97

393

PRECAUTIONARY STATEMENT

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark.

STORAGE & DISPOSAL

STORAGE: Store only in original container, in a dry place inaccessible to children, pets and domestic animals. Store apart from other pesticides, fertilizers, food or feed that may cause cross-contamination from odor.

Store above 32° F. Keep container closed when not in use.

REPELLENT DISPOSAL: Product or rinse water that cannot be used according to label instructions must be disposed of according to Federal or approved State procedures. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse. Then offer for recycling or reconditioning, or puncture and dispose in a sanitary landfill, or by other procedures approved by State and local authorities.

ReJeX-iT® AG-145 for agricultural use

ACTIVE INGREDIENT:

Methyl Anthranilate (MA), methyl 2-aminobenzoate 14.5%

INERT INGREDIENTS 85.5%

TOTAL 100.0%

This product contains 1.23 pounds of Methyl Anthranilate per gallon of product.

ReJeX-iT® AG-145 has been formulated from Food Grade Ingredients with an active ingredient that exceeds US Food Chemical Codex (FCC) specifications.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See Left Panel for additional precautionary statements.

Manufactured for
RJ Advantage, Inc.

A Subsidiary of PMC Specialties Group, Inc.
501 Murray Road, Cincinnati, Ohio 45217
EPA EST. No. 68607-OH-001 EPA Reg. No. 58035-13

NET WEIGHT / CONTENT: 2.5 gallons
with COMMENTS
in EPA Letter Dated

FEB 28 1997

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, the pesticide registered under EPA Reg. No.

58035-13

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

COMPATIBILITY WITH OTHER INGREDIENTS: Compatibility with other adjuvants has not been tested. If adding other adjuvants or products, test for compatibility first or ask your field man for advice. Do not mix with dry formulations such as wettable powder formulations which may result in residue on the fruit.

CHERRIES

USE RESTRICTIONS: This product may be used to limit damage to ripening cherries caused by American robins (*Turdus migratorius*), European starlings (*Sturnus vulgaris*), mockingbirds (*Mimus polyglottos*), sparrows (*Spizella, Melospiza, and Zonotrichia* spp.), gray catbirds (*Dumetella carolinensis*), cardinals (*Cardinalis cardinalis*), American goldfinches (*Carduelis tristis*), house finches (*Carpodacus mexicanus*), and gulls (*Larus* spp.). This product must be applied using appropriate spray equipment, such as a pump sprayer or a power-blast sprayer. Do not apply when fruit is wet or rain is expected. Withhold treatment until fruit is susceptible to bird damage.

DILUTION DIRECTIONS: Shake or stir ReJeX-iT® AG-145 container well prior to diluting. For aerial application, mix AG-145 with water at a ratio of 1 part product to 1-3 parts water. For aerial applications (1:3 dilution with water), apply at rates of 10-20 gallons of spray mixture per acre. For ground application, mix AG-145 with water at a ratio of 1 part product to 10-20 parts water. For ground applications of 1:10 dilution with water, apply at rates of 27.5-55 gallons of spray mixture. At 1:20 dilution, apply at rates of 52.5-105 gallons of spray mixture per acre. Mix product outside or in ventilated areas.

APPLICATION DIRECTIONS: Apply spray mixture so that fruit receives thorough coverage. Apply at a rate of 2.5-5 gallons AG-145 (3.1-6.2 lbs a.i.) per acre. Repeat treatment if significant bird damage begins to occur. This product may be applied up to the day of harvest.

BLUEBERRIES AND GRAPES

USE RESTRICTIONS: This product may be used to limit damage to ripening blueberries and grapes caused by American robins (*Turdus migratorius*), cedar waxwings (*Bombicilla cedrorum*), European starlings (*Sturnus vulgaris*), common grackles (*Quiscalus quiscula*), brown thrushes (*Toxostoma rufum*), mockingbirds (*Mimus polyglottos*), mourning doves (*Zenaidura macroura*), cardinals (*Cardinalis cardinalis*), blackbirds (*Icterus* spp.), house finches (*Carpodacus mexicanus*), and gulls (*Larus* spp.). This product must be applied using appropriate spray equipment, such as a pump sprayer or a power-blast sprayer. Do not apply when fruit is wet or rain is expected. Withhold treatment until fruit is susceptible to bird damage.

DILUTION DIRECTIONS: Shake or stir ReJeX-iT® AG-145 container well prior to diluting. For better coverage, add a super wetter (such as Silwet L-77® from Loveland Industries, Inc.) as recommended on the label. For aerial application, mix AG-145 with water at a ratio of 1 part product to 1-3 parts water. For aerial applications (1:3 dilution with water), apply at rates of 10-20 gallons of spray mixture per acre. For ground application, mix AG-145 with water at a ratio of 1 part product to 10-20 parts water. For ground applications of 1:10 dilution with water, apply at rates of 27.5-55 gallons of spray mixture. At 1:20 dilution, apply at rates of 52.5-105 gallons of spray mixture per acre. Mix product outside or in ventilated areas.

APPLICATION DIRECTIONS: Apply spray mixture so that fruit receives thorough coverage. Apply at a rate of 2.5-5 gallons AG-145 (3.1-6.2 lbs a.i.) per acre. Repeat treatment if significant bird damage begins to occur. This product may be applied up to the day of harvest.

WARRANTY STATEMENT

NOTICE: The manufacturer warrants that this product conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this product is fit for its intended purpose. Since timing, method of application, weather and ground conditions, mixture with other chemicals, and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions, or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

11/26/96

BEST AVAILABLE COPY