

PP 04

58035-13

8/11/98

Page 1 of 4

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

AUG 11 1998

Catherine B. Rice
Manager, Regulatory Affairs
RJ Advantage Wildlife Management
501 Murray Road
Cincinnati, Ohio 45217-1014

Dear Ms. Rice:

Subject: Revised labeling, Master Label
ReJeX-iT AG-145
EPA Reg. No. 58035-13
Your submissions dated May 15, June 12, June 17, and June 29, 1998

Revised labeling referred to above, and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, to develop a master label and include homeowner use directions, is acceptable, provided you make the label changes listed below:

1. Add the general statement, "Do not contaminate water, food, or feed by storage and disposal" at the top of the Storage and disposal paragraph.
2. Delete the statement "for agricultural use" from the front panel, right below the product name. Since you have converted this into a master label, this statement is not appropriate. You may modify this statement to reflect both agricultural and homeowner use sites.
3. Delete the prohibition for aerial application in New York from the homeowner use section.

CONCURRENCES

SYMBOL	Kumars	8/11/98						
SURNAME	7511C	8/11/98						
DATE	8-5-98	8-5-98						


584

2

Submit five copies of the final printed label incorporating the changes listed above, before you release the product for shipment bearing the amended labeling.

A stamped copy of the label is enclosed for your records. Any questions regarding this correspondence should be directed to Rita Kumar at (703) 308-8291.

Sincerely yours,



Janet L. Andersen, Ph.D.
Director
Biopesticides and Pollution
Prevention Division (7511C)

Enclosure

PRECAUTIONARY STATEMENT

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark.

STORAGE & DISPOSAL

STORAGE: Store only in original container, in a dry place inaccessible to children, pets and domestic animals. Store apart from other pesticides, fertilizers, food or feed that may cause cross-contamination from odor.

Store above 32° F. Keep container closed when not in use.

REPELLENT DISPOSAL: Product or rinse water that cannot be used according to label instructions must be disposed of according to Federal or approved State procedures. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse. Then offer for recycling or reconditioning, or puncture and dispose in a sanitary landfill, or by other procedures approved by State and local authorities.

MASTER LABEL

This master label bears directions for both agricultural/commercial and homeowner use with various application rates specific to the associated container sizes.

ReJeX-iT® AG-145

for agricultural use

ACTIVE INGREDIENT:

Methyl Anthranilate (MA),
methyl 2-aminobenzoate 14.5%

INERT INGREDIENTS 85.5%

TOTAL 100.0%

This product contains 1.23 pounds/0.31 pounds/0.15 pounds of Methyl Anthranilate per gallon/quart/pint of product.

ReJeX-iT® AG-145 has been formulated from Food Grade Ingredients with an active ingredient that exceeds US Food Chemical Codex (FCC) specifications.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Left Panel for additional precautionary statements.

Manufactured for

RJ Advantage, Inc.

501 Murray Road, Cincinnati, Ohio 45217

EPA EST. No. 71638-OH-001 EPA REG. No. 58035-13

NET WEIGHT: _____ - CONTENT: _____

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

COMPATIBILITY WITH OTHER INGREDIENTS: Compatibility with other adjuvants has not been tested. If adding other adjuvants or products, test for compatibility first or ask your field man for advice. Do not mix with dry formulations such as wettable powder formulations which may result in residue on the fruit.

AGRICULTURAL/COMMERCIAL USE

CHERRIES

USE RESTRICTIONS: This product may be used to limit damage to ripening cherries caused by American robins (*Turdus migratorius*), European starlings (*Sturnus vulgaris*), mockingbirds (*Mimus polyglottos*), sparrows (*Spizella*, *Melospiza*, and *Zonotrichia* spp.), gray catbirds (*Dumetella carolinensis*), cardinals (*Cardinalis cardinalis*), American goldfinches (*Carduelis tristis*), house finches (*Carpodacus mexicanus*), and gulls (*Larus* spp.). This product must be applied using appropriate spray equipment, such as a pump sprayer or a power-blast sprayer. Do not apply when fruit is wet or rain is expected. Withhold treatment until fruit is susceptible to bird damage.

[New York only - Aerial application is prohibited.]

DILUTION DIRECTIONS: Shake or stir ReJeX-iT® AG-145 container well prior to diluting. For aerial application, mix AG-145 with water at a ratio of 1 part product to 1-3 parts water. For ground application, mix AG-145 with water at a ratio of 1 part product to 10-20 parts water. Mix product outside or in ventilated areas.

APPLICATION DIRECTIONS: Apply spray mixture so that fruit receives thorough coverage. Apply at a rate of 2.5-5 gallons AG-145 (3.1-6.2 lbs a.i.) per acre. For aerial applications (1:3 dilutions with water), apply at rates of 10-20 gallons of spray mixture per acre. For ground applications of 1:10 dilutions with water, apply at rates of 27.5-55 gallons of spray mixture. At 1:20 dilutions, apply at rates of 52.5-105 gallons of spray mixture per acre. Repeat treatment if significant bird damage begins to occur. This product may be applied up to the day of harvest.

BLUEBERRIES AND GRAPES

USE RESTRICTIONS: This product may be used to limit damage to ripening blueberries and grapes caused by American robins (*Turdus migratorius*), cedar waxwings (*Bombus cedrorum*), European starlings (*Sturnus vulgaris*), common grackles (*Quiscalus quiscula*), brown thrashers (*Toxostoma rufum*), mockingbirds (*Mimus polyglottos*), mourning doves (*Zenaidura macroura*), cardinals (*Cardinalis cardinalis*), blackbirds (*Icteridae*), house finches (*Carpodacus mexicanus*), and gulls (*Larus* spp.). This product must be applied using appropriate spray equipment, such as a pump sprayer or a power-blast sprayer. Do not apply when fruit is wet or rain is expected. Withhold treatment until fruit is susceptible to bird damage.

[New York only - Aerial application is prohibited. This product is not to be used on grapes within Suffolk County.]

DILUTION DIRECTIONS: Shake or stir ReJeX-iT® AG-145 container well prior to diluting. For better coverage, add a super wetter (such as Silwet L-77® from Loveland Industries, Inc.) as recommended on the label. For aerial application, mix AG-145 with water at a ratio of 1 part product to 1-3 parts water. For ground application, mix AG-145 with water at a ratio of 1 part product to 10-20 parts water. Mix product outside or in ventilated areas.

APPLICATION DIRECTIONS: Apply spray mixture so that fruit receives thorough coverage. Apply at a rate of 2.5-5 gallons AG-145 (3.1-6.2 lbs a.i.) per acre. For aerial applications (1:3 dilutions with water), apply at rates of 10-20 gallons of spray mixture per acre. For ground applications of 1:10 dilutions with water, apply at rates of 27.5-55 gallons of spray mixture. At 1:20 dilutions, apply at rates of 52.5-105 gallons of spray mixture per acre. Repeat treatment if significant bird damage begins to occur. This product may be applied up to the day of harvest.

ACCEPTED
with COMMENTS
In EPA Letter Dated

AUG 11 1998

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

58035-13

3
4
E

HOMEOWNER USE

CHERRIES

USE RESTRICTIONS: This product may be used to limit damage to ripening cherries caused by American robins (*Turdus migratorius*), European starlings (*Stumus vulgaris*), mockingbirds (*Mimus polyglottos*), sparrows (*Spizella*, *Melospiza*, and *Zonotrichia* spp.), gray catbirds (*Dumetella carolinensis*), cardinals (*Cardinalis cardinalis*), American goldfinches (*Carduelis tristis*), house finches (*Carpodacus mexicanus*), and gulls (*Larus* spp.). This product must be applied using appropriate spray equipment, such as a pump sprayer. Do not apply when fruit is wet or rain is expected. Withhold treatment until fruit is susceptible to bird damage.

[New York only - Aerial application is prohibited.]

DILUTION DIRECTIONS: Shake or stir ReJoX-IT® AG-145 container well prior to diluting. Mix AG-145 with water at a ratio of 1 part product to 10-20 parts water (for example, mix 1 ounce of AG-145 with 2 cups of water to make a spray mixture of 1 part product and 16 parts water). Mix product outside or in ventilated areas.

APPLICATION DIRECTIONS: Apply spray mixture so that fruit receives thorough coverage. To cover 1000 square feet, 1 ounce of AG-145 with the appropriate dilution of water is required. Repeat treatment if significant bird damage begins to occur. This product may be applied up to the day of harvest.

BLUEBERRIES AND GRAPES

USE RESTRICTIONS: This product may be used to limit damage to ripening blueberries and grapes caused by American robins (*Turdus migratorius*), cedar waxwings (*Bombus cedrorum*), European starlings (*Stumus vulgaris*), common grackles (*Quiscalus quiscula*), brown thrashers (*Toxostoma rufum*), mockingbirds (*Mimus polyglottos*), mourning doves (*Zenaidura macroura*), cardinals (*Cardinalis cardinalis*), blackbirds (*Icteridae*), house finches (*Carpodacus mexicanus*), and gulls (*Larus* spp.). This product must be applied using appropriate spray equipment such as a pump sprayer. Do not apply when fruit is wet or rain is expected. Withhold treatment until fruit is susceptible to bird damage.

[New York only - Aerial application is prohibited. This product is not to be used on grapes within Suffolk County.]

DILUTION DIRECTIONS: Shake or stir ReJoX-IT® AG-145 container well prior to diluting. For better coverage, add a super wetter (such as Silwet L-77® from Loveland Industries, Inc.) as recommended on the label. Mix AG-145 with water at a ratio of 1 part product to 10-20 parts water (for example, mix 1 ounce of AG-145 with 2 cups of water to make a spray mixture of 1 part product and 16 parts water). Mix product outside or in ventilated areas.

APPLICATION DIRECTIONS: Apply spray mixture so that fruit receives thorough coverage. To cover 1000 square feet, 1 ounce of AG-145 with the appropriate dilution of water is required. Repeat treatment if significant bird damage begins to occur. This product may be applied up to the day of harvest.

WARRANTY STATEMENT

NOTICE: The manufacturer warrants that this product conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this product is fit for its intended purpose. Since timing, method of application, weather and ground conditions, mixture with other chemicals, and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions, or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

Submitted 6/17/98

h to A