

# FUNGO® 50

## SYSTEMIC TURF FUNGICIDE

AUG 1985

### ACTIVE INGREDIENT:

Dimethyl (1,2-phenylene)bis (iminocarbonothioyl)bis (carbamate) . . . . . 50%

INERT INGREDIENT: . . . . . 50%

TOTAL . . . . . 100%

KEEP OUT OF REACH OF CHILDREN

# CAUTION

STATEMENT OF PRACTICAL TREATMENT

**IF IN EYES,** Flush eyes with water for 15 minutes. Get medical attention.

**IF INHALED,** Remove victim to fresh air. Apply artificial respiration if indicated.

**IF ON SKIN,** Wash off with soap and water. Remove contaminated clothing, and wash before reuse.

**IF SWALLOWED,** Induce vomiting immediately by giving two glasses of water and sticking finger down throat. Never give anything by mouth to an unconscious person. **OBTAIN MEDICAL ATTENTION IMMEDIATELY.**

SEE BACK PANEL FOR  
ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 58185-9

EPA EST. NO. 228-IL-1

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

In case of contact, immediately flush eyes or skin with water. Get medical attention if irritation persists. May be harmful if swallowed or inhaled. Avoid breathing dust or spray mist. Wash thoroughly with soap and water after handling. Wash contaminated clothing before reuse. Do not contaminate feed or food. Keep children and pets off treated area until dry or material has washed into the soil. Always wear protective clothing and respirator when mixing and applying pesticides.

#### ENVIRONMENTAL HAZARDS

**THIS PRODUCT IS TOXIC TO FISH.** Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

#### DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL (USA) LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. Add recommended amount to a partially filled spray tank under agitation. Complete filling to the desired volume. Keep agitated to insure even suspension of material. Make fresh daily. Spray uniformly over the area with a boom type or other power sprayer. Apply after mowing or avoid mowing for 12 hours after application. Apply in sufficient water to obtain thorough coverage, usually 5 gal per 1000 sq ft of turf area. Always treat aprons and approaches to golf greens. Under conditions of severe disease, apply every 5 to 7 days. Note to user: Do not apply to areas likely to be grazed by livestock and do not feed clippings to livestock or poultry.

FUNGO® 50 is a wettable powder for use on golf course greens, tees, lawn and lawns of Bentgrass, Bluegrass, Fescue, Bermudagrass, Ryegrass, St. Augustine grass or their mixtures to be used for the prevention and control of the below mentioned diseases. NOTE: For the following diseases, the spray should dry on the leaves, with no "watering in."

**ANTHRACNOSE** (*Colletotricum graminicola*), **DOLLAR SPOT** (*Sclerotinia homoeocarpa*) - Apply at 1 oz per 1000 sq ft beginning when the disease first appears and continue at 10-14 day intervals as needed. As a curative where preventive schedule is not followed, use 2 oz per 1000 sq ft.

**COPPER SPOT** (*Gloeocercospora sorghi*), **LARGE BROWN PATCH** (*Rhizoctonia solani*), **RED THREAD** (*Corticium fuckelii*), **FUSARIUM PATCH** (*Fusarium nivale*) - Apply at 2 oz per 1000 sq ft beginning when the disease first appears and continue at 10-14 day intervals. Re treat at 5 to 7 day intervals under severe disease conditions.

NOTE: For the following diseases "watering in" is required

**FUSARIUM BLIGHT** (*Fusarium roseum*), **NECROTIC RING SPOT** (*Leptophtheria korrae*), and **SUNDER PATCH** (*Phialophora graminicola*)

Apply at 4-8 oz per 1000 sq ft. Water into root zone by irrigating thoroughly immediately after application. Apply two applications at 10-14 day intervals beginning application when the disease first appears.

**STRIPE SMUT** (*Ustilago striiformis*) - Apply at 4-8 oz per 1000 sq ft. Water into the root zone thoroughly immediately after application. Apply in October or early spring just before grass begins growth.

### STORAGE AND DISPOSAL

**STORAGE:** Store in a dry place in a well-closed container. Keep away from fire or sparks. Do not allow to become wet or overheated in storage. This may cause chemical changes which will impair the fungicidal effectiveness of FUNGO® 50.

**PROHIBITIONS:** Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

**PESTICIDE DISPOSAL:** Pesticide spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, State, or local procedures.

**CONTAINER DISPOSAL:** Completely empty bag by shaking and tipping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities.

### WARRANTY

This material conforms to the description of this label and is reasonably fit for the purposes referred to in the directions for use. Timing and method of application, as well as weathering practices, nature of soil, the disease problem, condition of the crop, irrigation, and other chemical or other factors not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage, or handling of this material and in strict accordance with directions given hereon. THE BUYER ASSUMES ALL RISK OF LOSS OF WARRANTY OF FITNESS OR MERCHANTABILITY IS MADE.

FUNGO® 50 is a registered trademark of Sierra Crop Protection Company for its brand of Fungicide products.

**BEST AVAILABLE COPY**

081

2 LB.  
(0.9 Kg)

Manufactured For

**SIERRA**

CROP PROTECTION COMPANY  
3001 Yosemite Drive, Milpitas, CA 95045