

OCT 22 1993

Mr. John A. Francis, Jr.
Grace-Sierra Crop Protection Co.
1001 Yosemite Drive
Milpitas, CA 95035


Dear Mr. Francis:

Subject: Banrot Broad Spectrum Fungicide 40% Wettable Powder
EPA Reg. No. 58185-10
Your Submission of September 27, 1993

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

This acceptance of your label does not relieve you of any obligation to comply with the Worker Protection Standard (WPS). Under the WPS labeling regulations at 40 CFR Part 158, subpart K, section 156.200(c)(3), you are prohibited from distributing or selling any product within the scope of the WPS requirements after April 21, 1994, without amended labeling accepted by the Agency.

Sincerely yours,



Sidney C. Jackson
Acting Product Manager (21)
Fungicide-Herbicide Branch
Registration Division (H-7505C)

enclosure

H7505C:C.Grable:cg:10/15/93

and *Thielaviopsis* in ornamental and nursery crops.

ACTIVE INGREDIENTS:

5-Ethoxy-3-trichloromethyl-1,2,4-thiadiazole*	15%
Thiophanate-methyl, (Dimethyl [(1,2-phenylene) bis (imino-carbonothioyl)] bis (carbamate))	25%
INERT INGREDIENTS	60%
TOTAL	100%

**KEEP OUT OF REACH OF CHILDREN
DANGER**
CORROSIVE. CAUSES EYE DAMAGE AND SKIN IRRITATION
HARMFUL IF SWALLOWED

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush eyes immediately with plenty of water for at least 15 minutes. **Get medical attention.**

IF SWALLOWED: Drink promptly a large quantity of milk, egg white, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. **Call a physician immediately.**

IF INHALED: Induce coughing immediately.

IF ON SKIN: Wash thoroughly with soap and water. If irritation persists, get **medical attention.**

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

**SEE BACK PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS**

*Licensed under U.S. Patent No. 3,260,588; 3,260,725

EPA REG. NO. 58185-10

EPA EST. NO. 228-IL-1

4322

**NET CONTENTS 32 OZ (2 LB)
(900 g)**

GRACE SIERRA

Grace-Sierra Crop Protection Company • 1001 Yosemite Drive, Milpitas, CA 95035
Banrot® is a registered trademark of Grace-Sierra Crop Protection Company for its brand of fungicide products

ACCEPTED
with COMMENTS
by EPA Letter Dated
OCT 22 1993
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered with EPA Reg. No.
58185-10

as a periodic drench throughout the plant's life cycle for the control of problems such as greenhouses and outdoors.

BANROT suspensions in water should be constantly applied in water uniform spaced lawn. BANROT suspensions do not need to be s. absorbed from soil. No application, if necessary in total permeation will not result in buildup, which may cause plant damage. Use at recommended rates. Overdosage may result in injury to sensitive plants. If plants are damaged by being sensitive, apply BANROT to a single number to insure safety. BANROT should not be mixed with other pesticides or nutrients.

Thoroughly rinse BANROT residues out of all equipment after use.

Woody
Acer - Aurobe - Abies -
Bambusa - Carotia - Cedar
Chamaecyparis -
Carya - Elm - Ginkgo -
Juniper - Picea - Fir -
Taxus - Thuja - Yucca -
Laurus - Ligustrum
Philadelphus -
Pithecelobium -
Poncirus -
Sapindus -
Syringa -
Tilia -
Thuja

Do not use on the ground or on plants for control of insects. Do not use on plants for control of insects. Do not use on plants for control of insects. Do not use on plants for control of insects.

COPPER SEEDLINGS
(Propagation)

NOTE: For non-bearing seedlings only. Do not apply after field transplanting and within two years of the first coffee bean harvest. Do not apply to bearing coffee plants."

Seedlings
Tree Seedlings - Max 113 grams (4 ounces) of BANROT with 190 liters (50 gallons) of water and apply 3.5 liters (one gallon) per meter² (1.2 square yards) two days prior to sowing seeds, thoroughly watering the planting medium.

Seedling Establishment - Max 85 grams (3 ounces) of BANROT with 190 liters (50 gallons) of water and apply 3.5 liters (one gallon) per meter² (1.2 square yards) if seedlings have emerged, follow application immediately by irrigating with additional water equal to at least 1/2 the volume of the fungicide drench for improved soil penetration and to rinse foliage. Re-treat at 4 to 12 week intervals if necessary.

CONTAINERIZED CITRUS SEEDLINGS AND LINERS
(Propagation)

NOTE: For non-bearing containerized citrus seedling and liner use only. Do not apply after field transplanting and within two years of the first citrus fruit harvest. Do not apply to fruit-bearing citrus trees."

Containerized Seedlings and Liners

Max 6-10 oz. of BANROT in 100 gal. water and apply to 800 sq. ft. of bench area (1 pint of solution per sq. ft.) at one and two months after seeding.

With seedlings scheduled for planting in ground later both out-of-doors, apply an additional drench, in the greenhouse, 1 to 2 weeks prior to field liner to planting.

With plants scheduled for replanting in large containers in greenhouse culture for the budding process, apply an additional drench soon transplanting. Max 6 to 10 oz. BANROT with 100 gal. of water per 400 sq. ft. Re-treat at 6 to 12 week intervals if necessary. Discontinue any applications 2 months prior to grove planting.

Follow-up all applications by immediately irrigating with additional water equal to at least 1/2 the volume of the fungicide drench for improved soil penetration and to rinse foliage.

BEST AVAILABLE COPY

STORAGE AND DISPOSAL

STORAGE: Not sensitive to light. Stable at normal temperatures and humidity. Store in a dry place in a well-closed container.

Preparation: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE PREPARATION: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, State or local procedures.

CONTAINER DISPOSAL: Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities.

WARNING

This material contains no description or the label and is intended to be the primary reference to the directions for use. Mixing and method of application, whether spraying, dusting, or other means, the disease pattern, method of the use, responsibility with other chemicals not specifically mentioned, and other information factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, change in handling of this material and in your application with directions given hereon. All agents carrying this material are authorized to provide an information only to the user.

BANROT is a registered trademark of Green-Stone Crop Protection Company, Inc. in the United States and other countries.

05 0049 000