# After application, rake-in or lightly cultivate the soil.

## TRUBAN° 5 · G FUNGICIDE

For control of damping-off root and stem rot diseases caused by Pythium and Phytophthora in ornamental and nursery crops.

ACCEPTED

Use the Pederal Insection in 1 stage to 2, died Rodenticide Act, on ar straight for the pesticide in that red under EPA Reg. No. 5

#### **ACTIVE INGREDIENTS**

### CAUTION

#### STATEMENT OF PRACTICAL TREATMENT

IF 'N EYES, Flush eyes with water for 15 minutes. IF INHALED, Induce coughing. IF ON SKIN, Wash off with soap and water. IF INHALED, Induce coughing. IF ON SKIN, Wash off with soap and water.

#### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if inhaled or absorbed through skin. Avoid contact with skin, eyes or clothing. May cause irritation. Avoid breathing dust IN CASE OF CONTACT, immediately flush eyes or skin with plenty of water. GET MEDICAL ATTENTION if irritation persists. Always wear protective clothing and respirator when mixing and applying pesticides. Wash tho bughly with soap and water after handling. Wash contaminated clothing before reuse. Do not contaminate feed or food. Keep children and pets out of treated area during application.

#### **ENVIRONMENTAL HAZARDS**

THIS PRODUCT IS TOXIC TO FISH. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

#### **DIRECTIONS FOR USE**

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

DRY SOIL MIX: Incorporate recommended amount of TRUBAN 5 • G in each cu, yd. when seeding or transplanting for control of damping-off and root rot diseases, caysed by Phythium and Phytophthora. Overdosage may result in phytotoxicity. If presented of being sensitive, apply fungicide to a small number to insure to use in greenho uses and outdoors. Where applicable, incorporate TRUBAN 5 • G after steam sterilization of soil mix.

BROADCAST AND SIDE-DRESS APPLICATIONS: Incorporate recommended amount of TRUBAN 5 • G. When side-dressing, use sharp sand, calcined clay or similar diluent to achieve good distribution. Open furrow—apply product—close—firm with packer wheel and irrigate to activate. When broadcasting, distribute evenly using cyclone or drop spreader to apply recommended amount. Harrow or till into the soil to incorporate the granules into the upper 2 to 6 in of soil.

#### STORAGE AND DISPOSAL

STORAGE: Do not store near seeds or fertilizer. Keep container closed when not in use

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal Open dumping is prohibited.

PESTICIDE DISPOSAL: Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, State, or local procedure

CONTAINER DISPOSAL: Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner in a sanitary landfill or by incineration if allowed by State and local authorities, if drum cannot be reused, dispose of in the same mariner.

#### RECOMMENDATIONS

PLANT TYPE						
BEDDING PI Ageratum Alyssum Aster Balsam Begonia Calendula Carnation Celosia	Coleus Dahlia Dianthus Foxglove Impatients Larkspur Marigold Pansy	Petunia Portulaca Salvia Snapdragon Stokesia Verbena Vinca Zinnia	CONTAINER & BEDGROWN HERBACEOUS African Violet-Begonia-Calceolaria-Carnation-Chrysanthemum-Easter Lily-Geranium-Gerbera-Gloxinia- Gypsophila-Orchids-Sinningia WOODY			
FOLIAGE PLANTS & INTERI Aglaonema Cieffenbachia Anthurium Dracaena Brassaia Echeveria Caladium Epipremnum Chinese Everç en		ORSCAPES Peperomia Pilea Philodendron Zygocactus Nephilhytis	Arctostaphylos-Azalea-Camellia- Canssa-Ceanothus-Cornus-Ficus- Fir-Hemlock-flex-Juniper-Leucothe- Magnolia-Poinsettia-Rhamnus- Southern Pine-Western Pine			

DRY SOIL MIX for BEDDING PLANTS: Mix 5 oz. TRUBAN 5 • G with each cu. yd. of soil mixture.

DRY SOIL MIX for FOLIAGE and CONTAINER PLANTS: Mix 10 oz. TRUBAN 5 • G with each cu, yd, of soil mixture, Ingredients for the soil mix should be free of large lumps and dry enough to prevent agglomeration during mixing. Re-treat as needed with Sierra's TRUBAN® 30% Wettable Powder, TRUBAN® 25% Emulsifiable Concentrate or BANROT® as indicated under appropriate soil drench recommendations.

BROADCAST APPLICATIONS: Apply 6 to 8 lb. of TRUBAN 5 • G per 1,000 sq. ft. using a cyclone or drop spreader as follows.

Make of	Desired Rate	Desired Rate	Spreader Settings	Spread
Spreader	Per acre	Per 1,000 Sq. Ft.	Twice Over	Width
Cyclone	260 Lb	6 lb	3.8	6½ Ft.
	350 Lb	8 Lb	4.2	6½ Ft
Scott	260 Lb 350 Lb.	6 Lb 8 Lb. • • •	40	20 ln 20 ln

SIDE DRESS APPLICATIONS: Open a furrow® to 10 ft. to the side of the plant to a depth of 4 to 6 in. Band 4 oz. of TRUBAN § ¿ ⑤, per 100 impers®, širjing suitable equipment and a diluent. Cover the band and packfirmly. RepeatShe operations on the other side of the plant bed lirrigate to ensure activatine of the TRUBAN 5 • G.

#### WARRANT/: ' '

This material conforms to the description on the label and ig reasonably fit for the pyriposes referred to in the directions for use. Timing and method of applicat 3.2, w.a.her, watering practices, nature of soil or poting medium, the disease problem, condition of the crop, incompatibility yillificipier chemicals not specifically recommended, and other influencing factors in the use of this privilegate beyond the control of the setter. Buyer assumes all risks of use, storage, or handling of this material not in strict accordance, with directions given herewith. NO OTHER EXPRESS OR ILLIFLIED: MARRANTY OF FITNESS OR MERCHANTABILITY IS MADE.

TRUBAN® is a registered trademark of Grace-Sierra Crop Protection Company for its brand of Fungicide products.

EPA REG. NO. 58185-13 PEST AVAILABLE COPY EPA EST. NO. 228-IL-1

6937

GRACE SIERRA

40 LB. (18 Kg)