

58185-30

11/17/2005

1/4

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval expires 05-31-98

EPA	United States	<input type="checkbox"/> Registration	OPP Identifier Number
	Environmental Protection Agency	<input type="checkbox"/> Amendment	
	Washington, DC 20460	<input checked="" type="checkbox"/> Other	

Application for Pesticide - Section I

1. Company/Product Number 58185-30	2. EPA Product Manager Mary Waller	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) FUNGO 50 WSB	PM# PM-21	
5. Name and Address of Applicant (Include ZIP Code) Scotts-Sierra Crop Protection Company 14111 Scottslawn Road Marysville, OH 43041 <input type="checkbox"/> Check if this is a new address	6. Expedited Review In accordance with FIFRA Section 3(c)(3)(b)(I), my product is similar or identical in composition and labeling to: EPA Reg. No _____ Product Name _____	

Section II

<input type="checkbox"/> Amendment - Explain below	<input type="checkbox"/> Final printed labels in response to Agency letter dated yy-yy-yy
<input type="checkbox"/> Resubmission in response to Agency letter dated XX-XX-XX	<input type="checkbox"/> "Me Too" Application
<input checked="" type="checkbox"/> Notification - Explain below	<input type="checkbox"/> Other - Explain below.

NOTIFICATION
NOV 17 2005

Explanation Use additional page(s) if necessary (For section I and Section II.)


Notification of general label revisions per PR Notice 98-10.

Section III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
*Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per Container	If "Yes" Package wgt.	No per Container
				<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
				<input type="checkbox"/> Other (Specify)	
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Other <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled					

Section IV

1. Contact Point (Complete items directly below for identification of individual to be contacted if necessary to process this application)		
Name Sheila R Kendrick	Title Senior Regulatory Specialist	Telephone No (Include Area Code) (937) 644-7214

Certification		6. Date Application Received (Stamped)
I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		
2. Signature (Scotts Sierra Crop Protection Company) BY 	3. Title Senior Regulatory Specialist	
4. Typed Name Sheila R Kendrick	5. Date November 3, 2005	

2/4

NOTIFICATION

NOV 17 2005



AllBan 50 WSB

Systemic Turf and Ornamental Fungicide

Controls a broad spectrum of diseases of containerized woody shrubs and trees, herbaceous, bedding, flowering and tropical foliage plants, shrubs, trees and flowers in the landscape, interiorscape, greenhouse, siat and screen house, nursery and diseases of turfgrasses.

ACTIVE INGREDIENTS:

Triphenylmethy	50%
OTHER INGREDIENTS	50%
TOTAL	100%

EPA Reg. No. 22564-95-8

KEEP OUT OF REACH OF CHILDREN CAUTION

See label for complete instructions. Do not use if you are pregnant or nursing. Do not use if you are allergic to any of the ingredients listed on the label. Do not use if you are allergic to any of the ingredients listed on the label. Do not use if you are allergic to any of the ingredients listed on the label.

FIRST AID

IF SWALLOWED: Call a poison control center immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF ON SKIN: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call poison control center or doctor for treatment advice.
IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes then continue rinsing eye. Call a poison control center or doctor for further treatment advice.
IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, absorbed through skin, or inhaled. Causes moderate eye irritation. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. Remove contaminated clothing and wash clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical resistance selection chart.
Applicators and other handlers must wear:
• Long-sleeved shirt and long pants,
• Chemical-resistant gloves made of any water proof material,
• Shoes plus socks.
Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS:

- Users should:
 - Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
 - Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
 - Remove PPE as soon as possible after handling the product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

THIS PESTICIDE IS TOXIC TO FISH. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply where runoff is likely to occur. Do not contaminate water by disposing of equipment washwaters.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

Not for homeowner use. For use by individuals/firms licensed or registered by the state to apply commercial or turf pest control products. Do not apply with forced wing or rotary aircraft.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and prohibitions pertaining to the requirements on the label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

Exemptive: If the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

- PPE required for each entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as:
 - Coveralls,
 - Chemical-resistant gloves made of any waterproof material,
 - Shoes plus socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Do not enter treated areas without protective clothing until spray has dried.

FOR PROFESSIONAL USE ONLY

CHEMIGATION INFORMATION

Chemigation is prohibited in the state of California.

Introduction

Pesticide labels contain directions for use which are necessary for effecting the purpose for which the product is intended and to protect health and the environment. The following information is intended to decrease environmental risks of pesticide contamination of ground water and will decrease direct human exposure to pesticide treated irrigation water by providing appropriate directions for use.

Apply the product only through pressurized drench (flood), or drip (trickle) irrigation systems. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Pressurized Drench (Flood) System

Systems utilizing a pressurized water and pesticide injection system must meet the following requirements:

- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the injection pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Sprinkler Chemigation

The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the injection pipeline to prevent water source contamination from backflow.

- The pesticide injection pipeline must contain a functional, automatic quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to

the point where pesticide distribution is adversely affected.

- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Do not apply when wind speed favors drift beyond the area intended for treatment.

Drip (Trickle) Chemigation

The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the injection pipeline to prevent water source contamination from backflow.

- The pesticide injection pipeline must contain a functional automatic quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

General Information

Pesticide supply tanks are recommended for the application of these products. See label instructions for dilution use rates and timing of application. For amenable concentrations - apply prior to use. For wettable powders - agitate continuously during application.

GENERAL INFORMATION

AllBan™ 50 WSB provides broad spectrum disease control on containerized woody, flowering, herbaceous, and tropical foliage ornamental plants, trees, ornamentals in the landscape, and turfgrasses.

The bag contains four 5 ounce AllBan™ 50 WSB water soluble bags. Add 2/3 of the required amount of water to the spray or mixing tank. With agitator running, place required amount of unopened water soluble bags into the spray tank. Do not use a spray tank or closed system's mix tank. Keep applied to areas to be treated with a boom type or other power sprayer.

Do not handle water soluble bags for longer than required to place into sprayer tank. Do not expose water soluble bags to moisture.

Apply AllBan™ 50 WSB with ground equipment, using sufficient volume of spray to provide thorough coverage. Do not tank mix with copper containing materials or with highly alkaline pesticides, such as Bordeaux mixture or lime sulfur. No claim of compatibility with other pesticides is made. Use the higher concentration or rates under conditions of severe disease pressure or when application rates are shorter than 14 days due to persistent rainfall. Also, see local State Extension Service Recommendations for application schedules.

IMPORTANT: If, after using AllBan™ 50 WSB as recommended, treatment is not effective, a tolerant strain of fungus may be present. Consult your local Scotts representative or distributor, your State Agricultural Experiment Station or your State Agricultural Extension Service for advice on the prompt use of some other labeled fungicide.

Distributed by: Scotts-Sierra Crop Protection Company • 14111 Scottslawn Road • Marysville, OH 43041
If you have any questions regarding the use of this product, call TOLL FREE 1-800-492-8255

95710

NET WEIGHT: 2 lb (907 g) (Four - 8 oz. Water-Soluble Bags)

DO NOT REMOVE PACKAGES FROM CONTAINER EXCEPT FOR IMMEDIATE USE

HORTICULTURAL APPLICATIONS

GREENHOUSE, SLAT AND SARAN HOUSES, NURSERY, LANDSCAPE AND INTERIORESCAPE.

AlBam™ 50 WSB is a broad spectrum active controlling foliar, stem and below ground diseases on a wide range of horticultural plants grown or maintained under a wide variety of conditions. Broad label applications 14 to 21 days prior to when a particular disease usually appears, and at the very latest, upon first signs of disease. Usual spray intervals range from 7 to 14 days (the latter are preventative and the former under times when conditions are judged acceptable for disease development. With hard to wet plant foliage, an acceptable wetting agent added to the spray tank according to label may increase product efficacy. Where excessive and repeated foliar wetting occurs, use of a spreader/sticker is suggested as labeled concentration. Where previously untreated applications with AlBam™ 50 WSB are considered, follow the trial suggestions under the "Wide Coverage Advantages" below below full scale use. AlBam™ 50 WSB may be used to control listed diseases on non-commercial fruit and nut trees but do not use fruit or nuts from trees treated with this product as food. Do not apply to home orchards, backyard fruit trees after fruit set.

NOTE: AlBam™ 50 WSB has been determined to be safe for use on the plant types listed in these directions based on cumulative data derived from research product trials and historical field use. Since all species and cultivars have not been tested, it is suggested that trial applications be performed if a user wishes to make an application to a plant type not listed on the label but listed on a similar use site and for a disease that is listed on the label. To all least 25 trial plants, make at least two applications at the highest concentration 7 days apart and evaluate 7 days after the last application before making full scale application. This product is not recommended for the following plants: *Nepenthes escholtzii*, *Plectranthus australis*, and *Historia quortoni*.

PLANT TYPE	such as but not limited to...
Herbaceous bedding	Ageratum, Begonia, Carna, Coleus, Dahlia, Daily Milla, Fuchsia, Fuchsia, Gazanther, Impatiens, Lonicera, Marigold, Petal, Petunia, Pinks, Primrose, Salvia, Saffron, Stachytarax, L. leaved, Verbena
Flowering	Chrysanthemum, Hydrangea, Hollyhock, Iris, Lily, Parrotia
Tropical foliage	Delonix, Dracaena, English Ivy, Philodendron, Portulaca
Woody Dicotyledons	Asplen, Hibiscus, Holly, Ilex, Juniper, Rhododendron, Rose, Pterocarya
Evergreen trees	Cycas, H. H. Palm, Palm, Spruce
Deciduous trees	Ash, London Plane, Maple, Oak, Spruce, Walnut
Flowering trees	Cherry, Crabapple, Hawthorn, Mountain Ash, Pear

Do not use fruit or nuts from treated trees as food or feed.

FOLIAR SPRAY APPLICATIONS

Application Instructions: After shaking, add the required amount of AlBam™ 50 WSB to a partially filled spray tank agitated by mechanical or hydraulic means and then add remaining amount of water. Maintain continuous agitation to keep the material in suspension and apply with properly calibrated spray equipment.

Application Concentrations (Mechanical or Hand Held): Use the recommended amount of AlBam™ 50 WSB per 100 gallons of water for the prevention and control of the disease shown below.

Disease/Pest/Insect Controlled	Concentration of AlBam™ 50 WSB fl. oz./100 gal.*
Asterisporium Colletotrichum	12-16
Black Spot of Rose Dulciorosa rose	12-16
Brown Rot and Blight Monilinia Sclerotinia Phoma	12-16
Fusarium and Verticillium Leaf Lesions: Colletotrichum, Phoma, Mycelium, Pyrenopeziza, Pyrenopeziza, Pyrenopeziza	12-16
Leaf Spots and Blights caused by: Ascochyta, Blumeriella, Botrytis, Cercospora, Colletotrichum, Corynespora, Curvularia, Didymopanax, Entomosporeum, Fusicladium, Fusarium, Ramularia, Rhizoctonia, Marssonina, Mycosphaerella, Myrothecium, Phoma, Pyrenopeziza, Scheuchzeria, Septoria, Spilocladium	12-16

G C C F L G
C C C
C C
C C C C C
C C C C C
C C C C C
C C C C C

FOLIAR DISEASES (continued)

Disease/Pest/Insect Controlled	Concentration of AlBam™ 50 WSB fl. oz./100 gal.*
Olivefruit Blight	8-16
Powdery Mildews Erysiphe, Microsphaera, Phyllosticta, Phyllosticta, Oidium, Sphaerotheca	8-16
Rust Diseases caused by: Puccinia, Gymnosporangium, Uromyces	12-16
Tip Blight Pine Sporenceps sapinae, Diplolepis	12-16
Tring Blights, Cankers, and Diebacks Dothiorella, Fusarium, Phoma	16-22

* For California label applications use the highest labeled concentration except for Olivefruit Blight and Powdery Mildews, use the lowest labeled concentration. For Tring Blights, Cankers and Diebacks use the 24 to 32 ounce concentration.

Wide Coverage Advantages: In situations where rainfall and/or overhead irrigation is the norm, use of a compatible spreader/sticker is suggested. Where wetting of foliage is difficult, use instead a compatible wetting agent. Follow the wetting agent directions stated previously under "Horticultural Applications" note.

SOIL DRENCH APPLICATIONS

Mixing Instructions: Add required amount of AlBam™ 50 WSB to partially filled tank agitated by mechanical or hydraulic means. Then add remaining required amount of water. Maintain continuous agitation to keep the material in suspension.

Application Rates and Timing for Disease Control: Use 8 to 16 ounces of AlBam™ 50 WSB per 100 gallons of water. Apply as a heavy spray or drench at the rate of 1/2 to 2 pints per square foot (100 gallons per 1,600 to 400 square feet) for shallow flats and small pots up to 4 inch deep, use 8 ounces/100 gallons applied at 1 pint per square foot. For pots and containers 4 inches or larger, refer to following table for the volume to apply. Applications may be repeated at 4 to 8 week intervals, the frequency depending upon presence of disease and optimum conditions for disease development.

Container Type	Volume to Apply/Container @ 1 1/2 fl. oz./Rate @ 2 fl. oz./Rate
4 inch	2 fl. oz.
5 inch	2-1/2 fl. oz.
6 inch	3-1/2 fl. oz.
7 inch	4-1/2 fl. oz.
8 inch	5-1/2 fl. oz.
9 inch	6-1/2 fl. oz.
10 inch	7-1/2 fl. oz.

Above 10 inch in nursery container production, a drench volume of 2-1/2 to 3 pints per square foot of surface area may be required.

Plant Types: containerized woody shrubs, trees, herbaceous bedding, flowering, and tropical foliage plants and flowers and bedding plants in the landscape.

Note: Application of this product to plug trays or seedling flats at time of seeding is not recommended.

Soil Diseases Controlled: Botrytis, Cythrodactylum, Fusarium, Gliocladium, Myrothecium, Penicillium, Ramularia, Rhizoctonia, Sclerotinia and Thielaviopsis.

Note: Pythium, Phytophthora and Cythrodactylum spathophila are not controlled by AlBam™ 50 WSB.

PREPLANT/POST HARVEST DIPS

Mixing Instructions: Mix as previously instructed for foliar spray and drench applications. Keep solution in agitation.

Application Concentration and Dipping Time: Plantlet Cuttings - Use 12 to 16 ounces of AlBam™ 50 WSB per 100 gallons of water. Immerse plants or cuttings for 10 to 15 minutes, remove and allow to drain and preferably dry. Wear clothing outlined in "PPE Requirements" for applications and other handlers.

Bulbs, Corms Tubers and Rhizomes - Use 16 to 32 ounces of AlBam™ 50 WSB per 100 gallons of water. Soak cleaned bulbs for 15 to 30 minutes in warm dry (80-85 degrees F) preferably within 48 hours of digging. Dry bulbs after treatment. If bulbs are for forcing, treat bulbs that have been heat-cured.

* For California applications use 24-32 ounces of AlBam™ 50 WSB per 100 gallons of water.

Plant Types: Propagated units of woody herbaceous, flowering and tropical foliage plants. Bulbs, corms, tubers, and rhizomes of plants such as but not limited to Calladium, Easter Lily, Tulip, Gladiolus, Daffodil, Iris.

Diseases Controlled: Botrytis, Cythrodactylum, Fusarium, Gliocladium, Myrothecium, Penicillium, Ramularia, Rhizoctonia, Sclerotinia, Thielaviopsis.

INTERIORESCAPE AND PLANTScape USE

Spray Applications: Mix well before spraying. It is important to assure complete coverage by spraying foliage to runoff, thoroughly wetting upper and lower leaf surfaces. Use concentrations recommended under "Ornamentals" for foliage, twig and branch application. When applied in public areas, use a coarse, low pressure spray. Apply during non-business hours or conduct off area during spraying and seal spray has dried. Avoid spraying wall coverings, plastic (vinyl) surfaces, drapes, carpets, and upholstery.

TURF APPLICATIONS

AlBam™ 50 WSB is a fungicide with broad spectrum activity against certain foliar and root diseases for use on all turf types such as golf course greens, lawns and fairways, athletic fields, cemeteries, parks, and commercial and home lawns. AlBam™ 50 WSB can be used both preventatively and curatively and is not phytotoxic. Not for use on turf being grown for sale or other commercial uses as seed.

Mixing Instructions: Add required amount of AlBam™ 50 WSB to partially filled tank agitated by mechanical or hydraulic means. Then add remaining required amount of water. Maintain continuous agitation to keep the material in suspension. For best results use spray machine the same day it is prepared.

Turf Types: All cool season and warm season grasses (such as but not limited to Bentgrass, Bermudagrass, Bluegrass, Fescue, Ryegrass, St. Augustinegrass, and Zoysia) and other grasses.

TURF DISEASES

Disease(s) Controlled	Rate of AlBam™ 50 WSB oz./1000 sq. ft.
Asterisporium Colletotrichum granivora	2-4
Duffia	4-6*
Sclerotinia Anthracnose	2-4
Copper Spot Gloeosporium Sclerotinia Brown Patch and Zoysia Patch Rhizoctonia solani Red Thread	2-4
Liberaria Anthracnose	2-4
Pink Snow Mold Microbotryum	2-4
Apply middle-late November before turf has stopped all growth activity. Lightly water the application into the soil zone for best results. A second application may be used in combination with FOMLI. Second spray should dry on leaf surface with no "watering in."	
(Only in areas where snow cover is not present the entire winter)	
Spry Leaf Spot Pyricularia grisea	4-6
Apply when conditions are favorable for disease development. Continue at 14 day intervals. Spray should dry on leaves with no "watering in."	
Sweeney Patch Magnaporthe Phaeo	4-6
For prevention, apply 3 applications beginning early May using 21 day intervals. Water material into root zone thoroughly after application. For suppression, apply two applications at 14 day intervals beginning when disease first appears.	
Fusarium Blight Fusicladium spp. Necrotic Ring Spot and Spring Dead Spot Leptochytrium Anther	4-6
Apply two applications at 14 day intervals beginning when disease first appears.	
Stripe Smut Ustilago sphaerogera	4-6
Make two applications at 14 day intervals when disease first appears. Water material into root zone after application. For prevention, apply in spring just before grass begins growth, and fall.	

* Refer to "Use Sites and Maximum Application Rates" to determine allowable rates for each application site.

** Note: Use 4 - 6 oz. rate for curative response to Basal Stem Anthracnose.

TURF APPLICATION DIRECTIONS

Spray uniformly over the area to be treated with a properly calibrated power sprayer.

Apply after mowing or avoid mowing for 12 hours after application. Apply sufficient water to obtain thorough coverage, usually 1-1/2 to 2-1/2 gallons per 1,000 square feet of turf area. Always treat before and approach to golf greens.

Use Sites and Maximum Application Rates

Site	Maximum Single Application Rate fl. oz./1000 sq. ft.	Maximum Seasonal Application Rate fl. oz./1000 sq. ft.
Golf Course Greens, Tees and Aprons	6	16
Golf Course Fairways (except Florida)	4	4
Golf Course Fairways (Florida Only)	2	7
Residential and Public Areas (home lawns, parks, athletic fields, schools, day care centers)	2	8

NOTE TO USER: Do not graze animals on treated turf. Do not feed clippings to livestock or poultry.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in the original container in a dry, temperature controlled area. Do not store in a manner where cross-contamination with other pesticides, herbicides, food or feed could occur. If spilled during storage or handling, contain/capture spillage and dispose of in accordance with the Pesticide Disposal Instructions listed below.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product are toxic and may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Do not re-use empty containers. Triple rinse (or equivalent), then offer for recycling or reconditioning, or puncture and dispose of as a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

This product contains a chemical known to the state of California to cause cancer, birth defects or other reproductive harm.

IMPORTANT NOTICE: DISCLAIMER AND LIMITATION OF LIABILITY

The product has been researched to provide necessary data to support its uses listed on the label. The user should always follow the label directions and exercise judgment and caution when using this product under their growing conditions. NO WARRANTY OR REPRESENTATION IS MADE, EXPRESS OR IMPLIED, CONCERNING THE RESULTS OBTAINED FROM THE USE OF THIS PRODUCT. The exclusive remedy of the user or Buyer, and the liability of The Scotts Company LLC, Scotts-Sierra Horticultural Products Company or their affiliates, for any and all losses, injuries or damages resulting from the use or handling of the product shall be the purchase price paid by the user or Buyer for the quantity of this product involved. The Buyer and all users are deemed to have accepted the terms of this Notice, which may be varied only by agreement in writing, signed by a duly authorized representative of The Scotts Company LLC or Scotts-Sierra Horticultural Products Company.



4/4

November 3, 2005

Document Processing Desk (NOTIF)
Atten: Ms. Sherada Hobgood
Registration Division, H7504C
Office of Pesticide Programs
Environmental Protection Agency
401 M Street, SW
Washington, DC 20460

Subject: Fungo 50 WSB
EPA Reg. No.: 58185-30

Dear Ms. Waller:

We are notifying the Agency of the following revisions to the subject product label:

1. Alternate brand name, AllBan 50 WSB
2. We have added the following statement to the front panel:
 - o For Professional Use Only
 - o Do Not Remove Packages from container except for immediate use.
3. We have revised the liability to note the following company information:
 - o 'The Scotts Company LLC, Scotts-Sierra Horticultural Products Company or their affiliates'
4. The Chemigation information is now contained within the 'Directions for Use section.

"This notification is consistent with the provisions of PR Notice 95-2, 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 95-2, 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA".

Enclosed are two (2) copies of the final printed label with changes highlighted to facilitate your review. Should you have any questions regarding this NOTIFICATION, please contact me at 937-644-7214.

Respectfully,
SCOTTS-SIERRA CROP PROTECTION COMPANY

BY: 
Sheila R. Kendrick
Senior Regulatory Specialist

14111 Scottslawn Road Marysville, Ohio 43041 937-644-0011

www.scotts.com