

58185-30

10-2-2006

1/4

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval expires 05-31-95

<b>EPA</b>	United States	<input type="checkbox"/> Registration	OPP Identifier Number
	<b>Environmental Protection Agency</b>	<input type="checkbox"/> Amendment	
	Washington, DC 20460	<input checked="" type="checkbox"/> Other	

**Application for Pesticide - Section I**

1. Company/Product Number 58185-30	2. EPA Product Manager Mary Waller	3. Proposed Classification  <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Fungo 50 WSB	PM# PM-21	
5. Name and Address of Applicant (Include ZIP Code) Scotts-Sierra Crop Protection Company 14111 Scottslawn Road Marysville, OH 43041  <input type="checkbox"/> Check if this is a new address	6. <b>Expedited Review.</b> In accordance with FIFRA Section 3(c)(3)(b)(I), my product is similar or identical in composition and labeling to:  EPA Reg. No. _____ Product Name _____	

**Section II**

<input type="checkbox"/> Amendment - Explain below. <input type="checkbox"/> Resubmission in response to Agency letter dated XX-XX-XX <input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____ <input type="checkbox"/> "Me Too" Application <input type="checkbox"/> Other - Explain below.
---	--

**NOTIFICATION**  
OCT 2 2006

Explanation: Use additional page(s) if necessary. (For section I and Section II.)  
  
Notification- Label updates per PR Notice 98-10 (see attached letter)


**Section III**

1. Material This Product Will Be Packaged In:			
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No  If "Yes"    No. per Unit Packaging wgt.    Container	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No  If "Yes"    No. per Package wgt.    Container	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify)
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container	
5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product		6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Other _____ <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled	

**Section IV**

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)

Name Lisa R. McClellan	Title Senior Regulatory Scientist	Telephone No. (Include Area Code) (937) 644-7691
---------------------------	--------------------------------------	---

<b>Certification</b>		<b>6. Date Application Received (Stamped)</b>
I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		
2. Signature/ Scotts Sierra Crop Protection Company BY: 	3. Title Sr. Product Regulatory Scientist	
4. Typed Name Lisa R. McClellan	5. Date September 13, 2006	

2/4

NOTIFICATION

OCT 2 2006



# Fungo 50 WSB

## Systemic Turf and Ornamental Fungicide

Controls a broad spectrum of diseases of containerized woody shrubs and trees, herbaceous, bedding, flowering and tropical foliage plants, shrubs, trees and flowers in the landscape, interiorscape, greenhouse, stat and saran house, nursery and diseases of turfgrasses.

<b>ACTIVE INGREDIENT:</b>	
Thiophanate-Methyl <sup>1</sup>	50%
<b>OTHER INGREDIENTS:</b>	50%
<b>TOTAL</b>	<b>100%</b>

1 CAS # 23564-05-8

### KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle (if you do not understand the label, find someone to explain it to you in detail).

EPA REG. NO. 58145-30 EPA EST. NO. 67845-AZ-001  
Product of U.S.A.

©2006, Scotts-Sierra Horticultural Products Company. World rights reserved. Fungo<sup>®</sup> is a trademark of OMC Investments, Inc. for its brand of Fungicide products.

#### FIRST AID

**IF SWALLOWED:** Call a poison control center immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**IF ON SKIN:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present after first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for further treatment advice.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed, absorbed through skin, or inhaled. Causes moderate eye irritation. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. Remove contaminated clothing and wash clothing before reuse.

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance selection chart.

- Applicators and other handlers must wear:
  - Long-sleeved shirt and long pants
  - Chemical-resistant gloves made of any water proof material.
  - Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### USER SAFETY RECOMMENDATIONS:

- Users should:
- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
  - Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
  - Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible wash thoroughly and change into clean clothing.

#### ENVIRONMENTAL HAZARDS

**THIS PESTICIDE IS TOXIC TO FISH.** Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply where runoff is likely to occur. Do not contaminate water by disposing of equipment washwaters.

#### DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

Not for homeowner use. For use by individuals/firms licensed or registered by the state to apply ornamental or turf pest control products. Do not apply with fixed wing or military aircraft.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. The standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

Exemption: If the product is applied by drenching the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls,
- Chemical-resistant gloves made of any waterproof material,
- Shoes plus socks

#### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides, 40 CFR Part 170. The WPS applies when this product is used to protect agricultural plants on farms, forests, nurseries or greenhouses.

- Do not enter treated areas without protective clothing until spray has dried.

#### FOR PROFESSIONAL USE ONLY

#### CHEMIGATION INFORMATION

Chemigation is prohibited in the State of California.

#### Introduction

Pesticide labels contain Directions For Use that are necessary for effecting the purpose for which the product is intended. Further, they protect health and environmental risks of pesticide contamination of ground water and decrease direct human exposure to pesticide-treated irrigation water.

Do not connect chemigation system to a public water system.

Apply this product only through pressurized (flood), or drip (trickle) irrigation systems. Do not apply this product through any other type of irrigation systems.

Drop injury, lack of effectiveness or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect any irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person shall shut the system down and make necessary adjustments should the need arise.

#### Pressurized Drench (Flood) System

Systems utilizing a pressurized water and pesticide injection system must meet the following requirements:

- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- Systems using a gravity flow pesticide dispensing system must meter the pesticide into the water at the head of the field and downstream of a hydraulic discontinuity such as a drop structure or weir box to decrease potential for water source contamination from backflow if water flow stops.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

#### Sprinkler Chemigation

The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

- The pesticide injection pipeline must contain a functional, automatic quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

• The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

• Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

• Do not apply when wind speed favors drift beyond the area intended for treatment.

#### Drip (Trickle) Chemigation

The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

- The pesticide injection pipeline must contain a functional, automatic quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

#### General Information

Pesticide supply tanks are recommended for the application of these products. See label instructions for dilution use rates and timing of application. For emulsifiable concentrates - agitate prior to use. For wettable powders - agitate continuously during application.

#### GENERAL INFORMATION

Fungo 50 WSB provides broad spectrum disease control on containerized woody, flowering, herbaceous, and tropical foliage ornamental plants, trees and shrubs in the landscape and nurseries.

This bag contains four 5 ounce Fungo 50 WSB water soluble bags. Add 2/3 of the required amount of water to the spray or mixing tank. With agitator running, place required amount of unopened water soluble bags into the partially filled spray-tank or closed-system mix tank. Keep agitated to insure even suspension of material. Make fresh daily. Spray uniformly over the area to be treated with a boom type or other power sprayer.

Do not handle water soluble bags for longer than required to place into spray tank. Do not expose water soluble bags to moisture.

Apply Fungo 50 WSB with ground equipment, using sufficient volume of spray to provide thorough coverage. Do not tank mix with copper containing materials or with highly alkaline pesticides, such as Bordeaux mixture or lime sulfur. No claim of compatibility with other pesticides is implied. Use the higher concentration or rates under conditions of severe disease pressure or when application intervals are shorter than 14 days due to persistent rainfall. Also, see local State Extension Service Recommendations for application schedules.

**IMPORTANT:** If, after using Fungo 50 WSB as recommended, treatment is not effective, a tolerant strain of fungus may be present. Consult your local Scotts representative or distributor, your State Agricultural Experiment Station or your State Agricultural Extension Service for advice on the prompt use of some other labeled fungicide.

Distributed by: Scotts-Sierra Crop Protection Company • 14111 Scottslawn Road • Marysville, OH 43041

If you have any questions regarding the use of this product, call TOLL FREE 1-800-492-8255

# 95710

## NET WEIGHT: 2 lb (907 g) (Four - 8 oz. Water-Soluble Bags)

DO NOT REMOVE PACKAGES  
FROM CONTAINER EXCEPT FOR  
IMMEDIATE USE

**HORTICULTURAL APPLICATIONS**

GREENHOUSE, SLAT AND SARAN HOUSES, NURSERY, LANDSCAPE AND INTERIORESCAPE.  
Fungo50 WSB is broad spectrum in activity controlling foliar, stem and below ground diseases on a wide range of horticultural plants grown or maintained under a wide variety of conditions. Begin foliar applications 14 to 21 days prior to when a particular disease usually appears, and if the very latest, upon first signs of disease. Usual spray intervals range from 7 to 14 days, the latter as preventative and the former under times when conditions are judged applicable for disease development. With hard to wet plant foliage, an acceptable wetting agent added to the spray tank according to label may increase product efficacy. Where excessive and repeated rotting occurs, use of a spreader-sticker is suggested at labeled concentration. Where previously untested adjuvants with Fungo 50 WSB are considered, follow the trial suggestions under the "Note Concerning Adjuvants" below before full scale use. Fungo 50 WSB may be used to control listed diseases on non-commercial fruit and nut trees but do not use fruit or nuts from trees treated with this product as food. Do not apply to home orchard/backyard fruit trees after fruit set!

NOTE: Fungo 50 WSB has been determined to be safe for use on the plant types listed in these directions based on cumulative data derived from research product trials and historical field use. Since all species and cultivars have not been tested, it is suggested that trial applications be performed if a user wishes to make an application to a plant type not listed on the label but found on a similar site and for a disease that is listed on the label. To all least 25 trial plants, make at least two applications at the highest concentration 7 days apart and evaluate 7 days after the last application before instituting full scale application. This product is not recommended for the following plants: *Nepenthes spathulata*, *Platanus australis*, and *Hibiscus quercin*.

PLANT TYPE	which are but not limited to:
Herbaceous Bedding	<i>Agrostis</i> , <i>Diopsis</i> , <i>Canna</i> , <i>Delphinium</i> , <i>Dusty Miller</i> , <i>Fuchsia</i> , <i>Geranium</i> , <i>Impatiens</i> , <i>Lavender</i> , <i>Margold</i> , <i>Penstemon</i> , <i>Peonies</i> , <i>Primulas</i> , <i>Salvia</i> , <i>Statice</i> , <i>Strawflowers</i> , <i>Tickseed</i> , <i>Verbena</i>
Flowering	<i>Chrysanthemum</i> , <i>Hydrangea</i> , <i>Hollyhock</i> , <i>Iris</i> , <i>Lily</i> , <i>Floribunda</i>
Tropical Foliage	<i>Dieffenbachia</i> , <i>Dieffenbachia</i> , <i>English Ivy</i> , <i>Philodendron</i> , <i>Pothos</i>
Woody Ornamentals	<i>Azalea</i> , <i>Hibiscus</i> , <i>Holly</i> , <i>Liquidambar</i> , <i>Rhododendron</i> , <i>Rose</i> , <i>Pyracantha</i>
Evergreen Trees	<i>Douglas Fir</i> , <i>Pinus</i> , <i>Larch</i> , <i>Pine</i> , <i>Service</i>
Deciduous Trees	<i>London Plane</i> , <i>Maple</i> , <i>Oak</i> , <i>Sycamore</i> , <i>Willow</i>
Flowering Trees	<i>Cherry</i> , <i>Crapeholme</i> , <i>Hawthorn</i> , <i>Mountain Ash</i> , <i>Pear</i>

Do not use fruit or nuts from treated trees as food or feed.

**FOLIAR SPRAY APPLICATIONS**

Hydraulic Application: Mixing Instructions  
Add the required amount of Fungo 50 WSB to a partially filled spray tank agitated by mechanical or hydraulic means and then add remaining amount of water. Maintain continuous agitation to keep the material in suspension and apply with properly calibrated spray equipment.

Application Concentration (Mechanical or Hand Held)  
Use the recommended amount of Fungo 50 WSB per 100 gallons of water for the prevention and control of the disease shown below.

DISEASES	Concentration of Fungo 50 WSB oz./100 gal.*
<b>DISEASES/PATHOGENS CONTROLLED</b>	
<b>Anthracnose</b>	
<i>Colletotrichum</i>	
Apply as buds break or at first sign of disease. Repeat at 7-14 day intervals as needed during disease period.	12-16
<b>Black Spot of Rose</b>	
<i>Diplocarpon roseae</i>	
Apply early summer or at first sign of disease. Repeat every 7-14 days as needed during disease period.	12-16
<b>Brown Rot and Blight</b>	
<i>Botrytis</i> , <i>Sclerotinia</i> , <i>Witchblight</i>	
Apply late Spring or at first sign of disease. Repeat every 7-14 days as needed during disease period.	12-16
<b>Fusicladium and Yenteria Leaf Scabs on</b>	
<i>Crapeholme</i> , <i>Hawthorn</i> , <i>Pear</i> , <i>Mountain Ash</i> , <i>Pyracantha</i> , etc.	
Apply as buds break. Repeat every 7-14 days during disease period. Effective control requires coverage during expansion. Rotations with chlorothalonil or propiconazole can be utilized.	12-16
<b>Leaf Spots and Blights caused by</b>	
<i>Ascochyta</i> , <i>Bufovia</i> , <i>Botrytis</i> , <i>Cercospora</i> , <i>Dothidea</i> , <i>Corynespora</i> , <i>Chytrium</i> , <i>Dactyloctenium</i> , <i>Entomospium</i> , <i>Fabraea</i> , <i>Fusarium</i> , <i>Ramularia</i> , <i>Rhizoctonia</i> , <i>Marcossonia</i> , <i>Mycosphaerella</i> , <i>Myrothecium</i> , <i>Phoma</i> , <i>Phyllosticta</i> , <i>Schroeterium</i> , <i>Septoria</i> , <i>Sphaeria</i>	
Apply when disease symptoms first appear. Repeat every 7-14 days during disease period. Rotations with chlorothalonil can be utilized.	12-16

**FOLIAR DISEASES (continued)**

Disease/Pathogen Controlled	Concentration of Fungo 50 WSB oz./100 gal.*
<b>Oxleya Blight</b>	
Apply as flowers open. Repeat every 7-14 days during disease period.	8-16
<b>Powdery Mildews</b>	
<i>Erysiphe</i> , <i>Microsporangium</i> , <i>Phyllosticta</i> , <i>Podosphaera</i> , <i>Oidium</i> , <i>Sphaerotheca</i>	
Apply when disease first appears and repeat every 7-14 days. Rotations with other effective products is recommended.	8-16
<b>Rust Diseases caused by</b>	
<i>Puccinia</i> , <i>Gymnosporangium</i> , <i>Uromyces</i>	
Apply late Spring or when symptoms first appear. Repeat every 7-14 days during disease period. Rotations with other effective products is recommended.	12-16
<b>Tip Blight of Pine</b>	
<i>Sphaeropsis sapinea</i> , <i>Diplodia pinea</i>	
Begin application in Spring when new growth starts. Make a second application just before needles emerge from the sheath and a third application 7 days later. Thorough coverage is essential for optimal disease control.	12-16
<b>Twig Blights, Cankers, and Diebacks</b>	
<i>Diaporthe</i> , <i>Botrytis</i> , <i>Phoma</i> , <i>Phomopsis</i>	
Apply when symptoms first appear. Repeat every 7-14 days during disease period.	16-32

\* For California foliar applications use the highest labeled concentration except for Oxleya Blight and Powdery Mildews, use the lowest labeled concentration. For Twig Blights, Cankers and Diebacks use the 24 to 32 ounce concentration.

Note Concerning Adjuvants: In situations where rainfall and/or overhead irrigation is the norm, use of a corrosion spreader-sticker is suggested. Where wetting of foliage is difficult, use instead a compatible wetting agent. Follow the phytoxicity precautions stated previously under "Horticultural Applications" note.

**SOIL DRENCH APPLICATIONS**

Mixing Instructions: Add required amount of Fungo 50 WSB to partially filled tank agitated by mechanical or hydraulic means. Then add remaining required amount of water. Maintain continuous agitation to keep the material in suspension.

Application Rates and Timing for Disease Control: Use 8 to 16 ounces of Fungo 50 WSB per 100 gallons of water. Apply as a heavy spray or drench at the rate of 1/2 to 2 pints per square foot (100 gallons for 1,000 to 400 square feet). For shallow flats and small pots up to 4 inch size, use 8 ounces/100 gallons applied at 1 pint per square foot. For pots and containers 4 inches or larger, refer to following table for the volume to apply. Applications may be repeated at 4 to 8 week intervals, the frequency depending upon presence of disease and optimum conditions for disease development.

Container Type	Volume to Apply/Container @ 1.0 LPL/L Rate	@ 2.0 LPL/L Rate
4 inch	2.0 fl. oz.	
5 inch	2-1/2 fl. oz.	
6 inch	3-1/2 fl. oz.	
7 inch	4-1/2 fl. oz.	
8 inch	5-1/2 fl. oz.	
9 inch	6-1/2 fl. oz.	
10 inch	7-1/2 fl. oz.	

Above 10 inch in nursery container production, a drench volume of 2-1/2 to 3 pints per square foot of surface area may be required.

Plant Types: containerized woody shrubs, trees, herbaceous bedding, flowering and tropical foliage plants and flowers and bedding plants in the landscape.

Note: Application of the product to plug trays or seedling flats at time of seeding is not recommended.

Soil Diseases Controlled: *Botrytis*, *Cylindrocladum*, *Fusarium*, *Gloeosporium*, *Myrothecium*, *Penicillium*, *Ramularia*, *Rhizoctonia*, *Sclerotinia* and *Thelephora*.

Note: Phythium, *Phytophthora* and *Cylindrocladum apothecium* are not controlled by Fungo 50 WSB.

**PREPLANT/POST HARVEST DIPS**

Mixing Instructions: Mix as previously instructed for foliar spray and drench applications. Keep solution in agitation.

Application Concentration and Dipping Time: Plants or Cuttings - Use 12 to 16 ounces of Fungo 50 WSB per 100 gallons of water. Immerse plants or cuttings for 10 to 15 minutes, remove and allow to drain and preferably dry. Wear clothing outlined in "PPE Requirements" for applications and other handlers.

Bulbs, Corms, Tubers and Rhizomes - Use 15 to 32 ounces of Fungo 50 WSB per 100 gallons of water. Soak cleaned bulbs for 15 to 30 minutes in warm dip (80-85 degrees F) preferably within 48 hours of digging. Dry bulbs after treatment. It bulbs are for forcing, treat bulbs that have been heat-treated.

\* For California applications use 24-32 ounces of Fungo 50 WSB per 100 gallons of water.

Plant Types: Propagated units of woody herbaceous, flowering and tropical foliage plants. Bulbs, corms, tubers, and rhizomes of plants such as but not limited to *Calladium*, *Easter Lily*, *Hyacinth*, *Gladiolus*, *Daffodil*, *Iris*.

Diseases Controlled: *Botrytis*, *Cylindrocladum*, *Fusarium*, *Gloeosporium*, *Myrothecium*, *Penicillium*, *Ramularia*, *Rhizoctonia*, *Sclerotinia*, *Thelephora*.

**INTERIORESCAPE AND PLANTSCAPE USE**

Spray Application: Mix well before spraying. It is important to ensure complete coverage by spraying foliage to runoff, thoroughly wetting upper and lower leaf surfaces. Use concentrations recommended under "Ornamentals" for foliar, twig and drench application. When applied in public areas, use a coarse, low pressure spray. Apply during non-business hours or cordon off area during spraying and until spray has dried. Avoid spraying wall coverings, plastic (vinyl) surfaces, drapes, carpets, and upholstery.

**TURF APPLICATIONS**

Fungo 50 WSB is a fungicide with broad spectrum activity against certain foliar and soil diseases for use on all turf types such as golf course greens, tees and fairways, athletic fields, cemeteries, parks, and commercial and home lawns. Fungo 50 WSB can be used both preventatively and curatively and is not phytotoxic. Not for use on turf being grown for sale or other commercial uses as seed.

Mixing Instructions: Add required amount of Fungo 50 WSB to partially filled tank agitated by mechanical or hydraulic means. Then add remaining required amount of water. Maintain continuous agitation to keep the material in suspension. For best results use spray mixture the same day it is prepared.

Turf Types: All cool season and warm season grasses (such as but not limited to *Bentgrass*, *Bermudagrass*, *Boutelougrasses*, *Festucas*, *Ryegrasses*, *St. Augustinegrasses*, and *Zoysiagrasses*) or their mixtures.

**TURF DISEASES**

Disease(s) Controlled	Rate of Fungo 50 WSB oz./1000 sq. ft.*
<b>Anthracnose</b>	2-4
<i>Colletotrichum graminicola</i>	(4-8)**
<b>Dothier Spot</b>	2-4
<i>Sclerotinia homoeocarpa</i>	
Apply when disease first appears. Continue at 14 day intervals as needed. Spray should dry on leaves with no "watering in."	
<b>Copper Spot</b>	2-4
<i>Gloeosporium sorghii</i>	
<b>Brown Patch and Zoysia Patch</b>	2-4
<i>Rhizoctonia solani</i>	
<b>Rust Thread</b>	2-4
<i>Laetisaria aculeiformis</i>	
Apply when disease first appears. Continue at 14 day intervals as needed. Spray should dry on leaves with no "watering in."	
<b>Peak Sewer Mold</b>	2-4
<i>Microdochium nivale</i>	
(Only in areas where snow cover is not present the entire winter)	
Apply middle-late November before turf has stopped all growth activity. Lightly water this application into the root zone for best results. A second application may be used in combination with PCNB. Second spray should dry on leaf surfaces with no "watering in."	
<b>Gray Leaf Spot</b>	4-6
<i>Pyricularia grisea</i>	
Apply when conditions are favorable for disease development. Continue at 14 day intervals. Spray should dry on leaves with no "watering in."	
<b>Sweeney Patch</b>	4-6
<i>Aspergillus Rose</i>	
For prevention, apply 3 applications beginning early May using 21 day intervals. Water material into root zone thoroughly after application. For suppression, apply two applications at 14 day intervals beginning when disease first appears.	
<b>Fusarium Blight</b>	4-6
<i>Fusarium spp.</i>	
<b>Necrotic Ring Spot and Spring Dead Spot</b>	
<i>Leptophthalma acuta</i>	
Apply two applications at 14 day intervals beginning when disease first appears.	
<b>Striped Stem</b>	4-6
<i>Helminthosporium</i>	
Make two applications at 14 day intervals when disease first appears. Water material into root zone after application. For prevention, apply in spring (just before grass begins growth), and fall.	

\* Refer to "Use Sites and Maximum Application Rates" to determine allowable rates for each application etc.

\*\* Note: Use 4 - 6 oz rate for curative response to Basal Stem Anthracnose.

**TURF APPLICATION DIRECTIONS**

Spray uniformly over the area to be treated with a properly calibrated power sprayer.

Apply after mowing or avoid mowing for 12 hours after application. Apply sufficient water to obtain thorough coverage, usually 1-1/2 to 2-1/2 gallons per 1,000 square feet of turf area. Always treat aprons and approaches to golf greens.

Use Sites and Maximum Application Rates

Site	Maximum Single Application Rate oz./1000 sq. ft.	Maximum Seasonal Application Rate oz./1000 sq. ft.
Golf Course Greens, Tees and Aprons	6	16
Golf Course Fairways (Except Florida)	4	4
Golf Course Fairways (Florida Only)	2	2
Residential and Public Areas (home lawns, parks, athletic fields, schools, day care centers)	2	8

NOTE TO USER: Do not graze animals on treated turf. Do not feed clippings to livestock or poultry.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in the original container in a dry, temperature controlled area. Do not store in a manner where cross-contamination with other pesticides, fertilizers, food or feed could occur if spilled during storage or handling, container/closure spillage and disposal in accordance with the Pesticide Disposal Instructions listed below.


PESTICIDE DISPOSAL: Wastes resulting from the use of this product are toxic and may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Do not re-use empty container. Dispose of empty container in a sanitary landfill or by incineration or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

This product contains a chemical known to the state of California to cause cancer, birth defects or other reproductive harm.

**IMPORTANT NOTICE: DISCLAIMER AND LIMITATION OF LIABILITY**

This product has been researched to provide necessary data to support its use listed on the label. The user should always follow the label directions and exercise judgment and caution when using this product under their growing conditions. NO WARRANTY OR REPRESENTATION IS MADE, EXPRESS OR IMPLIED, CONCERNING THE RESULTS OBTAINED FROM THE USE OF THIS PRODUCT. The exclusive remedy of the user or Buyer, and the liability of The Scotts Company LLC, Scotts-Sierra Horticultural Products Company or their affiliates, for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or Buyer for the quantity of this product involved. The Buyer and all users are deemed to have accepted the terms of this notice, which may be varied only by agreement in writing, signed by a duly authorized representative of The Scotts Company LLC or Scotts-Sierra Horticultural Products Company.



**The Scotts Company**

and Subsidiaries

4/4  
**Scotts**

September 13, 2006

Ms. Sherada Hobgood  
Document Processing Desk (NOTIF)  
Office of Pesticide Programs (7504P)  
US Environmental Protection Agency  
Room S-4900, One Potomac Yard  
2777 South Crystal Drive  
Arlington, VA 22202-4501

Subject: Fungo 50 WSB  
EPA Reg. No.: 58185-30

Dear Ms. Hobgood:

We are notifying of the Agency of the following addition to the subject label:

- We have added a Prop 65 statement per California requirements, "This product contains a chemicals know to the state of California to cause cancer, birth defeats or other reproductive harm."
- We also requesting that "AllBan 50 WSB" be added as an additional brand name for the subject product.

**"This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA"**

Should you have any questions regarding this submission, please contact me at (937) 644-7691.

Respectfully,

SCOTTS-SIERRA CROP  
PROTECTION COMPANY

BY: 

Lisa R. McClellan  
Sr. Regulatory Scientist