to thempore to the communicate water by dicaming of equipment The Discourse of the More than the part of the transfer to water when Committee to the committee the transfer of the committee of the committee

ATT \$31 18 4 1 18 N 中央 1 元 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	:ek:de
SIGNATURE OF APPROVING OFFICIAL	DATE O/ A/A
DA C 0570 / (D 5.3/)	

- e. Change DIRECTIONS FOR FEEDING USE to DIRECTIONS FOR USE.
- f. Use the full product name or "this product" within the Directions for Use.
- g. Insert single and double asterisks at appropriate points on front panel to agree with footnotes.
- 3. Please provide the chemical identity and Material Safety Data Sheet for the following:
 - a. Animal protein products;
 - b. Plant protein products;
 - c. Grain products; and
 - d. Processed grain byproducts.
 - 4. Please provide the following Data Matrix requirements:
 - a. Discussion of Formation of [Unintentional] Ingredients—a brief description of impurities formed during the manufacturing/formulation process, in packaging, or during storage. If you do not expect any impurities during these stages, please so state.
 - b. Analytical Method--provide the methods used to analyze for the active ingredients. Provide either the MRID or Accession Number, not the Pesticide Petition Number.
- 5. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

George T. LaRocca Product Manager (15) Insecticide-Rodenticide Branch Registration Division (TS-767C)

Enclosures

NET WEIGHT 50-1bs.

WENDLAND'S FLY CONTROL RANGE MEAL

(contains Rabon Oral Larvicide)

Cattle (beef & dairy): Control of fecal flies in manure of treated cattle. Prevents development of face flies, horn flies, house flies and stable flies in the manure of treated cattle.

ACTIVE INGREDIENT

2-Chloro-1-(2,4,5-trichlorophenyl)	by Weight	ACCEPTED with COMMINTS
vinyl dimethyl phosphate	0.087%	in EPA Letter Dated:
Inert Ingredients	99.913%	
TOTAL	100 00%	SEP 12 1983

*Contains 0.395 grams of Rabon per pound.
**Refers only to ingredients which are not larvicidal.

Under the Federal Insecticide Fungicide, and Rodenticide Act a amended. for the pesticide registered under EPA Reg. No

GUARANTEED ANALYSIS

58349-7

Crude Protein, minimum	15.00%
(This includes not more than 5.25% equivalent protein	n from non-protein
nitrogen.)	0.500
Crude Fat, minimum	2.50%
Crude Fiber, maximum	7.50%
Calcium (Ca), minimum	2.50%
Calcium (Ca), maximum	3.50%
Phosphorus (P), minimum	1.00%
Salt (NaCl), minimum	23.00%
Salt (NaCl), maximum	27.50%

INGREDIENTS

Grain products, plant protein products, processed grain by-products, forage products, animal protein products, roughage products (5%), molasses product, calcium carbonate, monocalcium phosphate, salt, calcium sulfate, potassium sulfate, magnesium sulfate, sodium bicarbonate, magnesium oxide, vitamin A supplement, d-activated animal sterol (source of vitamin P3), vitamin E supplement, manganous oxide, zinc oxide, ferrous sulfate, copper oxide, ethylenediamine dihydriodide, cobalt sulfate and sodium selenite....

CAUTION USE AS DIRECTED.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Precautionary Statements Hazards to Humans

CAUTION: Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water. If irritation persists, see a physician.

WENDLAND'S FLY CONTROL RANGE MEAL.
Page 2

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE and DISPOSAL
Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a dry place in original container.

CONTAINER DISPOSAL: Completely empty bag of contents. Then dispose of empty packaging in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Pesticide Disposal: Wastes resulting from the use of this may be disposed of on-site or at an approved waste disposal facility.

DIRECTIONS FOR FEEDING USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Rations containing this product may be fed up to slaughter and to lactating dairy cows without withholding the milk from market during or after treatment.

Start feeding Rabon larvicidal feeds early in the spring before flies begin to appear and continue feeding throughout the summer and into the fall until cold weather restricts fly activity.

When fed, this product passes through the digestive system into the animal's manure where it kills fly larvae on contact shortly after fly eggs hatch.

It prevents the development of fly larvae in the manure of treated animals, but is not effective against existing adult flies.

In some cases, supplemental fly control measures may be needed in and around cattle lots and barns to control adult house flies and stable thies which can breed not only in manure but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, this product should he used in conjunction with other good management and sanitation practices.

Wendland's Fly Control Range Meal should be supplied free choice for cattle, but no less than 0.18 lbs. per/cwt/day should be consumed. An 1100 pound brood cow should consume a minimum of 2-lbs. per head per day. Adjust intake by movement of feed sources closer or farther from sources of water.

Manufactured by WENDLAND'S FARM PRODUCTS, INC. TEMPLE, TEXAS 76503

EPA Reg. No. 58349-

EPA Est. No. 58349-TX-001