


PM13 58866-10

10/14

 <p>U.S. ENVIRONMENTAL PROTECTION AGENCY          UNITED STATES OFFICE of Pesticide Programs          Registration Division (H7505C)          401 M St., S.W.          Washington, D.C. 20460</p> <p><b>NOTICE OF PESTICIDE:</b>  <input checked="" type="checkbox"/> Registration  <input type="checkbox"/> Reregistration</p> <p>(under FIFRA, as amended)</p>	EPA Reg. Number: 58866-10	Date of Issuance: APR 5 1991
	Term of Issuance: Conditional	
	Name of Pesticide Product: Diazinon 25 Insect Control	
Name and Address of Registrant (include ZIP Code): ProGuard, Inc. P.O. Box 550 Suisun, CA 94585-0550		
<p>Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.</p>		
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.</p> <p>Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:</p> <ol style="list-style-type: none"> <li>Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.</li> <li>Make the labeling changes listed below before you release the product for shipment:           <ol style="list-style-type: none"> <li>Add the phase "EPA Registration No. 58866-10"</li> <li>Under the active ingredient statement change "For outdoor use on the home lawn and home garden only." to read "For outdoor use only on the home lawn and home garden."</li> </ol> </li> </ol>		
Signature of Approving Official:		Date:

EPA Form 8570-6

BEST AVAILABLE COPY

CONCURRENCES						
SYMBOL	H-7505C					
SURNAME	RND					
DATE	7/23/91					

OFFICIAL FILE

EPA Form 1320-1A (1/90)

Printed on Recycled Paper

20/14

c. Under Measurement Conversions change "1 1/2 tablespoons in 3 gals. of water = 1 1/2 teaspoons in 1 gal. of water 2 tablespoons in 3 gals. of water= 2 teaspoons in 1 gal. of water" to read "2 tablespoons in 3 gals. of water= 2 teaspoons in 1 gal. of water or 6 ozs. in 18 gals. of water".

d. Under Lawn Pest Control change ".....water lawn with at least inch of water immediately after spraying this product....." to read ".....water lawn with at least 1/4 inch of water immediately after spraying this product....." .

e. Under Fruits and Nuts: PHI add for "Raspberries 7, for Strawberries 5 and for Walnuts 45".

3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

) If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Enclosures

)

CONTAINER 30/14

# DIAZINON-25

## INSECT CONTROL

This product must not be used on golf courses and sod farms.

**ACTIVE INGREDIENTS:**

Diazinon: O, O-Diethyl O(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate ..... 25.0%

INERT INGREDIENTS\*: ..... 75.0%

TOTAL: ..... 100.0%

\*Contains aromatic petroleum distillates.

EPA Reg. No. 58866  
EPA Est. 58866-CA-1

For outdoor use on the home lawn and the home garden only.  
Spray concentrate. Contains diazinon insecticide.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

See side panel for additional precautionary statements.

**PROTECTS:** Lawns, Vegetables, Fruits and Nuts, Roses, Flowers, Trees, Shrubs.  
**CONTROLS:** Ants, Aphids, Chinch Bugs, Cutworms, Fire Ants<sup>1</sup>, Flea Beetles, Grubs, Mites, Sod Webworms, and many others.

<sup>1</sup> Aids in control.

**NET CONTENTS: 8 oz.**

ACCEPTED  
with COMMENTS  
in EPA Letter Dated

APR 5 1991  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.  
58866-CA

**ProGuard, Inc.  
Suisun, CA 94585**

**BEST AVAILABLE COPY**

**Ornamental Trees & Shrubs**

Arborvitae. Aphids. Mites. Scale Crawlers. Whiteflies. Tent Caterpillars and others \*  
 Azaleas. Birch.  
 Boxwood. Camellias.  
 Douglas Fir. E.m.  
 Hawthorn Holly.  
 Juniper. Liriod.  
 Locust. Maple.  
 Mimosa. Oak. Pine.  
 Ornamental Plum.  
 Poplar.  
 Rhododendron.  
 Spruce. Willow

**Storage and Disposal****Storage**

Do not store near heat or open flame. Store in original container in an area inaccessible to children or pets. **USE CONTENTS WITHIN 6 MONTHS AFTER OPENING.**

**Disposal**

Do not reuse empty container. Thoroughly rinse container, securely wrap in several layers of newspaper, and discard in trash. If disposal of a partially full container is necessary, make sure cap is tightly closed, securely wrap the container in several layers of newspaper and discard in trash.

**Precautionary Statements****Hazards to Humans and Domestic Animals****CAUTION**

Harmful if swallowed, inhaled, or absorbed through skin. Causes moderate eye injury. Avoid breathing spray mist. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Do not use inside the home. Avoid contamination of feed and foodstuffs. Do not use on nousehold pets or humans. Do not allow children or pets to go onto a treated area until the treated area, such as grass or soil, has dried. Do not contaminate ornamental fish ponds. Food utensils such as spoons and measuring cups must not be used for food purposes after use in measuring pesticides.

**Statement of Practical Treatment**

**If swallowed:** Call a physician immediately. **DO NOT INDUCE VOMITING UNLESS NO OTHER TREATMENT IS AVAILABLE.** Vomiting may cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing.

**If inhaled:** Remove victim to fresh air and apply artificial respiration if indicated. Get medical attention.

**If on skin:** Wash promptly with soap and water. Rinse thoroughly. Get medical attention.

**If in eyes:** Rinse eyes with plenty of water and call a physician immediately.

**Note to Physician:** This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine. This product contains aromatic petroleum

distillates which may present an aspiration hazard. Gastric lavage may be indicated if product was taken internally.

**Environmental Hazards**

This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of sprays. For lawns, where irrigation (watering) is recommended in the directions, or if waterfowl, i.e., ducks or geese, can be expected in the treated area after treatment (except spot treatment), apply at least 1/4 inch of water immediately after spraying this product, however, stop watering before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

This pesticide is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this pesticide or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

**Physical or Chemical Hazards**

Do not use, pour, spill, or store near heat or open flame.

**BEST AVAILABLE COPY**

50714

## DIAZINON-25

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

### GENERAL INFORMATION

) Diazinon-25 is an emulsifiable concentrate product which must be diluted with water before application. This product kills many outdoor pests which damage lawns, roses, flowers, trees, shrubs, vegetables and fruits and insects which annoy pets and people.

) SEE THE BOOKLET FOR COMPLETE DIRECTIONS FOR USE AND PESTS CONTROLLED.

# DIAZINON-25

## INSECT CONTROL

This product must not be used on golf courses and sod farms.

**ACTIVE INGREDIENTS:**

Diazinon: O, O-Diethyl O(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate ..... 25.0%

INERT INGREDIENTS\*: ..... 75.0%

TOTAL: ..... 100.0%

\*Contains aromatic petroleum distillates.

EPA Reg. No. 58866  
EPA Est. 58866-CA-1

For outdoor use on the home lawn and the home garden only.  
Spray concentrate. Contains diazinon insecticide.

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

See side panel for additional precautionary statements.

**PROTECTS:** Lawns, Vegetables, Fruits and Nuts, Roses, Flowers, Trees, Shrubs.

**CONTROLS:** Ants, Aphids, Chinch Bugs, Cutworms, Fire Ants<sup>1</sup>, Flea Beetles, Grubs, Mites, Sod Webworms, and many others.

<sup>1</sup> Aids in control.

**NET CONTENTS: 8 oz.**

**ProGuard, Inc.**  
**Suisun, CA 94585**

Peppers. Tomatoes	Aphids. Colorado Potato Beetles. Fall Armyworms (Tomatoes). Flea Beetles. Leafhoppers. Southern Armyworms (Tomatoes). Vinegar Flies - <i>Drosophila</i> spp. (Tomatoes)
Onions -Bulb and Green)	Onion Thrips
Radishes. Turnips. Carrots. Parsnips. Potatoes	Aphids. Flea Beetles. Leafhoppers. Dipterous Leafminers. Colorado Potato Beetles (Potatoes). Southern Armyworms (Potatoes)
Sweet Corn	Aphids. Flea Beetles. Grasshoppers. Spider Mites
Melons. Squash (Summer, Winter)	Aphids. Thrips. Spider Mites
Vegetables: Perennial Soil Application	
Beans. Peas (except dried varieties)	Cutworms. Mole Crickets. Wireworms (Snap Beans)
Beets. Red (Table)	Cutworms. Mole Crickets
Cabbage	Cutworms. Root Maggots. Wireworms
Carrots. Sweet Corn	Cutworms. Mole Crickets. Wireworms
Lettuce. Tomatoes	Cutworms. Wireworms
Radishes. Turnips	Cutworms
Fruits & Nuts	
Almonds (California only)	Scale Crawlers
Walnuts (California only)	Aphids. Scale Crawlers. Spider Mites and others*
Apricots. Peaches. Nectarines	Aphids. Apricot Mealybugs (Apricots). Spider Mites. Scale Crawlers
Cherries	Black Cherry Aphids. Cherry Fruitflies. Cherry Rust Mites. Scale Crawlers and others*
Plums Prunes	Aphids. Spider Mites
Caneberries (CA, OR, WA only) Blackberries. Boysenberries. Dewberries. Loganberries. Raspberries	Aphids. Leafhoppers. Two-Spotted Spider Mites. Thrips. Raspberry Fruitworms
Strawberries	Aphids. Two-Spotted Spider Mites. Strawberry Leafrollers
Grapes	Aphids. Grape Berry Moths. Grape Leafrollers. Leafhoppers
Roses & Flowers	
Azaleas. Camellias. Chrysanthemums. Carnations. Gladioli. Lilac. Rhododendron. Roses	Aphids. Flea Beetles. Dipterous Leafminers. Leafhoppers. Mealybugs. Two-spotted Spider Mites. Thrips. Whiteflies and others*

80714

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.**

**General Information**

Diazinon-25 is an emulsifiable concentrate product which must be diluted with water before application. Refer to the section How to Mix and Apply. This product kills many outdoor insects which damage lawns, roses, flowers, trees, shrubs, vegetables, fruits and nuts, and insects which annoy pets and people. Contains diazinon insecticide which provides positive control of many "hard-to-kill" sucking and chewing insects. Diazinon-25 works fast and applications should be repeated only as directed to maintain control.

**Notes:** (1) USE CONTENTS WITHIN 6 MONTHS AFTER OPENING. SEE STORAGE AND DISPOSAL DIRECTIONS. (2) Do not use in the home. (3) Do not use on ferns, poinsettia, hibiscus, papaya, filix, gardenia or orchids since injury may occur. (4) Use this product on succulent peas and beans only. Do not use on dried pea or bean varieties. (5) Do not apply to food crops grown in greenhouses.

**Precautions:** (1) Wear a long-sleeved shirt and long-legged pants when spraying. (2) Spray with the wind to your back. (3) Do not spray on windy days. (4) If clothing becomes wet from spray during use, remove clothing after spraying, wash affected body areas thoroughly with soap and water, and launder clothing before wearing again.

**How to Mix and Apply**

**Step #1:** Determine size of spraying job. For lawns and vegetables, measure the area to be sprayed. For all other applications, apply as a thorough cover spray. Make applications when insects first appear. Applications should be repeated only as directed to maintain control.

**Step #2:** Use clean sprayer. Carefully measure and mix the amount of Diazinon-25 and water as suggested in the rate chart.

**Step #3:** Spray as directed in rate chart. Thorough coverage is important. Flush sprayer with clean water after each use.

See chart for complete list of use sites, crops, and insects controlled. Use the rate given in the chart for the type of insect you want to control on the site or crop you are spraying.

**Measurement Conversions**

- 1 tablespoon = 3 teaspoons
- 1 fl. oz. = 6 teaspoons or 2 tablespoons
- 5 fl. oz. = 1 cup
- 1 1/2 tablespoons in 3 gals. of water = 1 1/2 teaspoons in 1 gal. of water
- 1 1/2 tablespoons in 3 gals. of water = 2 teaspoons in 1 gal. of water

**Lawn Pest Control**

**Precautions:** Birds, especially waterfowl, feeding or on tending on treated areas may be killed. Apply this product only as specified on this label. Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of sprays. Where irrigation (watering) is recommended in the directions, or if waterfowl, i.e., ducks or geese, can be expected in the treated area after treatment (except spot treatment), water lawn with at least 1/2 inch of water immediately after spraying this product; however, stop watering before puddling occurs.

Use on Bent, Bermuda, Bluegrass, Dichondra, Fescue, Merion and St. Augustine grass.

To control pests listed below, apply recommended rates.

Pests	Rate		Remarks
	Diazinon	Water	
Lawn Chinch Bugs	1 1/2 tablespoons per 125 sq. ft.	3 gals.	Thoroughly wet down grass a few hours before applying.  In the North, if Chinch Bugs are the only problem, a lower rate of application can be used: 3 teaspoons in 3 gals. of water covers 125 sq. ft. or 4 fl. oz. in 24 gals. of water per 1,000 sq. ft.
	6 fl. oz. per 1,000 sq. ft.	24 gals.	
Ants, Armyworms, Bermudagrass, Mites, Clover Mites, Springtails (Collembola), Crickets, Cutworms, Digger Wasps, Earwigs, Frit Flies, Lawn Billbugs, Leafhoppers, Sod Webworms (Lawn Moths), Sowbugs (Pillbugs)	1 1/2 tablespoons per 125 sq. ft.	3 gals.	Thoroughly wet down grass a few hours before applying.  For ants and digger wasps, spot spray ant hills and wasp ground nest openings. For digger wasps, apply at dusk, when wasps are not active.  Billbug control: Apply when activity is first observed or when chewed or brown grass is present indicating damage from the insect.  For frit flies, mow grass and water thoroughly before treatment, but delay application until grass is dry.
	6 fl. oz. per 1,000 sq. ft.	24 gals.	
Brown Dog Ticks, Chiggers, Fleas	1 1/2 tablespoons per 125 sq. ft.	3 gals.	<b>NOTE:</b> Do not apply this product to animals. For brown dog ticks, spray grass and under shrubbery, particularly near house.
	6 fl. oz. per 1,000 sq. ft.	24 gals.	
Hyperodes Weevil	1 1/2 tablespoons per 125 sq. ft.	3 gals.	Treat problem areas in mid-April and again in mid-May. Water grass thoroughly after application.
	5 fl. oz. per 1,000 sq. ft.	24 gals.	
Milkweeds	1 1/2 tablespoons per 125 sq. ft.	3 gals.	Thoroughly wet grass a few hours before applying.
	6 fl. oz. per 1,000 sq. ft.	24 gals.	
Rhodesgrass Scales	1 1/2 tablespoons per 125 sq. ft.	3 gals.	Apply when crawlers first emerge.
	6 fl. oz. per 1,000 sq. ft.	24 gals.	
White Grubs of Japanese Beetle, European Chaler, Southern Chaler	1 1/2 tablespoons per 125 sq. ft.	3 gals.	Apply any time between late July and early October. Water grass thoroughly after application.
	6 fl. oz. per 1,000 sq. ft.	24 gals.	



90714

Almonds	Aphids, Apricot Mealybugs, Mites (Brown, Two-spotted Spider), Scale Crawlers (Olive, San Jose)	4 teaspoons in 2 gals. of water	Apply as a full cover spray. Do not apply more than 4 gallons per 500 sq. ft. of orchard. Do not make more than 3 applica- tions per season a minimum of 14 days apart. Do not apply within 21 days of harvest.	Plums and Prunes	Aphids (Leaftunt Plum, Mealy Plum, Thistle), Mites (Brown, European Red)	4 teaspoons in 2 gals. of water	Apply as a full cover spray. Do not apply more than 4 gallons per 500 sq. ft. of orchard. Do not make more than 3 applica- tions per season a minimum of 7 days apart. Do not apply within 21 days of harvest.
Cane- berries (CA, OR, WA only): Black- berries, Boysen- berries, Dew- berries, Logan- berries, Rasp- berries	Aphids, Leafhoppers, Raspberry Fruitworms, Two-spotted Spider Mites, Thrips	2 tablespoons in 3 gals. of water	Apply as a full cover spray. Do not apply more than 5 gallons per 500 sq. ft. of field. Do not make more than 5 applications per season a minimum of 14 days apart. Do not apply within 7 days of harvest.  This product may be used on caneberrries only in CA, OR and WA. Do not use this product on caneberr- ries grown in other states.	Straw- berries	Aphids, Two-spotted Spider Mites, Strawberry Leafrollers	2 tablespoons in 3 gals. of water	Apply as a full cover spray. Do not apply more than 3 gallons per 500 sq. ft. of field. Do not make more than 3 applications per season a minimum of 7 days apart. Do not apply within 5 days of harvest.
Cherries	Black Cherry Aphids, Cherry Fruitflies, Cherry Fruitworms (Maggois), Fruitree Leafrollers, Mites (Cherry Rust), Scale Crawlers (San Jose)	4 teaspoons in 2 gals. of water	Apply as a full cover spray. Do not apply more than 4 gallons per 500 sq. ft. of orchard. Do not make more than 3 applica- tions per season a minimum of 7 days apart. Do not apply within 21 days of harvest.	Walnuts (CA only)	Aphids, Scale Crawlers (Walnut, Italian Pear, Fristed), Spider Mites, Twig Borers	2 tablespoons in 3 gals. of water	Apply as a full cover spray. Do not apply more than 6 gallons per 500 sq. ft. of orchard. Do not make more than 3 applica- tions per season 14 days apart. Do not apply after husks open or split. Do not apply within 45 days of harvest.  This product may be used on walnuts only in CA. Do not use this product on walnuts grown in other states.
Grapes	Aphids, Leafhoppers, Grape Berry Moths Grape Leafrollers	4 1/2 teaspoons in 3 gals. of water	Apply as a full cover spray. Do not apply more than 3 gallons per 500 sq. ft. of vineyard. Do not make more than 5 applica- tions per season a minimum of 7 days apart. Do not apply within 28 days of harvest.				
Nectarines	Aphids, Mites (Brown, Two-spotted Spider), Scale Crawlers (Olive, San Jose)	4 teaspoons in 2 gals. of water	Apply as a full cover spray. Do not apply more than 4 gallons per 500 sq. ft. of orchard. Do not make more than 3 applica- tions per season a minimum of 14 days apart. Do not apply within 21 days of harvest.				
Peaches	Aphids, Mites (Brown, Two-spotted Spider), Scale Crawlers (Olive, San Jose)	4 teaspoons in 2 gals. of water	Apply as a full cover spray. Do not apply more than 4 gallons per 500 sq. ft. of orchard. Do not make more than 3 applica- tions per season a minimum of 14 days apart. Do not apply within 21 days of harvest.				

100/14

Nuisance Pests Outside the Home

**Ants, Boxelder Bugs, Brown Dog Ticks, Chiggers, Clover Mites, Cockroaches, Crickets, Earwigs, Fleas, Flies, Millipedes, Sowbugs (Pillbugs), Spiders, Springtails (Collembola)**

Mix 6 fl. oz. in 3 gals. of water and thoroughly spray window and door screens, sills, foundations, porch and patio walls, entranceways, walls, garbage cans, tree trunks, and into cracks and other places where insects can hide.

To prevent entry into houses, spray a five-foot band of soil around the house next to the foundation and spray the foundation of the house to a height of 2-3 feet. Repeat application as necessary.

Do not treat animals with this product. Do not treat plants (other than tree trunks) at this nuisance pest rate.

**Fire Ants**

To aid in the control of fire ants in home lawns and other outdoor home recreation areas, apply 2 tablespoons in a minimum of 1 gal. of water as a drench. Apply over and 6 inches around each mound. Apply gently to avoid disturbing ants. Reapply as necessary. Larger quantities of solution can be prepared for multiple mound treatments. Use equipment capable of delivering the diluted product as a gentle rain at the rate of 1 gal. per mound. High pressure sprays may disturb the ants and cause migration, reducing product effectiveness.

NOTE: For best results, apply in cool weather 55-80°F. or in early morning or late evening hours. Treat new mounds as they appear.

Insect Control on Ornamental Trees, Shrubs and Flowers

Tree and Shrubs	Insects	Rate per 3 Gals. of Water	Timing/Special Directions
Arbormae, Azaleas, Birch, Boxwood, Camellias, Douglas Fir, Elm, Hawthorn, Holly, Juniper, Lilac, Locust, Madia, Mimosa, Oak, Pine, Ornamental Plum, Potted, Rhododendron, Spruce, Willow	Aphids, Bagworms, European Pine Shoot Moths, Holly Bud Moths, Dopterous Leafminers, Lepidoptera, Azalea, Birch, Boxwood, Holly, Oak, Leafhoppers, Mealybugs, Mites (Clover, Cyclamen, European Red, Privet, Two-spotted Spider, Obscure Root Weevils, Omnivorous Leaf-tiers, Scale Crawlers (Cottony-cushion), Euonymus, Lecanium, Pine Needle, Pit, San Jose, Soft)	2 table- spoons	Apply as a thorough foliar spray. Wet leaves and branches to the drooping point. Try to penetrate dense foliage with spray.
	Webworms (Juniper, Mimosa), Whiteflies	2 table- spoons	Apply as a thorough foliar spray. Mimosa Webworms: apply as a thorough foliar spray when adults first appear. Make two additional applications at 4-5 week intervals.
	Apple-and-Thorn Skeletonizers, Hemlock Chermes, Oak Looper, Obliquebanded Leafhopper, Pear Slugs, Tent Caterpillars, Webworms (Cotton-eater, Fall)	2 table- spoons	Apply as a thorough foliar spray.

Roses and Flowers	Insects	Rate per 3 Gals. of Water	Timing/Special Directions
Azaleas, Camellias, Chrysanthemums, Carnations, Gladiol, Lilac, Rhododendron, Roses	Aphids, Euonymus Scale Crawlers, Flea Beetles, Dipterous Leafminers, Leafhoppers, Mealybugs, Mites (Carnation Bud, Carnation Shoot, Cyclamen, European Red, Two-spotted Spider, Thrips, Whiteflies	2 table- spoons	Apply as a thorough foliar spray. Try to spray underside of leaves and penetrate dense foliage.

Fruits and Nuts

Start spraying after blooming. Applications should be made according to infestation of when aphids, mites, and crawler stages of scale insects are first noted.

Observe stated time intervals between last application and harvest of the crop, as well as intervals between applications and total number of applications that can be made per season.

Crop	PHI*
Almonds	**
Apricots	21
Blackberries	7
Boysenberries	7
Cherries	21
Dewberries	7
Grapes	28
Loganberries	7
Nectarines	21
Peaches	21
Plums	21
Prunes	21
Raspberries	
Strawberries	6
Walnuts	15

\* PHI = Pre-harvest interval, or the number of days between the last application and harvest.

\*\* Dormant spray only.

Note: To protect bees, do not apply to fruit trees when trees or substantial numbers of weeds around the trees are in bloom.

Crop	Pest	Rate	Timing/Special Directions
Almonds (CA only)	San Jose scale, Paratons scale	2 tablespoons in 3 gals. of water	Apply only as a dormant spray. Do not apply more than 6 gallons per 500 sq. ft. of orchard.  This product may be used on almonds only in CA. Do not use this product or other as shown in other states.

11/14

**DIAZINON-25**

**Vegetables**

For foliar or soil applications, Diazinon-25 should be applied in the volume of water recommended in the table below to assure thorough coverage of the foliage or soil.

Observe stated time intervals between last application and harvest as well as intervals between applications and maximum numbers of applications per season. Do not apply to leafy crops grown in greenhouses.

**Pre-Harvest Interval (PHI)\* for Foliar Applications**

Crop	PHI* (Days)
Beans <sup>1</sup> (succulent varieties only: Lima, Pole and Snap)	7
Broccoli	7
Cabbage	21
Cauliflower	3
Carrots	14
Cassava	3
Cauliflower	7
Chinese Broccoli, Chinese Cabbage, Collards	10
Cranberries	3
Cucumbers	7
Endive (Escarole)	14
Nonpareil Melons	3
Kale	10
Lettuce	14
Muskmelons	3
Onions (Bulb and Green), Parsnips	14
Peas <sup>1</sup>	7
Peppers	5
Potatoes (Irish)	35
Raspberries, Spinach	14
Squash, Summer	7
Squash, Winter	3
Sweet Corn	7
Tomatoes	1
Turnips (Roots and Tops)	14
Watermelons	3

\* PHI = Pre-harvest interval; or the number of days between the last application and harvest.

<sup>1</sup> Do not use this product on ched peas or bean varieties such as Ohio beans, dried limas, soy peas or Ohio black-eyed peas.

**Vegetables**

For application to the foliage, thoroughly spray all plant parts. Application to the soil should be mixed into the soil immediately after application.

Crop	Pest	Rate	Timing/Special Directions
Beans, Lima, Pole and Snap (succulent varieties only)	Aphids, Cucumber beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 5 days up to 3 applications per season. Do not apply within 7 days of harvest.
	Cutworms (surface, subterranean)	5 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil 1-2 days before planting. Immediately work into soil 1-2 inches.
	Wireworms (snap beans only)	5 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 4-8 inches.
NOTES: (1) Do not apply this product to dried bean varieties such as pinto beans or dried limas. (2) To protect beans, do not apply bean foliage treatments if the beans or weeds in the treatment area are in bloom.			
Beans, Red (Table)	Cutworms (surface, subterranean)	5 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil 1-2 days before planting. Immediately work into soil 1-2 inches.
Broccoli	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Diterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days of harvest.  NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
	Cutworms (surface, subterranean)	5 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
Cabbage	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Diterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 21 days of harvest.  NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
	Cutworms (surface, subterranean)	5 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Root Maggots	5 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches.
	Wireworms	5 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 4-8 inches.

**BEST AVAILABLE COPY**

DIAZINON-25

12/14

Carrots	Aphids. Flea Beetles. Leafhoppers. Dipterous Leafminers. Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 14 days of harvest.  NOTE: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecti- cides.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for sur- face cutworms, 3-6 inches for subterra- nean cutworms.
	More Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil 1-2 days before plant- ing. Immediately work into soil 1-2 inches.
	Worms	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 4-8 inches
Cauliflower	Aphids. Diamondback Moths Flea Beetles. Imported Cabbageworms. Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 7 days of harvest.  NOTE: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecti- cides.
	Chinese Broccoli	Aphids. Diamondback Moths Flea Beetles. Imported Cabbageworms Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden  Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 10 days of harvest.  NOTE: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecti- cides.
Chinese Cabbage (Bok Choy and Napa)	Aphids. Diamondback Moths. Flea Beetles. Imported Cabbageworms Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 10 days of harvest.  NOTE: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecti- cides.
	Collards	Aphids. Diamondback Moths. Flea Beetles. Imported Cabbageworms Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden  Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 10 days of harvest.  NOTE: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecti- cides.

Corn (see sweet corn)				
Cucumbers	Aphids. Thrips. Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 7 days of harvest.	
	Escove (Escarole)	Aphids. Flea Beetles. Leafhoppers. Dipterous Leafminers. Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden  Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 14 days of harvest.  NOTE: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecti- cides.	
Kale	Aphids. Diamondback Moths. Flea Beetles. Imported Cabbageworms. Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 10 days of harvest.  NOTE: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecti- cides.	
	Lettuce	Aphids. Flea Beetles. Leafhoppers. Dipterous Leafminers. Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 14 days of harvest.  NOTE: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecti- cides.
		Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
		Worms	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 4-8 inches.
Melons (Cantaloupe, Casabas, Cren- snaws, Honey- dews, Musk- melons, Persians, and Hydros of these)	Aphids. Thrips. Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 3 days of harvest.	
	Water- melons			

BEST AVAILABLE COPY

Onions (Bulb and Green)	Onion Thrips	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest.
Parsnips	Aphids. Flea Beetles. Leafhoppers. Dipterous Leafminers. Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat applications every 7 days up to 5 applications per season. Do not apply within 14 days of harvest.  NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Peas (succulent only)	Aphids. Cucumber Beetles. Grasshoppers. Leafhoppers. Mexican Bean Beetles. Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 5 days up to 3 applications per season. Allow at least 7 days between the last application and harvest.
	Cutworms (surface, subterranean)	5 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil 1-2 days before planting. Immediately work into soil 1-2 inches.
	Wireworms	5 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 4-8 inches.
NOTES. (1) Do not apply this product to dried pea varieties such as split peas or dried black-eyed peas. (2) To protect bees, do not apply foliage treatments if bees or weeds in the treatment area are in bloom.			
Peppers	Aphids. Colorado Potato Beetles. Flea Beetles. Leafhoppers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 5 days of harvest.
Potatoes	Aphids. Flea Beetles. Leafhoppers. Colorado Potato Beetles. Southern Armyworms. Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 35 days of harvest.  NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.

Radishes	Aphids. Flea Beetles. Leafhoppers. Imported Cabbageworms. Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 3 applications per season. Do not apply within 14 days of harvest.  NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
Squash	Aphids. Flea Beetles. Dipterous Leafminers. Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest.  NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Squash, Summer	Aphids. Thrips. Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days of harvest.
Squash, Winter	Aphids. Thrips. Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 3 days of harvest.
Sweet Corn	Aphids, Thrips. Flea Beetles. Grasshoppers. Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Corn may be picked 7 days following last application.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil 1-2 days before planting. Immediately work into soil 1-2 inches.
	Wireworms	5 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 4-8 inches.
NOTE: To protect bees, do not apply to corn foliage during the pollen shed period.			

BEST AVAILABLE COPY

Tomatoes	Aphids. Colorado Potato Beetles. Fall Armyworms. Flea Beetles. Leafhoppers. Southern Armyworms. Vinegar Flies -Drosophila spp.	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 1 day of harvest.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 4-8 inches.
Turnips and Turnip Tops	Aphids. Flea Beetles. Leafhoppers. Imported Cabbageworms Destructive Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest.  NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.

### Storage and Disposal

#### Storage

Do not store near heat or open flame. Store in original container in an area inaccessible to children and pets. USE CONTENTS WITHIN 6 MONTHS AFTER OPENING.

#### Disposal

Do not reuse empty container. Thoroughly rinse container, securely wrap it in several layers of newspaper and discard in trash. If disposal of a partially full container is necessary, make sure cap is tightly closed, securely wrap the container in several layers of newspaper and discard in trash.

### Precautionary Statements

#### Hazards to Humans and Domestic Animals

##### CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Causes moderate eye injury. Avoid breathing spray mist. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Do not use inside the home. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not allow children or pets to go onto a treated area until the treated area, such as grass or soil, has dried. Do not contaminate ornamental fish ponds. Food utensils such as spoons and measuring cups must not be used for food purposes after use in measuring pesticides.

#### Statement of Practical Treatment

If swallowed: Call a physician immediately. DO NOT INDUCE VOMITING UNLESS NO OTHER TREATMENT IS AVAILABLE. Vomiting may cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing.

If inhaled: Remove victim to fresh air and apply artificial respiration if indicated. Get medical attention.

If on skin: Wash promptly with soap and water. Rinse thoroughly. Get medical attention.

If in eyes: Rinse eyes with plenty of water and call a physician immediately.

Note to Physician: This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. Z-PAM is also antidotal and may be administered, but only in conjunction with atropine. This product contains aromatic petroleum distillates which may present an aspiration hazard. Gastric lavage may be indicated if product was taken internally.

#### Environmental Hazards

This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. A. C. C. overlapping of sprays. For lawns, where irrigation (watering) is recommended in the directions for waterfowl, i.e., ducks or geese, can be expected in the treated area after treatment (except spot treatment), water lawn with at least 1/2 inch of water immediately after spraying this product, however, stop watering before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

This pesticide is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this pesticide or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

#### Physical or Chemical Hazards

Do not use, pour, spill, or store near heat or open flame.

PROGUARD, INC.  
Suisun, CA 94585

**BEST AVAILABLE COPY**