

AS-130

ACCEPTED
with COMMENTS
in EPA Letter Dated:

SEP 22 1988

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

588 19-3

ACTIVE INGREDIENT:

Poly(oxyethylene (dimethyliminio) ethylene-
(dimethyliminio) ethylene dichloride) 30.0 %

INERT INGREDIENT 70.0 %

This product contains 2.67 lb. of active ingredient per gallon and weighs
8.9 lb. per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid breathing vapors. Avoid contact with skin, eyes, or clothing.

FIRST AID: If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Keep out of lakes, streams, or ponds. Permits may be required for discharges containing this pesticide into lakes, streams, ponds, or public water. For guidance, contact the regional office of the Environmental Protection Agency.

DIRECTIONS

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

AS-130
and fungi in recirculating commercial and industrial air wash systems. Prior to its use, systems must be cleaned to remove slime, and other deposits. An initial slug of AS-130 fluid ounces of AS-130

of water to provide a concentration of 1 AS-130

in the system, is recommended. Repeat initial AS-130 Subsequent slug additions of 0.47 AS-130

per AS-130
40 parts per million of AS-130

employed every 2 to 5 days, or as needed upon the relative amount of bleedoff and the severity of the problem. Slug additions should be made in the AS-130

is for air-washing systems that maintain effectiveness to its use, systems should be cleaned to remove deposits. An initial slug dose of 7.1 AS-130

is recommended. Repeat initial dosage until concentrations of 4.8 to 11.9 fluid ounces per 1000 gallons of water should be employed. Frequency of addition depends upon the relative severity of the bacterial problem. Slug additions may be made in collection trays of the airwash system.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. **STORAGE:** Do not stack more than four drums high. Leaking drums should be disposed of. Spills should be absorbed in sawdust or other material and the container closed when not in use.

PESTICIDE DISPOSAL: Wastes resulting from the use of this pesticide should be disposed of at a government approved waste disposal facility.

CONTAINER DISPOSAL

PLASTIC: Triple rinse (or equivalent). Then offer for recycling, if available, or for sanitary landfill, by incineration, or, if allowed by state and local authorities, by other procedures approved by state and local authorities.

METAL: Triple rinse (or equivalent). Then offer for recycling, if available, or for sanitary landfill, or by other procedures approved by state and local authorities.

Manufacturer

A.S., INC.

138 SKELLY

HERCULES, CA 94547

EPA REG. NO. _____

NET CONTENTS _____