US ENVIRONMENTAL PROTECTION AGENCY MOFFICE OF PESTICIDES PROGRAMS. REGISTRATION DIVISION (TS-767) WASHINGTON, DC 20460

 \vdash

L

NOTICE OF PESTICIDE: REGISTRATION

(Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)

EPA REGISTRATION NO. 59144-3 TERM OF ISSUANCE

NAME OF PESTICIDE PRODUC

5% Sevin Brand Carbaryl Insecticide

NAME AND ADDRESS OF REGISTRANT (.nclude ZIP code)

d.R. Mallane, The. 7316 9/3 E 34, 53150 2127 diami, FI 33143

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Tederal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

ing the first of the major of the first of the second and the second second second second second second second estion 3(a) to there we although the

- to Surgery to the Long Pathological Con-The Contrast Convents of Strat. St. 1 your product units (IPP) is only in 3(min.) where the Amency requires all remistrants of samilar projuments so not such data.
- 2. Harry the Carl Dear Harry Officer action before you release th product for the fruit.
 - Add the phrase "TEA Redistration to. 59144-3."
 - b. Add the ichingon to the paragrams for flew control does and Cats:

to hid use it program dogs.

o. Transferthe claim to be water, of your Burase ticks. There are no lata for a last formulation to support this claim. Yes ave to mainst of school data of the want to append other claim on your label.

SIGNATURE OF APPROVING OFFICIAL ennis A. Edward

EPA Form 8570-6 (Rev. 5-76)

PREVIOUS EDITION MAY BE USED UNTIL SUPPLY IS EXHAUS

- 3. A 1-year storage stability is needed for this product. The formulated product must be analyzed for its active ingredient at time zero and during a year of storage. The storage should be in warehouse conditions of temperature and humidity and stored in similar containers as you will be using in the trade. For the storage stability you cannot reference the concentrate you are using to formulate your product. Submit a storage stability study within 15 months from the date of this Registration Notice.
- 4. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the λ -79 enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with PIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Dennis H. Edwards, Jr. Product Manager (12) Insecticide-Rodenticide Branch Registration Division (H7505C)

Enclosures

15

5% SEVIN Brand Carbaryl Insecticide DUST

CONTROLS INSECTS ON VEGETABLES. LAWNS, ORNAMENTALS AND FRUITS

FOR FLEAS ON DOGS AND CATS **CONTROLS LYME DISEASE TICKS**

ACTIVE INGREDIENT:

Carbaryl:1-Napthyl N-methylcarbamate - -5.0% INERT INGREDIENTS: - - - - -95.0% Total 100.0%

SEVIN is a registered Trademark of Phone Poulenc for Carbaryl Insecticides.

KEEP OUT OF REACH OF CHILDREN CAUTION

"PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicac'o ampliamente" STATEMENTS OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician of Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF IN EYES: Flush eyes with plenty of water. Get medical attention if

irritation persists.
#F ON SKIN: Remove contaminated clothing and wash affected area with soap and water. Get medical attention if irritation persists. F <u>INHALED: Remove victim to fresh air. Apply Artificial Respiration if</u> IF INHALED: Hemove victim to maintained. Get medical attention.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION: May be harmful if swallowed. Do not breathe dust. Avoid contact with eyes, skir. and clothing. Wash thoroughly after handling. Change contaminated clothing daily. Avoid contamination of feed and foodstuffs. NOTE TO PHYSICIAN Carbarylife a moderate reversible cholinesterase inhibitor. Atraping is agricotal.

SEE SIDE/BACK PANEL FOR ADOMONAL PSECAN BRIVARY STATEMENTS EPA Reg. No. 59144-EPA Est. No. 59144-GA-1

> Net-Weight 4 Pound GRO TEC; INC; P.O. BOX 290 MADISON, GA 30650

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic invertebrates. Do not apply directly to water or wetlands. Do not contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the proposed treatment areas.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. STORAGE: Store product in original container in a cool, dry, locked place out of reach of children. DISPOSAL: Do not re-use empty container, Rinse thoroughly before discarding in trash.

READ ENTIRE LABEL BEFORE USING

5% SEVIN Brand Carbaryl Insecticide DUST may be applied with any suitable dust applicator. For best results make application in the morning or evening when the air is still. Be sure to treat both upper and lower surfaces of leaves. Apply (down wind) to give thorough coverage but avoid applying in excess. Applications should be started when insects first appear. Repeat as necessary. Apply at 7 to 10 day intervals, or 3 to 5 day intervals in rainly weather. Listed below are the crops with the pests controlled followed by the time interval to stop applications prior to harvest. As a general rule apply enough dust to give thorough coverage to all plant parts or 11 to 15 ounces per 1,000 Sq.Ft. (equal to 30 to 40 lbs. dust per acre.)

VEGETABLES and FRUIT

ASPARAGUS: Asparagus Beetles. Do not apply later than 1 day before harvest. Do not apply more often than once every 3 days. BEANS: Mexican Bean Beetle, Bean Leaf Beetle, Leafhoppers, Flea Beetles, Com Earworms, Lygus Bugs, Stink Bugs, Lima Bean Pod Borer, Japanese Beetle. Can be applied day of harvest. BEETS (table): Armyworms, Flea Beetles, Leafhoppers, 3 days before harvest. 14 days before harvest if tope are used for food or feed.

BRUSSELS SPROUTS, BROCCOL, CABBAGE, CALUFLOWER:
Small Cabbage Looper, (treat every 5 to 7 days for control),
Armyworms, Flea Beetise. Do not apply later 3 days before harvest.

CARROTS: Leafhoppers, Flea Beetles. Can be applied day of harvest. CITHUS: Black Scale, California Red Scale, Soft Brown Scale, Orange Tortrot, California Orange Dog, Fruit Tree Leaf Rollers. For Scale Insects best results are obtained when crawlers are present. Apply 5 days before harvest.

COLLARDS, KALE, MUSTARD GREENS, RADISHES, RUTABAGAS, CHINESE CABBAGE, HORSERADISH and TURNIPS: Flea Boetles, Harlequid Bug, Leathoppers, Armyworm, Imported Cabbageworm, Tarnished Plant Bug and Stink Bugs, (3 days of harvest on horseradish, radishes, rutabagas and turnip roots; 14 days on chinese cabbage, collards, kale, mustard greens and turnip tops).

CORN(Sweet): Corn Earworm, Flea Beetles, Leafhoppers, Fall Armyworms. For Corn Earworm control treat corn silks every 3 days

until silke dry. Can apply day of harvest.

<u>CUCUMBERS</u>, <u>MELONS</u>, <u>SQUASH</u>, <u>PUMPKINS</u> <u>CANTALOUPES</u>:

Squash Bugs, Leafhoppers, Cucumber Beetles, Flea Beetles. Do not apply to Melons or Cucumbers when temperatures exceed 80° F or

when vines are wet. Can be applied day of harvest.

<u>EGGPLANT</u>: Corn Earworm, Flea Bestles, Hornworms, Cucumber Beetles. Can be applied day of harvest.

LETTUCE: Armyworms, Small Cabbage Looper (treat every 5 to 7 days for control), Leafhoppers, Salt Marsh Caterpillars. For Head Lettuce; 3 days before harvest. Leaf Lettuce; 14 days before harvest.

PEPPERS: Corn Ean applied up to day of he POTATOES: (White); harvest. TOMATOES: Tomato Stink Bug, Leafhoppers STRAWBERRIES: Leaf

One day before harves

OR **ORNAMENTAL**

Apply this product to th the following pests. Re Apple Aphid Bagworm **Blister Beetles** Boxelder Bug Elm Leaf Beetle Flea Beetle Japanese Beelte June Beetle Leathoppers For best scale insect co present. Do not apply

TURF & P

Apply 8 oz. to cover 25 area to control Earwig: and Ants. Area should dust off foliage. Reper treated areas until gras

FLEAS

For control of Fleas, at spoonful into pets skin. quarters. Replace with area where insects are pet several hours after t old. Use lower rate and

LYME

Home Lawns a Areas, Parks, Par

Càmp Grou For control of bigides sp Tick) and Amblyomma Disease apply 3 1/2 lb spring or early summer ticks apply in late sumr entire lawn and perime where exposure to tick and do not allow reent Ticks may be reintredu Retreat as necessary t contaminate water. NOTICE: This product reasonably fit for the

accordance with directi turer le not responsible instructions, or under a reasonably foreseeable buyer assumes the risk