


**GREEN CHARM®**

**SEVIN® 5%**

brand carbaryl insecticide

**DUST**

**Controls Insects on Vegetables, Lawns,  
Ornamentals and Fruits  
For Fleas on Dogs and Cats**



**ACTIVE INGREDIENT:**

Carbaryl; 1 Naphthyl N-methylcarbamate - - - 5.0%

**INERT INGREDIENTS:** - - - 95.0%

Total 100.0%

SEVIN® is a Registered Trademark of Rhone-Poulenc for Carbaryl Insecticides

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

**PRECAUCION AL USUARIO: Si usted no lee ingles, no use este  
producto hasta que la etiqueta haya sido explicado completamente.  
See Back Panel for Additional Precautionary Statements**

EPA Reg. No. 59144-3

EPA EST. NO. 59144-GA-1; 6728-FL-1; 61936-GA-1; 788-GA-1; 44818-MO-1; 19713-GA-1  
Subscript is first letter of run code on container (59-30eq.prm4(12-15-92))

**NET WEIGHT 4 POUNDS**

**PROUDLY**  
P.O. BOX 290 - Madison, GA 30650

# GREEN CHARM® SEVIN® 5% brand carbaryl insecticide DUST

EPA Reg. No. 59144-3

NET WEIGHT 4 POUNDS

## STATEMENTS OF PRACTICAL TREATMENT

**IF SWALLOWED:** Call a physician of Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person. **IF IN EYES:** Flush eyes with plenty of water. Get medical attention if irritation persists. **IF ON SKIN:** Remove contaminated clothing and wash affected area with soap and water. Get medical attention if irritation persists. **IF INHALED:** Remove victim to fresh air. Apply Artificial Respiration if indicated. Get medical attention.

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed. Do not breathe dust. Avoid contact with eyes, skin and clothing. Wash thoroughly after handling. Change contaminated clothing daily. Avoid contamination of feed and foodstuffs. **NOTE TO PHYSICIAN:** Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal.

### ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic invertebrates. Do not apply directly to water or wetlands. Do not contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the proposed treatment areas.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store product in original container in a cool, dry, locked place out of reach of children. **DISPOSAL:** Do not re-use empty container. Rinse thoroughly before discarding in trash.

## READ ENTIRE LABEL BEFORE USING

GREEN CHARM SEVIN 5% Brand Carbaryl Insecticide DUST may be applied with any suitable dust applicator. For best results make application in the morning or evening when the air is still. Be sure to treat both upper and lower surfaces of leaves. Apply (down wind) to give thorough coverage but avoid applying in excess. Applications should be started when insects first appear. Repeat as necessary. Apply at 7 to 10 day intervals, or 3 to 5 day intervals in rainy weather. Listed below are the crops with the pests controlled followed by the time interval to stop applications prior to harvest. As a general rule apply enough dust to give thorough coverage to all plant parts or 11 to 15 ounces per 1,000 sq. ft. (equal to 40 lbs. dust per acre.)

## TURF PERIMETER INSECTS

Apply 1/2 oz. to cover 250 sq. ft. of turf or around building foundation area to control Earwigs - Millipedes - Sod Webworms - Leafhoppers and Ants. Area should be lightly sprinkled after application to wash dust off foliage. Repeat as necessary. Keep children and pets off treated areas until grass has been sprinkled and area is dry.

## FLEAS ON DOGS & CATS

For control of Fleas, and Brown Dog Ticks, rub 1/2 to 1 level tablespoonful into pets skin. After treating, bathing and other infested quarters. Replace with fresh bedding. Treat entire pet especially in area where insects are seen. Avoid getting into pets eyes. Groom the pet several hours after treatment. Do not treat kittens under 2 months old. Use low rate and apply sparingly for cats. Repeat as needed. Do not use on pregnant dogs.

**WARRANTY AND DISCLAIMER:** Subject to the limitations hereinafter set forth, Green Charm, Inc. warrants that the product sold is as described on the container, within recognized tolerances. Liability on this warranty is limited to the purchase price of the product. In no event shall Green Charm, Inc. be liable for indirect, consequential, or other damages in excess of the purchase price. Green Charm, Inc. makes no other warranty either expressed or implied as to merchantability or fitness for a particular purpose of the product. The product is sold at the price quoted only upon the condition that the above disclaimer of warranty and the limitation of damages are part of the basis of the bargain. Products not accepted under 14-day return should be returned within 30 days in the original, unopened container and buyer will receive his or her purchase price or a price quotation which includes shipping charges as desired on the basis of the terms provided each quote is obtainable which is determined upon the nature of the products sold. [EPA-36AG2 PM4 (03-01-82) except 08-03-8012-15-82.]

## VEGETABLES and FRUIT

**ASPARAGUS:** Asparagus Beetles. Do not apply later than 1 day before harvest. Do not apply more often than once every 3 days.

**BEANS:** Mexican Bean Beetle, Bean Leaf Beetle, Leafhoppers, Flea Beetles, Corn Earworms, Lygus Bugs, Stink Bugs, Lima Bean Pod Borer, Japanese Beetle. Can be applied day of harvest.

**BEETS (table):** Armyworms, Flea Beetles, Leafhoppers. 3 days before harvest. 14 days before harvest if tops are used for food or feed.

**BRUSSELS SPROUTS, BROCCOLI, CABBAGE, CAULIFLOWER:** Small Cabbage Looper, (treat every 5 to 7 days for control), Armyworms, Flea Beetles. Do not apply later than 3 days before harvest.

**CARROTS:** Leafhoppers, Flea Beetles. Can be applied day of harvest.

**CITRUS:** Black Scale, California Red Scale, Soft Brown Scale, Orange Tortrix, California Orange Dog, Fruit Tree Leaf Rollers. For Scale insects best results are obtained when crawling are present. Apply 5 days before harvest.

**COLLARDS, KALE, MUSTARD GREENS, RADISHES, RUTABAGAS, CHINESE CABBAGE, HORSERADISH and TURNIPS:** Flea Beetles, Harlequin Bug, Leafhoppers, Armyworm, Imported Cabbageworm, Tarnished Plant Bug and Stink Bugs, (3 days of harvest on horseradish, radishes, rutabagas and turnip roots; 14 days on Chinese cabbage, collards, kale, mustard greens and turnip tops).

**CORN (Sweet):** Corn Earworm, Flea Beetles, Leafhoppers, Fall Armyworms. For Corn Earworm control treat corn silks every 3 days until silks dry. Can apply day of harvest.

**CUCUMBERS, MELONS, SQUASH, PUMPKINS, CANTALOUPES:** Squash Bugs, Leafhoppers, Cucumber Beetles, Flea Beetles. Do not apply to Melons or Cucumbers when temperatures exceed 80° F or when vines are wet. Can be applied day of harvest.

**EGGPLANT:** Corn Earworm, Flea Beetles, Hornworms, Cucumber Beetles. Can be applied day of harvest.

**LETTUCE:** Armyworms, Small Cabbage Looper (treat every 5 to 7 days for control), Leafhoppers, Salt Marsh Caterpillars. For Head Lettuce; 3 days before harvest. Leaf Lettuce; 14 days before harvest.

**PEPPERS:** Corn Earworm, Leafhoppers, Flea Beetles. Can be applied up to day of harvest.

**POTATOES (White):** Flea Beetles, Leafhoppers. Can apply day of harvest.

**TOMATOES:** Tomato Fruit Worm, Tomato Hornworm, Flea Beetles, Stink Bug, Leafhoppers. Apply day of harvest.

**STRAWBERRIES:** Leaf Rollers, Lygus Bugs, Salt Marsh Caterpillar. One day before harvest.

## ORNAMENTALS

### ORNAMENTAL FLOWERS, SHRUBS AND TREES

Apply this product to thoroughly cover the plant foliage for control of the following pests. Repeat as needed for continued control.

Apple Aphid	Leaf Rollers	Rose Bug
Begonia	Mealy Bugs	SCALE -
Black Beetles	Minnow Weevils	Soft Brown
Colorado Bug	Oak Leaf Miner	European Elm
Elm Leaf Beetle	Orange Tortrix	Cat Kermes
Flea Beetle	Perforated Cuckoo	Oystershell
Japanese Beetle	Plant Bug	Black
June Beetle	Pyralis	Colony Maple
Leafhoppers	Rose Aphids	Thrips (Exposed)

For best scale insect control apply when newly hatched "crawlers" are present. Do not apply to BOSTON IVY as plant injury will result.

**GREEN CHARM**

P.O. BOX 290 • Madison, GA 30650