

2419 59144-5 107

GREEN CHARM[®] SEVIN[®] 10%

brand carbaryl insecticide

DUST

For Control of Insect Pests on
Vegetables, Fruit and
Ornamental Plants

Fire Ant Control



ACCEPTED
FEB 16 1993
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 59144-5

ACTIVE INGREDIENT:

Carbaryl; 1 Naphthyl N-methylcarbamate . . . 10.0%

INERT INGREDIENTS: 90.0%

Total 100.0%

SEVIN[®] is a Registered Trademark of Rhone Poulenc for Carbaryl Insecticides

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este
producto hasta que la etiqueta haya sido explicado ampliamente

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No. 59144-5

EPA EST. NO. 59144-GA-1; 6728-FL-1; 51036 GA-1; 758-GA-1; 44818-MO-1; 19713-GA-1

Subscript is first letter of run code on container (59-58AG.PM4(12-15-92))

NET WEIGHT 4 POUNDS



P.O. BOX 290 - Madison, GA 30650

GREEN CHARM® SEVIN® 10% DUST

EPA Reg. No. 59144-5

Net Weight 4 Pounds

STATEMENTS OF PRACTICAL TREATMENT
IF SWALLOWED: Call a physician of Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person. **IF IN EYES:** Flush eyes with plenty of water. Get medical attention if irritation persists. **IF ON SKIN:** Remove contaminated clothing and wash affected area with soap and water. Get medical attention if irritation persists. **IF INHALED:** Remove victim to fresh air. Apply Artificial Respiration if indicated. Get medical attention.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION: Harmful if swallowed. Do not breathe dust. Avoid contact with eyes, skin and clothing. Wash thoroughly after using. Change contaminated clothing daily. Avoid contamination of feed and foodstuffs. **NOTE TO PHYSICIAN:** Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal.

ENVIRONMENTAL HAZARDS
This product is extremely toxic to aquatic invertebrates. Do not apply directly to water or wetlands. Do not contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the proposed treatment areas.

DIRECTIONS FOR USE
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL
Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store product in original container in a cool, dry, locked place out of reach of children. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

READ ENTIRE LABEL BEFORE USING
GREEN CHARM SEVIN 10% Brand Carbaryl Insecticide DUST can be applied with any suitable hand or power duster. For best results make application in the morning or evening when the air is still. Treat both upper and lower surfaces of leaves. Applications should be started when insects first appear and repeated at 7 to 10 day intervals, or 5 to 7 days in rainy weather. Crops with the pests are listed below. Observe number of days to stop treatment before harvest. Apply to deposit a light visible dusting to all plant and fruit parts or 4 to 8 ounces per 1,000 sq. ft.; which is equal to approximately 10 to 20 lbs./acre. Use the low rate when insects are first seen or are anticipated; the high rate when insects are present in damaging numbers.

VEGETABLES and FRUIT

ASPARAGUS: Asparagus Beetles. Do not apply later than 1 day before harvest. Do not apply more often than once every 3 days.

BEANS: Mexican Bean Beetle, Bean Leaf Beetle, Leafhoppers, Flea Beetles, Corn Earworms, Lygus Bugs, Stink Bugs, Lima Bean Pod Borer, Japanese Beetle. Can be applied day of harvest.

BEETS (table): Armyworms, Flea Beetles, Leafhoppers. 3 days. 14 days if tops are used for food or feed.

BRUSSELS SPROUTS, BROCCOLI, CABBAGE, CAULIFLOWER: Small Cabbage Looper, (treat every 5 to 7 days for control), Armyworms, Flea Beetles. 3 days before harvest.

CARROTS: Leafhoppers, Flea Beetles. Can be applied day of harvest.

CITRUS: Black Scale, California Red Scale, Soft Brown Scale, Orange Tortrix, California Orange Dog, Fruit Tree Leaf Rollers. For Scale insects best results are obtained when crawlers are present. 3 days before harvest.

COLLARDS, KALE, MUSTARD GREENS, RADISHES, RUTABAGAS, CHINESE CABBAGE, HORRER, CUCUMBER and TURNIPS: Flea Beetles, Harlequin Bug, Leafhoppers, Armyworm,

Imported Cabbageworm, Tarnished Plant Bug and Stink Bugs, (3 days of harvest on horseradish, radishes, rutabagas and turnip roots; 14 days on Chinese cabbage, collards, kale, mustard greens and turnip tops).

CORN (Sweet and Field): Corn Earworm, Flea Beetles, Leafhoppers, Fall Armyworms. For Corn Earworm control treat corn silks every 3 days until silks dry. Can apply day of harvest. Wait 7 days before foraging.

CUCUMBERS, MELONS, SQUASH, PUMPKINS: Squash Bugs, Leafhoppers, Cucumber Beetles, Flea Beetles. Do not apply to Melons or Cucumbers when temperatures exceed 80° F. or when vines are wet. Can be applied day of harvest.

EGGPLANT: Corn Earworm, Flea Beetles, Hornworms, Cucumber Beetles. Can be applied day of harvest.

GRAPES: Grape Leafhopper, Grape Leaf Folder, Grape Berry Moth, European Fruit Lecanium, Grape Leaf Skeletonizer, Japanese Beetles, June Beetles, Red Banded Leaf Roller. Day of Harvest.

HEAD and LEAF LETTUCE: Armyworms, Small Cabbage Looper (treat every 5 to 7 days for control), Leafhoppers, Salt Marsh Caterpillars. 3 days before harvest on Head Lettuce; 14 days of harvest on Leaf Lettuce.

OKRA: Corn Earworm, Flea Beetles, Japanese Beetles, Stink Bugs. May be applied day of harvest.

PEANUTS AND COWPEAS: For Blister Beetles, Mexican Bean Beetles, Thrips, Leafhoppers, Three Corned Alfalfa Hopper, Cucumber Beetles, Bean Leaf Beetles, Japanese Beetles, Alfalfa Caterpillar, Velvetbean Caterpillar, Green Cloverworm, Armyworms, Webworms, Corn Earworms, Stink Bugs, Grasshoppers. On Cowpeas do not apply within 3 days of harvest; peanuts, no limitation.

PEPPERS: Corn Earworm, Leafhoppers, Flea Beetles. Day of harvest.

POTATOES (White): Flea Beetles, Leafhoppers. Day of harvest.

TOMATOES: Tomato Fruit Worm, Tomato Hornworm, Flea Beetles, Stink Bug, Leafhoppers. Apply day of harvest.

STRAWBERRIES: Leaf Rollers, Lygus Bugs, Salt Marsh Caterpillar. One day before harvest.

SUGAR BEETS: Leafhoppers, Armyworms, Webworms, Grasshoppers. 14 day Pre-harvest interval.

ORNAMENTALS

ORNAMENTAL FLOWERS, SHRUBS AND TREES

Apply this product to thoroughly cover the plant foliage for control of the following pests. Repeat as needed for continued control.

Apple Aphid	Minors Webworm	Blister Beetles
Bombardier Bug	Oak Leafminer	- SCALE -
Elm Leaf Beetle	Orange Tortrix	Soft Brown
Flea Beetle	Periodical Cicadas	European Elm
June Beetle	Pyralids	Oak Kermes
Leafhoppers	Rose Aphids	Oystershell
Leaf Rollers	Rose Slug	Black
Mealy Bugs	Thrips (Exposed Stages)	Cottontail Maple

For best scale insect control apply when newly hatched "crawlers" are present. Do not apply to BOSTON IVY as plant injury will result.

TURF & PERIMETER INSECTS

Apply 4 oz. to cover 250 sq. ft. of smaller turf areas adjacent to patios and home recreational areas or foundation perimeter areas to control ANTS, CHIGGERS, EARWIGS, FLEAS, MILLIPEDES, SPIDERS, SOD WEBWORMS, SOWBUGS AND LEAFHOPPERS. Area should be lightly sprinkled after application to wash dust off foliage. Repeat as necessary. Keep children and pets off treated areas until grass has been sprinkled & area is dry.

IMPORTED FIRE ANT CONTROL

Apply 2 oz. (8 level tablespoonfuls) evenly over and around each mound. After application, gently sprinkle the treated area with a minimum of 1 quart of water per 8 inches of mound diameter. For best results; do not disturb the mound before or during the application; apply in cool weather 65-80 degrees or in early morning or late evening. Treat new mounds as they appear.

WARRANTY AND DISCLAIMER: Subject to the limitation of liability herein set forth, Gro-Tec, Inc. warrants that the product sold is as described on the container, within recognized tolerances. Liability on this warranty is limited to the purchase price of the product. In no event shall Gro-Tec, Inc. be liable for indirect, consequential, or other damages in excess of the purchase price. Gro-Tec, Inc. makes no other warranty either expressed or implied as to merchantability or fitness for a particular purpose of the product. The product is sold at the price quoted only upon the condition that the above disclaimer of warranty and the limitation of damages are part of the basis of the sale. Products not covered under these terms should be returned within 30 days in the original, unopened container and Buyer will receive his choice of either refund of purchase price or a price credit which includes such return as described on the back of the package provided such credit is obtainable which is dependent upon the nature of the products sold. (EPA-59144-5 FPM - EPA approved version 07-55-92 of the 05-01-88 Approved copy 12-15-92)

GRO-TEC
P.O. BOX 250 • Madison, GA 30650